

World Journal of *Clinical Cases*

World J Clin Cases 2023 June 26; 11(18): 4210-4457



REVIEW

- 4210 Should gastroenterologists prescribe cannabis? The highs, the lows and the unknowns
Samuel S, Michael M, Tadros M

MINIREVIEWS

- 4231 Application of artificial intelligence in trauma orthopedics: Limitation and prospects
Salimi M, Parry JA, Shahrokhi R, Mosalamiaghili S
- 4241 Weight loss maintenance after bariatric surgery
Cho YH, Lee Y, Choi JI, Lee SR, Lee SY
- 4251 Bicuspid aortic valve with associated aortopathy, significant left ventricular hypertrophy or concomitant hypertrophic cardiomyopathy: A diagnostic and therapeutic challenge
Sopek Merkaš I, Lakušić N, Predrijevac M, Štambuk K, Hrabak Paar M
- 4258 Application experience and research progress of different emerging technologies in plastic surgery
Yang B, Yang L, Huang WL, Zhou QZ, He J, Zhao X

ORIGINAL ARTICLE

Case Control Study

- 4267 Multimodal integrated intervention for children with attention-deficit/hyperactivity disorder
Lv YB, Cheng W, Wang MH, Wang XM, Hu YL, Lv LQ

Retrospective Study

- 4277 Portal vein computed tomography imaging characteristics and their relationship with bleeding risk in patients with liver cirrhosis undergoing interventional therapy
Song XJ, Liu JL, Jia SY, Zhang K

Observational Study

- 4287 Wrist-ankle acupuncture combined with pain nursing for the treatment of urinary calculi with acute pain
Wu LM, Liu Q, Yin XH, Yang LP, Yuan J, Zhang XQ, Wang YL

CASE REPORT

- 4295 Coexistence of diffuse large B-cell lymphoma, acute myeloid leukemia, and untreated lymphoplasmacytic lymphoma/waldenström macroglobulinemia in a same patient: A case report
Zhang LB, Zhang L, Xin HL, Wang Y, Bao HY, Meng QQ, Jiang SY, Han X, Chen WR, Wang JN, Shi XF
- 4306 Collagen fleece (Tachosil®) for treating testis torsion: A case report
Kim KM, Kim JH

- 4313** Morphological features and endovascular repair for type B multichanneled aortic dissection: A case report
Lu WF, Chen G, Wang LX
- 4318** Hepatic inflammatory myofibroblastic tumor: A case report
Tong M, Zhang BC, Jia FY, Wang J, Liu JH
- 4326** Endometriosis of the lung: A case report and review of literature
Yao J, Zheng H, Nie H, Li CF, Zhang W, Wang JJ
- 4334** Delayed dislocation of the radial head associated with malunion of distal radial fracture: A case report
Kim KB, Wang SI
- 4341** Synchronous endometrial and ovarian cancer: A case report
Žilovič D, Čiurlienė R, Šidlovská E, Vaicekauskaitė I, Sabaliauskaitė R, Jarmalaitė S
- 4350** Nivolumab-induced tumour-like gastritis: A case report
Cijauskaite E, Kazenaite E, Strainiene S, Sadauskaite G, Kurlinkus B
- 4360** Solitary thyroid gland metastasis from rectal cancer: A case report and review of the literature
Chen Y, Kang QS, Zheng Y, Li FB
- 4368** Anesthesia for extracorporeal membrane oxygenation-assisted thoracoscopic lower lobe subsegmental resection in a patient with a single left lung: A case report
Wang XF, Li ZY, Chen L, Chen LX, Xie F, Luo HQ
- 4377** Indium chloride bone marrow scintigraphy for hepatic myelolipoma: A case report
Sato A, Saito K, Abe K, Sugimoto K, Nagao T, Sukeda A, Yunaiyama D
- 4384** Fibromatosis-like metaplastic carcinoma of the breast: Two case reports
Bao WY, Zhou JH, Luo Y, Lu Y
- 4392** Perforating and ophthalmic artery variants from the anterior cerebral artery: Two case reports
Mo ZX, Li W, Wang DF
- 4397** Diagnostic use of superb microvascular imaging in evaluating septic arthritis of the manubriosternal joint: A case report
Seskute G, Kausaite D, Chalkovskaja A, Bulotaite E, Butrimiene I
- 4406** Primary prostate Burkitt's lymphoma resected with holmium laser enucleation of the prostate: A rare case report
Wu YF, Li X, Ma J, Ma DY, Zeng XM, Yu QW, Chen WG
- 4412** Pancreatitis, panniculitis and polyarthritides syndrome: A case report
Pichler H, Stumpner T, Schiller D, Bischofreiter M, Ortmaier R
- 4419** Acute neck tendonitis with dyspnea: A case report
Wu H, Liu W, Mi L, Liu Q

- 4425** Next-generation sequencing technology for the diagnosis of *Pneumocystis* pneumonia in an immunocompetent female: A case report
Huang JJ, Zhang SS, Liu ML, Yang EY, Pan Y, Wu J
- 4433** Superior laryngeal nerve block for treatment of throat pain and cough following laryngeal herpes zoster: A case report
Oh J, Park Y, Choi J, Jeon Y
- 4438** Removal of unexpected schwannoma with superficial parotidectomy using modified-Blair incision and superficial musculoaponeurotic system folding: A case report
Nam HJ, Choi HJ, Byeon JY, Wee SY
- 4446** Simultaneously metastatic cholangiocarcinoma and small intestine cancer from breast cancer misdiagnosed as primary cholangiocarcinoma: A case report
Jiao X, Zhai MM, Xing FZ, Wang XL

LETTER TO THE EDITOR

- 4454** Erroneous presentation of respiratory-hemodynamic disturbances and postsurgical inflammatory responses in patients having undergone abdominal cavity cancer surgery
Idrissov KS, Mynbaev OA

ABOUT COVER

Editorial Board Member of *World Journal of Clinical Cases*, Guoping Zheng, MD, PhD, Associate Professor, Faculty of Medicine and Health, Sydney Medical School-Westmead Clinical School, The University of Sydney, Sydney 2145, Australia. guoping.zheng@sydney.edu.au

AIMS AND SCOPE

The primary aim of *World Journal of Clinical Cases* (WJCC, *World J Clin Cases*) is to provide scholars and readers from various fields of clinical medicine with a platform to publish high-quality clinical research articles and communicate their research findings online.

WJCC mainly publishes articles reporting research results and findings obtained in the field of clinical medicine and covering a wide range of topics, including case control studies, retrospective cohort studies, retrospective studies, clinical trials studies, observational studies, prospective studies, randomized controlled trials, randomized clinical trials, systematic reviews, meta-analysis, and case reports.

INDEXING/ABSTRACTING

The WJCC is now abstracted and indexed in Science Citation Index Expanded (SCIE, also known as SciSearch®), Journal Citation Reports/Science Edition, Current Contents®/Clinical Medicine, PubMed, PubMed Central, Reference Citation Analysis, China National Knowledge Infrastructure, China Science and Technology Journal Database, and Superstar Journals Database. The 2022 Edition of Journal Citation Reports® cites the 2021 impact factor (IF) for WJCC as 1.534; IF without journal self cites: 1.491; 5-year IF: 1.599; Journal Citation Indicator: 0.28; Ranking: 135 among 172 journals in medicine, general and internal; and Quartile category: Q4.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: *Ying-Yi Yuan*; Production Department Director: *Xu Guo*; Editorial Office Director: *Jin-Lei Wang*.

NAME OF JOURNAL

World Journal of Clinical Cases

ISSN

ISSN 2307-8960 (online)

LAUNCH DATE

April 16, 2013

FREQUENCY

Thrice Monthly

EDITORS-IN-CHIEF

Bao-Gan Peng, Jerzy Tadeusz Chudek, George Kontogeorgos, Maurizio Serati, Ja Hyeon Ku

EDITORIAL BOARD MEMBERS

<https://www.wjgnet.com/2307-8960/editorialboard.htm>

PUBLICATION DATE

June 26, 2023

COPYRIGHT

© 2023 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjgnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>



Synchronous endometrial and ovarian cancer: A case report

Diana Žilovič, Rūta Čiurlienė, Evelina Šidlovskā, Ieva Vaicekauskaitė, Rasa Sabaliauskaitė, Sonata Jarmalaitė

Specialty type: Oncology

Provenance and peer review:

Unsolicited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's scientific quality classification

Grade A (Excellent): 0

Grade B (Very good): 0

Grade C (Good): 0

Grade D (Fair): 0

Grade E (Poor): 0

P-Reviewer: Imai Y, Japan

Received: February 6, 2023

Peer-review started: February 6, 2023

First decision: April 26, 2023

Revised: May 4, 2023

Accepted: May 22, 2023

Article in press: May 22, 2023

Published online: June 26, 2023



Diana Žilovič, Rūta Čiurlienė, Department of Oncogynecology, National Cancer Institute, Vilnius 08406, Lithuania

Diana Žilovič, Ieva Vaicekauskaitė, Sonata Jarmalaitė, Institute of Biosciences, Vilnius University, Vilnius 10222, Lithuania

Diana Žilovič, Laboratory of Clinical Oncology, National Cancer Institute, Vilnius 08406, Lithuania

Evelina Šidlovskā, National Center of Pathology, Affiliate of Vilnius University Hospital Santaros Klinikos, Vilnius 08406, Lithuania

Ieva Vaicekauskaitė, Rasa Sabaliauskaitė, Sonata Jarmalaitė, Laboratory of Genetic Diagnostic, National Cancer Institute, Vilnius 08406, Lithuania

Corresponding author: Diana Žilovič, Doctor, Doctor, Researcher, Department of Oncogynecology, National Cancer Institute, 1 Santariskiu Street, Vilnius 08406, Lithuania.

dianauljanionok@gmail.com

Abstract

BACKGROUND

Synchronous endometrial and ovarian cancer (SEOC) is a rare genital tract tumor. Precise diagnosis is crucial for the disease management since prognosis and overall survival differ substantially between metastatic endometrial cancer (EC) or OC. In this review we present 2 cases of women who were diagnosed with SEOC, and discuss the clinical characteristic of SEOC, diagnostic and molecular profiling issues. Next generation sequencing of 10 gene panel was performed on cancerous tissue and uterine lavage samples.

CASE SUMMARY

In our report patients with SEOC had endometrioid type histology with early stage and low-grade histology for both EC and OC. They underwent surgical treatment and staging. Next-generation sequencing of 10 gene-panel identified *CTNNB1*, *PIK3CA*, and *PTEN* gene mutations in ovarian tissue in one case, while none of these genes were mutated in other case. Literature review in support to our data suggest a good prognosis for SEOC diagnosed at early stage.

CONCLUSION

Accurate diagnosis of SEOC is essential for disease management and gene mutation analysis can be helpful as a complementary diagnostic and prognostic tool.

Key Words: Ovarian cancer; Endometrial cancer; Synchronous primary cancer; Uterine lavage; Case report

©The Author(s) 2023. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: Synchronous gynecological tumors are a rare entity. The most common synchronous tumor is synchronous endometrial and ovarian cancer (SEOC). The importance of distinguishing SEOC from either isolated endometrium or ovarian cancer with metastasis is crucial, as it determines management and prognosis. When molecular testing was carried out for SEOC cases, a large proportion was found to be metastatic disease. Testing of vimentin, molecular analyses of gene mutation of *CTNNB1*, *Paired box gene 8*, β -catenin expression may be helpful to categorize SEOC in cases where clinical and pathological parameters are inconclusive.

Citation: Žilović D, Čiurlienė R, Šidlovská E, Vaicekauskaitė I, Sabaliauskaitė R, Jarmalaitė S. Synchronous endometrial and ovarian cancer: A case report. *World J Clin Cases* 2023; 11(18): 4341-4349

URL: <https://www.wjgnet.com/2307-8960/full/v11/i18/4341.htm>

DOI: <https://dx.doi.org/10.12998/wjcc.v11.i18.4341>

INTRODUCTION

Synchronous gynecological malignant tumor is a rare entity. Synchronous endometrial and ovarian cancer (SEOC) is the most common, accounting for 50%-70% of all synchronous tumors. SEOC is described as the simultaneous presence of ovarian cancer (OC) and endometrial cancer (EC) at the same time of diagnosis. The rate of SEOC is approximately 3%-10%, 5% occur in patients with EC and 10% in patients with OC. Endometrioid histological type of EC is significantly associated with increased risk of secondary primary OC with endometrioid histology. Also, OC with endometrioid histology diagnosed under the age of 50 can be associated with loss of mismatch repair genes expression, Lynch syndrome and other genetic mutations. In these patients, the risk of synchronous primary malignancy is higher. Up to 86% of SEOC have concordant endometrioid histology in the two cancer sites[1-4].

For more than 20 years histological criteria by Ulbright and Roth have been used to distinguish metastatic cancer from double primary cancer[5]. Criteria for SEOC diagnosis were first documented in 1981 and refined substantially in 1998[6,7]. The Scully criteria are based on histologic similarity, size, the presence of precursor lesions, location, and invasion pattern. Later, several authors have proposed methods of molecular analysis for precise SEOC diagnostic, but no consensus has been reached yet. Sometimes it is complicated to distinguish between SEOC and metastatic cancer (metastasis from endometrium to the ovary or ovary to the endometrium), but it is highly significant for proper treatment strategy selection. As SEOC is uncommon, it can be misdiagnosed as FIGO stage III of EC or FIGO stage II of OC. When histology of two tumors is discordant, the diagnosis of SEOC is less complicated, but dilemma occurs for concordant histologic types. The treatment management and overall survival differs significantly from SEOC and metastatic disease. Recent evidence has suggested that criteria by Ulbright and Roth for SEOC diagnostics may not always be applicable. When molecular testing was carried out for SEOC cases, a large proportion was found to be metastatic disease[8-12]. Molecular analyses of gene mutation of *CTNNB1*, *TP53*, *KRAS*, *PIK3CA*, and *PTEN* as well as microsatellite instability, β -catenin expression may be helpful to categorize SEOC in cases where clinical and pathological parameters are inconclusive[13]. In problematic cases molecular analysis of tumor foci may facilitate the precise diagnosis and identify the cell of origin, while mutation analysis in liquid biopsy has potential for early cancer detection.

In present study, next generation sequencing of 10 genes in DNA from uterine lavage and ovarian tissue samples from two cases with SEOC was performed aiming at improved understanding of the molecular profile of these rare tumors and evaluation of rationality of uterine lavage based genetic in SEOC diagnostics.

CASE PRESENTATION

Chief complaints

I clinical case: A 54-year-old, menopausal (gravida 0, para 0) woman presented with lower abdominal pain.

II clinical case: A 45-year-old woman (gravida 1, para 1) arrived to National Cancer Institute of Lithuania with histologically confirmed uterine cancer with no complaints.

History of present illness

I clinical case: The complaints lasted for last six months.

II clinical case: Hysteroscopy was performed because endometrial polyp was detected by ultrasound. Final histology confirmed well differentiated (G1) endometrioid endometrial cancer.

History of past illness

I clinical case: There were no other comorbidities, body mass index (BMI) 22.4 kg/m².

II clinical case: There were no comorbidities, she had a BMI of 27.8 kg/m².

Personal and family history

I clinical case: Her medical and family histories were unremarkable for any type of cancer.

II clinical case: There were no history of malignancy in the family.

Physical examination

I clinical case: The clinical examination revealed mobile pelvic mass 15 cm in diameter which was extended till the umbilicus. Pelvic ultrasound examination identified: Thickened endometrium up to 11 mm and left adnexal multilocular cystic tumor with papillary structures. Computed tomography (CT) scan of the thorax, abdomen and pelvis revealed left cystic ovary mass approximately 136 mm × 114 mm × 114 mm in size with several septum and papillary component.

II clinical case: On clinical examination left adnexal mass was identified. Transvaginal ultrasound showed normal sized uterus with endometrial thickness of 6 mm, and 7 cm × 8 cm left adnexal solid mass with mixed echogenicity and cystic inclusion, right ovary cyst up to 3 cm is likely to be functional cyst. Magnetic resonance imaging of the abdominal and pelvis showed irregularly shaped, solid, left adnexal mass up to 60 mm × 77 mm × 80 mm, suspected paraaortic and iliac lymph nodes.

Laboratory examinations

I clinical case: Cancer antigen 125 was evaluated 209 U/mL (normal range < 35 U/mL).

II clinical case: Cancer antigen 125 was not evaluated 32 U/mL (normal range < 35 U/mL).

Imaging examinations

I clinical case: **Figure 1** shows hematoxylin-eosin stains of the endometrial and **Figure 2** shows ovarian tumor specimens. Peritoneal cytological washing and additional peritoneal biopsies, as well as lymph nodes, were negative for malignant cells. Haematoxylin and eosin (H&E)-stained slide at low power (lens objective × 4) shows closely packed irregular glandular structures. H&E-stained slide at medium power (lens objective × 14) shows columnar epithelium with oval, stratified nuclei and visible nucleoli (**Figure 1A** and **B**). H&E-stained slide at low power (lens objective × 4) shows closely packed irregular glandular structures and focal solid growth. H&E-stained slide at medium power (lens objective × 14) shows tumor composed of moderate amount of eosinophilic cytoplasm with slightly polymorphous nuclei and visible nucleoli (**Figure 1C** and **D**).


II clinical case: **Figure 2A** and **B** shows hematoxylin-eosin stains of the endometrial and **Figure 3A** and **B** shows ovarian tumor specimens. H&E-stained slide at low power (lens objective × 4) shows closely packed irregular glandular structures. H&E-stained slide at medium power (lens objective × 14) shows tall columnar epithelium with oval, stratified nuclei with visible nucleoli (**Figure 2A** and **B**).

H&E-stained slide at low power (lens objective × 4) shows closely packed irregular glandular structures with focal necrosis. H&E-stained slide at medium power (lens objective × 14) shows columnar epithelium with oval, stratified nuclei, and visible nucleoli (**Figure 2C** and **D**).

FINAL DIAGNOSIS

I clinical case

On final histological result: Pathological findings revealed a well-differentiated (G1) endometrioid carcinoma of the uterus without myometrial invasion and a moderately-differentiated (G2) endometrioid tumor limited to left ovary. **Figure 1A** and **B** shows hematoxylin-eosin stains of the endometrial and **Figure 1C** and **D** shows ovarian tumor specimens. Peritoneal cytological washing and additional peritoneal biopsies, as well as lymph nodes, were negative for malignant cells.



DOI: 10.12998/wjcc.v11.i18.4341 Copyright ©The Author(s) 2023.

Figure 1 Haematoxylin and eosin staining in case I. A: Haematoxylin and eosin (H&E)-stained slide at low power (lens objective $\times 4$) shows closely packed irregular glandular structures; B: H&E-stained slide at medium power (lens objective $\times 14$) shows columnar epithelium with oval, stratified nuclei and visible nucleoli; C: H&E-stained slide at low power (lens objective $\times 4$) shows closely packed irregular glandular structures and focal solid growth; D: H&E-stained slide at medium power (lens objective $\times 14$) shows tumor composed of moderate amount of eosinophilic cytoplasm with slightly polymorphous nuclei and visible nucleoli.

Lymphovascular invasion was not seen. The final diagnosis of synchronous FIGO IA endometrioid grade 2 ovarian carcinoma and FIGO IA grade 1 endometrial carcinoma was made.

II clinical case

Histological examination confirmed well-differentiated endometrioid (G1) tumor limited to left ovary and a well-differentiated (G1) endometrioid carcinoma of the uterus without myometrial invasion. **Figure 2A** and **B** shows hematoxylin-eosin stains of the endometrial and **Figure 2C** and **D** shows ovarian tumor specimens. Peritoneal cytological washing and biopsies, as well as lymph nodes, were negative for malignant cells. The final diagnosis of synchronous FIGO IA endometrioid grade 1 ovarian carcinoma and FIGO IA grade 1 endometrial carcinoma was made. Lymphovascular invasion was not seen.

TREATMENT

I clinical case

The patient underwent an exploratory laparotomy, frozen section of ovarian tumors was highly suggestive of malignancy. A total abdominal hysterectomy, bilateral salpingo-oophorectomy, pelvic and para-aortic lymph node dissection, omentectomy, selective peritonectomy were conducted.


II clinical case

The patient underwent surgical staging: Hysterectomy with bilateral salpingo-oophorectomy, peritoneal biopsy, pelvic and para-aortic lymph node dissection.

OUTCOME AND FOLLOW-UP


I clinical case

To gain insight into the tumor biology, uterine lavage and ovarian tissue samples were sequenced using custom targeted 10 gene panel which included genes commonly associated with ovarian and endometrial cancers: *TP53*, *BRCA1*, *BRCA2*, *PIK3CA*, *KRAS*, *PTEN*, *ARID1A*, *CTNBNB1*, *FBXW7*, and



DOI: 10.12998/wjcc.v11.i18.4341 Copyright ©The Author(s) 2023.

Figure 2 Haematoxylin and eosin staining in case II. A: Haematoxylin and eosin-stained (H&E) slide at low power (lens objective $\times 4$) shows closely packed irregular glandular structures; B: H&E-stained slide at medium power (lens objective $\times 14$) shows tall columnar epithelium with oval, stratified nuclei with visible nucleoli; C: H&E-stained slide at low power (lens objective $\times 4$) shows closely packed irregular glandular structures with focal necrosis; D: H&E-stained slide at medium power (lens objective $\times 14$) shows columnar epithelium with oval, stratified nuclei, and visible nucleoli.



Transcription, cell proliferation

DOI: 10.12998/wjcc.v11.i18.4341 Copyright ©The Author(s) 2023.

Figure 3 Crosstalk of PI3K/AKT and WNT pathways in synchronous endometrial and ovarian. Both pathways converge at glycogen synthase kinase-3 beta (GSK3β): In PI3K/AKT pathway, activation of PI3K is required for AKT activation and downstream signaling. PTEN acts as deactivator of PI3K pathway. AKT also phosphorylates and inactivates GSK3β. Meanwhile in WNT signaling GSK3β/Axin complex is required for β -catenin phosphorylation and degradation. The numbers indicate TCGA published mutation data from uterine (right side data) and ovarian (left side data) cancers. Figure generated using PathwayMapper tool. GSK3β: Glycogen synthase kinase-3 beta.

PPP2R1A. No pathogenic mutations were detected in uterine lavage or tissue sample. After surgery, six cycles of chemotherapy of Carboplatin were performed. Patient was disease-free at 44 mo of follow-up.

II clinical case

As in previous case, uterine lavage and ovarian tissue samples were sequenced using custom targeted 10 gene panel. In ovarian tissue sample, somatic mutations of *PIK3CA* and *PTEN* were detected. Moreover, tumor sample had a mutation in β -catenin gene *CTNNB1*. No mutations were found in uterine lavage sample, showing low spread of cancerous cells. Mutation profile for both cases are shown in Table 1. No adjuvant treatment was considered. The patient is disease-free at 42 mo of follow-up.

DISCUSSION

SEOC is a rare variant of gynecological cancers. Usually, SEOC appears with different clinical characteristics compared to patients with EC or OC alone. SEOC is observed among the younger age women under 55 years and 40% of them are nulliparous. In most cases it can be diagnosed at an early stage and is associated with low-grade disease. The endometrioid subtype of the primary tumors is the most common histological finding with the rate of 50%-70% of cases. In our report, both women with SEOC had endometrioid histological type. The most common symptom of SEOC is abnormal uterine bleeding and pelvic pain due to the pelvic mass[9,11].

The importance of distinguishing SEOC from either isolated endometrium or ovarian cancer with metastasis is crucial, as it determines adjuvant treatment strategy and prognosis. The prognosis of patients with synchronous EC and OC is better than the patients with single-organ cancer with ovarian or endometrial spread. Median 5-year progression free survival rate is reported to be 65% for SEOC but is less than 50% for FIGO stage IIIA EC with ovarian spread. If SEOC is diagnosed in early stage the overall survival rate is excellent- up to 90%, in contrast to the poor prognosis noted in metastatic disease [1,2,4,13]. Both patients in our report were diagnosed with low grade endometrioid carcinomas of the ovary and uterus and are still alive more than 2 years after treatment. Based on histological findings alone, concurrent uterine and ovarian endometrioid carcinomas are considered synchronous primaries, when all of the following criteria are met: Both tumors are low grade (FIGO 1 or 2); less than 50% of myometrial invasion is present; ovarian tumor is unilateral, limited to parenchyma and no other site is involved in a malignant process; extensive lymphovascular invasion is absent at any location.

To determine cancer origin accurately, advancement of immunohistochemistry and molecular testing is needed. According to literature it is suggested that immunohistochemistry testing of vimentin can be carried out. Desouki *et al*[14] suggested that a negative stain has a sensitivity and specificity to predict primary OC at 97% and 82%, while positive vimentin staining had an 82% sensitivity and 97% specificity in predicting EC. Another study used different antibodies (ER, PR, HER2, p53, and Ki-67) and they found that ER, PR, BCL2 showed different immunostaining patterns between EC and OC and suggested that they can be used as a surrogate marker in the distinction of these tumors[15]. However, the practical meaning of immunohistochemistry in this setting is still the matter of debate due to SEOC heterogeneity and clonality. Immunohistochemical analysis of hormone receptors and markers is not always helpful, because they did not allow a comprehensive evaluation and comparison of possible clonal origin.

Molecular studies of SEOC used variable approaches-mutation analysis of single or group of genes, microsatellite instability (MSI), loss of heterozygosity[16-20]. Clonality of SEOCs can be confirmed by sequencing different genes using next-generation sequencing. Paired box gene 8 (*PAX8*) is a marker that can be useful in distinguishing between EC and OC, because primary OC express *PAX-8* but not EC metastases[18]. Nuclear localization of β -catenin and presence of *CTNNB1* mutations are associated with SEOC. Irving *et al*[21] identified that *CTNNB1* mutations were restricted to SEOC and were absent in all the metastatic tumors, providing direct evidence for a divergence of molecular oncogenetic mechanisms in the subset of SEOC. Genetic classification of SEOC *vs* metastatic tumors based on β -catenin expression/mutation correlates with clinical outcome, but screening of the tumor suppressor genes may be labor-intensive and expensive. Other studies analyze the DNA mismatch repair protein (MMR) expression in SEOC in comparison with the expression of these proteins in Lynch syndrome. The results of these studies showed that majority of SEOCs are sporadic cancers, but patients with endometrioid or clear cell OC under the age of 53 have higher risk for loss of MMR (*MLH1*, *MSH2*, *MSH6*) expression and Lynch syndrome[16,22]. Kobayashi *et al*[23] reported that the frequency of MSI in SEO was 24%, which is comparable to the frequency of endometrial cancer. In our report, none of the patients had a medical history of Lynch syndrome or other hereditary cancers.

Molecular profiling also can be beneficial for prognostic and predictive meaning. Studies analyzing SEOC mutational profiles identified *PI3KCA*, *PTEN*, *KRAS*, *CTNNB1*, and *ARID1A* gene mutations can be helpful for diagnosis of SEOC[16,22-28]. The study by Reijnen *et al*[22] comparing 50 SEOC cases with metastatic EC, OC and TCGA mutation data, showed that mutational profile of SEOC harbored a profoundly different molecular profile compared to metastatic disease. SEOC were enriched for *PTEN* and *CTNNB1* mutations and *TP53* was mutated less frequently in SEO than in ovarian metastatic cases. Although, SEOC closely resembles endometrial EC, but with slightly less frequent *ARID1A* mutations. In the endometrioid SEO subgroup, *PTEN* mutations were identified in 72% of ovarian tissue and in 75% of endometrial tissue. In contrast, *PTEN* mutations are less common in endometrioid OC, with

Table 1 Mutational profile of synchronous endometrial and ovarian cancer case 2

Gene	Uterine lavage mutation	Tissue mutation	dbSNP	Amino acid change
<i>PIK3CA</i>	-	c.1093G>A	rs1064793732	p.Glu365Lys
<i>PTEN</i>	-	c.517C>T	rs121913293	p.Arg173Cys
<i>CTNNB1</i>	-	c.101G>A	rs28931589	p.Gly34Glu

dbSNP: Single-Nucleotide Polymorphism database.

prevalence only 17%. In endometrioid endometrial carcinomas *PTEN* mutation was found in 67% of cases[22]. Other studies also found frequent mutation of *CTNNB1* in SEOC cases[22]. In our molecular analysis, one case was identified with mutations of the *CTNNB1*, *PIK3CA*, and *PTEN* genes, which are frequently detected in SEOC. Other case Mutation frequency from TCGA database and the crosstalk between PI3K/AKT and WNT pathways are visualized in Figure 3, showing possible molecular mechanism of SEOC development. No mutations were identified in other SEOC case in our study, showing further need of genome-wide analysis for improved understanding of molecular origins of these rare tumors.

CONCLUSION

SEOC predominantly occurs in younger premenopausal women. The identification of SEOC or metastatic endometrium/ovarian disease have great clinical significance, as the disease management, prognosis and overall survival differ. Precise diagnosis of SEOC may require additional molecular or IHC testing in addition to routine histopathologic assessment. According to literature review testing of vimentin, molecular analyses of gene mutation of *CTNNB1*, *PAX8*, and β -catenin expression may be helpful to categorize SEOC in cases where clinical and pathological parameters are inconclusive. Our two cases do not show that uterine lavage was suitable diagnostic test for SEOC, but further investigation is needed, and additional disease-specific biomarkers need to be discovered.

FOOTNOTES

Author contributions: Žilovič D, Čiurlienė R, and Jarmalaitė S contributed to conceptualization; Žilovič D and Čiurlienė R contributed to sample collection; Vaicekauskaitė I, Vaicekauskaitė I, and Sabaliauskaitė R contributed to sample preparation, and DNA extraction; Žilovič D, Čiurlienė R, and Šidlovskā E contributed to clinical data curation; Žilovič D contributed to writing-original draft preparation; Vaicekauskaitė I and Šidlovskā E contributed to visualization; Jarmalaitė S contributed to supervision; all authors contributed to formal analysis and writing-review and editing; and all authors have read and agreed to the published version of the manuscript.

Informed consent statement: Informed written consent was obtained from the patients for the publication of this report and any accompanying images.

Conflict-of-interest statement: The authors declare that they have no conflict of interest.

CARE Checklist (2016) statement: The authors have read the CARE Checklist (2016), and the manuscript was prepared and revised according to the CARE Checklist (2016).

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Country/Territory of origin: Lithuania

ORCID number: Diana Žilovič 0000-0001-8381-1551.

S-Editor: Chen YL

L-Editor: A

P-Editor: Chen YL

REFERENCES

- 1 **Zaino R**, Whitney C, Brady MF, DeGeest K, Burger RA, Buller RE. Simultaneously detected endometrial and ovarian carcinomas—a prospective clinicopathologic study of 74 cases: a gynecologic oncology group study. *Gynecol Oncol* 2001; **83**: 355-362 [PMID: [11606097](#) DOI: [10.1006/gyno.2001.6400](#)]
- 2 **Soliman PT**, Slomovitz BM, Broaddus RR, Sun CC, Oh JC, Eifel PJ, Gershenson DM, Lu KH. Synchronous primary cancers of the endometrium and ovary: a single institution review of 84 cases. *Gynecol Oncol* 2004; **94**: 456-462 [PMID: [15297188](#) DOI: [10.1016/j.ygyno.2004.05.006](#)]
- 3 **Chen T**, Brenner H, Fallah M, Jansen L, Castro FA, Geiss K, Holleczer B, Katalinic A, Luttmann S, Sundquist K, Rensing M, Xu L, Hemminki K; GEKID Cancer Survival Working Group+. Risk of second primary cancers in women diagnosed with endometrial cancer in German and Swedish cancer registries. *Int J Cancer* 2017; **141**: 2270-2280 [PMID: [28815572](#) DOI: [10.1002/ijc.30930](#)]
- 4 **Matsuo K**, Machida H, Blake EA, Holman LL, Rimel BJ, Roman LD, Wright JD. Trends and outcomes of women with synchronous endometrial and ovarian cancer. *Oncotarget* 2018; **9**: 28757-28771 [PMID: [29983894](#) DOI: [10.18632/oncotarget.25550](#)]
- 5 **Ulbright TM**, Roth LM. Metastatic and independent cancers of the endometrium and ovary: a clinicopathologic study of 34 cases. *Hum Pathol* 1985; **16**: 28-34 [PMID: [2982713](#) DOI: [10.1016/s0046-8177\(85\)80210-0](#)]
- 6 **Selvaggi SM**. Tumors of the ovary, maldeveloped gonads, fallopian tube, and broad ligament. *Arch Pathol Lab Med* 2000; **124**: 477 [PMID: [10705419](#)]
- 7 **Scully RE**, Young RH. Metastatic Tumors of the Ovary. In: Kurman RJ, Hedrick Ellenson L, Ronnett BM, editors. Blaustein's Pathology of the Female Genital Tract. Cham: Springer International Publishing; 2002: 987-90 [DOI: [10.1007/978-1-4614-3165-7](#)]
- 8 **Song T**, Seong SJ, Bae DS, Suh DH, Kim DY, Lee KH, Lim MC, Lee TS. Synchronous primary cancers of the endometrium and ovary in young women: a Korean Gynecologic Oncology Group Study. *Gynecol Oncol* 2013; **131**: 624-628 [PMID: [24051220](#) DOI: [10.1016/j.ygyno.2013.09.009](#)]
- 9 **Shin W**, Park SY, Kang S, Lim MC, Seo SS. How to manage synchronous endometrial and ovarian cancer patients? *BMC Cancer* 2021; **21**: 489 [PMID: [33933018](#) DOI: [10.1186/s12885-021-08220-w](#)]
- 10 **Matias-Guiu X**, Stewart CJR. Endometriosis-associated ovarian neoplasia. *Pathology* 2018; **50**: 190-204 [PMID: [29241974](#) DOI: [10.1016/j.pathol.2017.10.006](#)]
- 11 **Amaral PI**, Silva A, Lacerda A, Barros C. Synchronous endometrioid endometrial and ovarian cancer in a 34-year-old woman. *BMJ Case Rep* 2015; **2015** [PMID: [26351313](#) DOI: [10.1136/bcr-2015-210940](#)]
- 12 **Anglesio MS**, Wang YK, Maassen M, Horlings HM, Bashashati A, Senz J, Mackenzie R, Grewal DS, Li-Chang H, Karnezis AN, Sheffield BS, McConechy MK, Kommoss F, Taran FA, Staebler A, Shah SP, Wallwiener D, Brucker S, Gilks CB, Kommoss S, Huntsman DG. Synchronous Endometrial and Ovarian Carcinomas: Evidence of Clonality. *J Natl Cancer Inst* 2016; **108**: djv428 [PMID: [26832771](#) DOI: [10.1093/jnci/djv428](#)]
- 13 **Singh N**. Synchronous tumours of the female genital tract. *Histopathology* 2010; **56**: 277-285 [PMID: [20459528](#) DOI: [10.1111/j.1365-2559.2009.03367.x](#)]
- 14 **Desouki MM**, Kallas SJ, Khabele D, Crispens MA, Hameed O, Fadare O. Differential vimentin expression in ovarian and uterine corpus endometrioid adenocarcinomas: diagnostic utility in distinguishing double primaries from metastatic tumors. *Int J Gynecol Pathol* 2014; **33**: 274-281 [PMID: [24681739](#) DOI: [10.1097/PGP.0b013e31829040b5](#)]
- 15 **Halperin R**, Zehavi S, Hadas E, Habler L, Bukovsky I, Schneider D. Simultaneous carcinoma of the endometrium and ovary vs endometrial carcinoma with ovarian metastases: a clinical and immunohistochemical determination. *Int J Gynecol Cancer* 2003; **13**: 32-37 [PMID: [12631217](#) DOI: [10.1046/j.1525-1438.2003.13025.x](#)]
- 16 **Hájková N**, Tichá I, Hojný J, Němejcová K, Bárta M, Michálková R, Zikán M, Cibula D, Laco J, Geryk T, Méhes G, Dunder P. Synchronous endometrioid endometrial and ovarian carcinomas are biologically related: A clinico-pathological and molecular (next generation sequencing) study of 22 cases. *Oncol Lett* 2019; **17**: 2207-2214 [PMID: [30675285](#) DOI: [10.3892/ol.2018.9855](#)]
- 17 **AlHilli MM**, Dowdy SC, Weaver AL, St Sauver JL, Keeney GL, Mariani A, Podratz KC, Bakkum-Gamez JN. Incidence and factors associated with synchronous ovarian and endometrial cancer: a population-based case-control study. *Gynecol Oncol* 2012; **125**: 109-113 [PMID: [22210467](#) DOI: [10.1016/j.ygyno.2011.12.444](#)]
- 18 **Schultheis AM**, Ng CK, De Filippo MR, Piscuoglu S, Macedo GS, Gatiús S, Perez Mies B, Soslow RA, Lim RS, Viale A, Huberman KH, Palacios JC, Reis-Filho JS, Matias-Guiu X, Weigelt B. Massively Parallel Sequencing-Based Clonality Analysis of Synchronous Endometrioid Endometrial and Ovarian Carcinomas. *J Natl Cancer Inst* 2016; **108**: djv427 [PMID: [26832770](#) DOI: [10.1093/jnci/djv427](#)]
- 19 **Castro IM**, Connell PP, Waggoner S, Rotmensh J, Mundt AJ. Synchronous ovarian and endometrial malignancies. *Am J Clin Oncol* 2000; **23**: 521-525 [PMID: [11039516](#) DOI: [10.1097/00000421-200010000-00018](#)]
- 20 **Guerra F**, Girolimetti G, Perrone AM, Procaccini M, Kurelac I, Ceccarelli C, De Biase D, Caprara G, Zamagni C, De Iaco P, Santini D, Gasparre G. Mitochondrial DNA genotyping efficiently reveals clonality of synchronous endometrial and ovarian cancers. *Mod Pathol* 2014; **27**: 1412-1420 [PMID: [24633194](#) DOI: [10.1038/modpathol.2014.39](#)]
- 21 **Irving JA**, Catasús L, Gallardo A, Bussaglia E, Romero M, Matias-Guiu X, Prat J. Synchronous endometrioid carcinomas of the uterine corpus and ovary: alterations in the beta-catenin (CTNNB1) pathway are associated with independent primary tumors and favorable prognosis. *Hum Pathol* 2005; **36**: 605-619 [PMID: [16021566](#) DOI: [10.1016/j.humpath.2005.03.005](#)]
- 22 **Reijnen C**, Küsters-Vandeveldt HVN, Ligtenberg MJL, Bulten J, Oosterwegel M, Snijders MPLM, Sweepers S, de Hullu JA, Vos MC, van der Wurff AAM, van Altena AM, Eijkelenboom A, Pijnenborg JMA. Molecular profiling identifies synchronous endometrial and ovarian cancers as metastatic endometrial cancer with favorable clinical outcome. *Int J Cancer* 2020; **147**: 478-489 [PMID: [32022266](#) DOI: [10.1002/ijc.32907](#)]
- 23 **Kobayashi Y**, Nakamura K, Banno K, Aoki D. Current status of diagnosis for synchronous endometrial and ovarian carcinomas by molecular biological approach. *Eur J Gynaecol Oncol* 2021; **42**: 1300-1302

- 24 **Valtcheva N**, Lang FM, Noske A, Samartzis EP, Schmidt AM, Bellini E, Fink D, Moch H, Rechsteiner M, Dedes KJ, Wild PJ. Tracking the origin of simultaneous endometrial and ovarian cancer by next-generation sequencing—a case report. *BMC Cancer* 2017; **17**: 66 [PMID: [28103826](#) DOI: [10.1186/s12885-017-3054-6](#)]
- 25 **Bahceci I**, Dogrusoz U, La KC, Babur Ö, Gao J, Schultz N. PathwayMapper: a collaborative visual web editor for cancer pathways and genomic data. *Bioinformatics* 2017; **33**: 2238-2240 [PMID: [28334343](#) DOI: [10.1093/bioinformatics/btx149](#)]
- 26 **Chao A**, Wu RC, Jung SM, Lee YS, Chen SJ, Lu YL, Tsai CL, Lin CY, Tang YH, Chen MY, Huang HJ, Chou HH, Huang KG, Chang TC, Wang TH, Lai CH. Implication of genomic characterization in synchronous endometrial and ovarian cancers of endometrioid histology. *Gynecol Oncol* 2016; **143**: 60-67 [PMID: [27498588](#) DOI: [10.1016/j.ygyno.2016.07.114](#)]
- 27 **Ishikawa M**, Nakayama K, Nakamura K, Ono R, Yamashita H, Ishibashi T, Minamoto T, Iida K, Razia S, Ishikawa N, Kyo S. High frequency of POLE mutations in synchronous endometrial and ovarian carcinoma. *Hum Pathol* 2019; **85**: 92-100 [PMID: [30448219](#) DOI: [10.1016/j.humpath.2018.11.001](#)]
- 28 **Matsubayashi H**, Higashigawa S, Kiyozumi Y, Horiuchi Y, Hirashima Y, Kado N, Abe M, Ohishi T, Ohnami S, Urakami K, Yamaguchi K. Metachronous ovarian endometrioid carcinomas in a patient with a PTEN variant: case report of incidentally detected Cowden syndrome. *BMC Cancer* 2019; **19**: 1014 [PMID: [31664961](#) DOI: [10.1186/s12885-019-6272-2](#)]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

Telephone: +1-925-3991568

E-mail: bpgoffice@wjgnet.com

Help Desk: <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

