

# World Journal of *Gastroenterology*

*World J Gastroenterol* 2018 September 14; 24(34): 3813-3964



**EDITORIAL**

- 3813 Clinical impact of microbiome in patients with decompensated cirrhosis  
*Oikonomou T, Papatheodoridis GV, Samarkos M, Goulis I, Cholongitas E*

**REVIEW**

- 3821 Implication of neurohormonal-coupled mechanisms of gastric emptying and pancreatic secretory function in diabetic gastroparesis  
*Mussa BM, Sood S, Verberne AJ*
- 3834 Drug resistance and new therapies in colorectal cancer  
*Van der Jeught K, Xu HC, Li YJ, Lu XB, Ji G*

**MINIREVIEWS**

- 3849 Role of two-dimensional shear wave elastography in chronic liver diseases: A narrative review  
*Jeong JY, Cho YS, Sohn JH*

**ORIGINAL ARTICLE****Basic Study**

- 3861 Delta-like ligand 4 in hepatocellular carcinoma intrinsically promotes tumour growth and suppresses hepatitis B virus replication  
*Kunanopparat A, Issara-Amphorn J, Leelahavanichkul A, Sanpavat A, Patumraj S, Tangkijvanich P, Palaga T, Hirankarn N*
- 3871 Optimal immunosuppressor induces stable gut microbiota after liver transplantation  
*Jiang JW, Ren ZG, Lu HF, Zhang H, Li A, Cui GY, Jia JJ, Xie HY, Chen XH, He Y, Jiang L, Li LJ*
- 3884 Formin-like 3 regulates RhoC/FAK pathway and actin assembly to promote cell invasion in colorectal carcinoma  
*Zeng YF, Xiao YS, Liu Y, Luo XJ, Wen LD, Liu Q, Chen M*
- 3898 Low expression of CDK5RAP3 and DDRGK1 indicates a poor prognosis in patients with gastric cancer  
*Lin JX, Xie XS, Weng XF, Zheng CH, Xie JW, Wang JB, Lu J, Chen QY, Cao LL, Lin M, Tu RH, Li P, Huang CM*

**Retrospective Cohort Study**

- 3908 Gastroduodenal ulcer bleeding in elderly patients on low dose aspirin therapy  
*Fukushi K, Tominaga K, Nagashima K, Kanamori A, Izawa N, Kanazawa M, Sasai T, Hiraishi H*

**Retrospective Study**

- 3919 Predicting the presence of adenomatous polyps during colonoscopy with National Cancer Institute Colorectal Cancer Risk-Assessment Tool  
*Tariq H, Kamal MU, Patel H, Patel R, Ameen M, Shehi E, Khalifa M, Azam S, Zhang A, Kumar K, Baiomi B, Shaikh D, Makker J*

**META-ANALYSIS**

- 3927** Epidemiology of viral hepatitis in Somalia: Systematic review and meta-analysis study

*Hassan-Kadle MA, Mugtaba SO, Ogurtsov PP*

**CASE REPORT**

- 3958** Unicentric Castleman disease presenting as a retroperitoneal peripancreatic mass: A report of two cases and review of literature

*Cheng JL, Cui J, Wang Y, Xu ZZ, Liu F, Liang SB, Tian H*

## Contents

*World Journal of Gastroenterology*  
Volume 24 Number 34 September 14, 2018

### ABOUT COVER

Editorial board member of *World Journal of Gastroenterology*, Mitsushige Sugimoto, MD, PhD, Associate Professor, Division of Digestive Endoscopy, Shiga University of Medical Science Hospital, Otsu 520-2192, Japan

### AIMS AND SCOPE

*World Journal of Gastroenterology* (*World J Gastroenterol*, *WJG*, print ISSN 1007-9327, online ISSN 2219-2840, DOI: 10.3748) is a peer-reviewed open access journal. *WJG* was established on October 1, 1995. It is published weekly on the 7<sup>th</sup>, 14<sup>th</sup>, 21<sup>st</sup>, and 28<sup>th</sup> each month. The *WJG* Editorial Board consists of 642 experts in gastroenterology and hepatology from 59 countries.

The primary task of *WJG* is to rapidly publish high-quality original articles, reviews, and commentaries in the fields of gastroenterology, hepatology, gastrointestinal endoscopy, gastrointestinal surgery, hepatobiliary surgery, gastrointestinal oncology, gastrointestinal radiation oncology, gastrointestinal imaging, gastrointestinal interventional therapy, gastrointestinal infectious diseases, gastrointestinal pharmacology, gastrointestinal pathophysiology, gastrointestinal pathology, evidence-based medicine in gastroenterology, pancreatology, gastrointestinal laboratory medicine, gastrointestinal molecular biology, gastrointestinal immunology, gastrointestinal microbiology, gastrointestinal genetics, gastrointestinal translational medicine, gastrointestinal diagnostics, and gastrointestinal therapeutics. *WJG* is dedicated to become an influential and prestigious journal in gastroenterology and hepatology, to promote the development of above disciplines, and to improve the diagnostic and therapeutic skill and expertise of clinicians.

### INDEXING/ABSTRACTING

*World Journal of Gastroenterology* (*WJG*) is now indexed in Current Contents<sup>®</sup>/Clinical Medicine, Science Citation Index Expanded (also known as SciSearch<sup>®</sup>), Journal Citation Reports<sup>®</sup>, Index Medicus, MEDLINE, PubMed, PubMed Central and Directory of Open Access Journals. The 2018 edition of Journal Citation Reports<sup>®</sup> cites the 2017 impact factor for *WJG* as 3.300 (5-year impact factor: 3.387), ranking *WJG* as 35<sup>th</sup> among 80 journals in gastroenterology and hepatology (quartile in category Q2).

### EDITORS FOR THIS ISSUE

Responsible Assistant Editor: *Xiang Li*  
Responsible Electronic Editor: *Ying-Na Bian*  
Proofing Editor-in-Chief: *Lian-Sheng Ma*

Responsible Science Editor: *Ruo-Yu Ma*  
Proofing Editorial Office Director: *Ze-Mao Gong*

NAME OF JOURNAL  
*World Journal of Gastroenterology*

ISSN  
ISSN 1007-9327 (print)  
ISSN 2219-2840 (online)

LAUNCH DATE  
October 1, 1995

FREQUENCY  
Weekly

EDITORS-IN-CHIEF  
**Andrzej S Tarnawski, MD, PhD, DSc (Med),**  
Professor of Medicine, Chief Gastroenterology, VA Long Beach Health Care System, University of California, Irvine, CA, 5901 E. Seventh Str., Long Beach, CA 90822, United States

EDITORIAL BOARD MEMBERS  
All editorial board members resources online at <http://www.wjgnet.com/1007-9327/editorialboard.htm>

EDITORIAL OFFICE  
Ze-Mao Gong, Director  
*World Journal of Gastroenterology*  
Baishideng Publishing Group Inc  
7901 Stoneridge Drive, Suite 501,  
Pleasanton, CA 94588, USA  
Telephone: +1-925-2238242  
Fax: +1-925-2238243  
E-mail: [editorialoffice@wjgnet.com](mailto:editorialoffice@wjgnet.com)  
Help Desk: <http://www.f6publishing.com/helpdesk>  
<http://www.wjgnet.com>

PUBLISHER  
Baishideng Publishing Group Inc  
7901 Stoneridge Drive, Suite 501,  
Pleasanton, CA 94588, USA  
Telephone: +1-925-2238242  
Fax: +1-925-2238243  
E-mail: [bpgoffice@wjgnet.com](mailto:bpgoffice@wjgnet.com)  
Help Desk: <http://www.f6publishing.com/helpdesk>  
<http://www.wjgnet.com>

PUBLICATION DATE  
September 14, 2018

COPYRIGHT  
© 2018 Baishideng Publishing Group Inc. Articles published by this Open-Access journal are distributed under the terms of the Creative Commons Attribution Non-commercial License, which permits use, distribution, and reproduction in any medium, provided the original work is properly cited, the use is non commercial and is otherwise in compliance with the license.

SPECIAL STATEMENT  
All articles published in journals owned by the Baishideng Publishing Group (BPG) represent the views and opinions of their authors, and not the views, opinions or policies of the BPG, except where otherwise explicitly indicated.

INSTRUCTIONS TO AUTHORS  
Full instructions are available online at <http://www.wjgnet.com/bpg/gerinfo/204>

ONLINE SUBMISSION  
<http://www.f6publishing.com>

## Epidemiology of viral hepatitis in Somalia: Systematic review and meta-analysis study

Mohamed Abdulkadir Hassan-Kadle, Mugtaba Sulaiman Osman, Pavel Petrovich Ogurtsov

Mohamed Abdulkadir Hassan-Kadle, Pavel Petrovich Ogurtsov, Center For the study of the Liver, Department of Hospital Medicine, Peoples' Friendship University of Russia (RUDN University), Moscow 117198, Russia

Mohamed Abdulkadir Hassan-Kadle, Abrar Research and Training Center, Abrar University, Department of Internal Medicine, Division of Gastroenterology and Hepatology, Dufle Specialist Hospital, Mogadishu 25, Somalia

Mugtaba Sulaiman Osman, Department of Psychiatry, School of Medicine, Royal College of Surgeons in Ireland RCSI, Beaumont Hospital, Dublin 9, Ireland

ORCID number: Mohamed Abdulkadir Hassan-Kadle (0000-0003-2304-6135); Mugtaba Sulaiman Osman (0000-0002-5170-6199); Ogurtsov Pavel Petrovich (0000-0001-7939-891X).

**Author contributions:** Hassan-Kadle MA acquired, collected, extracted, and interpreted the data, drafted and revised the article and made the final approval; Mugtaba SO designed the study, analysis and interpreted the data, drafted the article and made the final approval; Ogurtsov PP revised the article and made the final approval.

Supported by RUDN University Program 5-100.

**Conflict-of-interest statement:** The authors deny any conflict of interest.

**PRISMA 2009 Checklist statement:** This systematic review and meta-analysis was conducted as PRISMA guidelines.

**Open-Access:** This article is an open-access article which was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

Manuscript source: Unsolicited Manuscript

Correspondence to: Mohamed Abdulkadir Hassan-Kadle,

MBChB, MD, Doctor, Abrar Research and Training Center, Abrar University, Department of Internal Medicine, Division of Gastroenterology and Hepatology, Dufle Specialist Hospital, 25 Mogadishu, Somalia. [dr.kadle@abrar.edu.so](mailto:dr.kadle@abrar.edu.so)  
Telephone: +252-6-15910409

Received: March 17, 2018

Peer-review started: March 17, 2018

First decision: April 11, 2018

Revised: May 25, 2018

Accepted: June 21, 2018

Article in press: June 21, 2018

Published online: September 14, 2018

### Abstract

#### AIM

To provide a clear understanding of viral hepatitis epidemiology and their clinical burdens in Somalia.

#### METHODS

A systematic review and meta-analysis was conducted as Preferred Reporting Items for Systematic Reviews and Meta-Analyses guidelines. A comprehensive literature search of published studies on viral hepatitis was performed from 1977-2016 in PubMed, Google Scholar, Science Direct, World Health Organization African *Index Medicus* and the Africa Journals Online databases, as well as on the Ministry of Health website. We also captured unpublished articles that were not available on online systems.

#### RESULTS

Twenty-nine studies from Somalia and Somali immigrants (United Kingdom, United States, Italy, Libya) with a combined sample size for each type of viral hepatitis [hepatitis A virus (HAV): 1564, hepatitis B virus (HBV): 8756, hepatitis C virus (HCV): 6257, hepatitis D virus (HDV): 375 and hepatitis E virus (HEV): 278] were analyzed. The overall pooled prevalence rate of HAV was 90.2% (95%CI: 77.8% to 96%). The



HAV prevalence among different age groups was as follows: < 1 year old, 61.54% (95%CI: 40.14% to 79.24%); 1-10 years old, 91.91% (95%CI: 87.76% to 94.73%); 11-19 years old, 96.31% (95%CI: 92.84% to 98.14%); 20-39 years old, 91.3% (95%CI: 83.07% to 95.73%); and > 40 years old, 86.96% (95%CI: 75.68% to 93.47%). The overall pooled prevalence of HBV was 18.9% (95%CI: 14% to 29%). The overall pooled prevalence among subgroups of HBV was 20.5% (95%CI: 5.1% to 55.4%) in pregnant women; 5.7% (95%CI: 2.7% to 11.5%) in children; 39.2% (95%CI: 33.4% to 45.4%) in patients with chronic liver disease, including hepatocellular carcinoma (HCC); 7.7% (95%CI: 4.2% to 13.6%), 12.4% (95%CI: 6.3% to 23.0%) and 11.8% (95%CI: 5.3% to 24.5%) in age groups < 20 years old, 20-39 years old and > 40 years old, respectively. The HBV prevalence among risk groups was 20% (95%CI: 7.19% to 44.64%) in female prostitutes, 21.28% (95%CI: 7.15% to 48.69%) in hospitalized adults, 5.56% (95%CI: 0.99% to 25.62%) in hospitalized children, 60% (95%CI: 31.66% to 82.92%) in patients with acute hepatitis, 33.55% (95%CI: 14.44% to 60.16%) in patients with ancylostomiasis, 12.34% (95%CI: 7.24% to 20.26%) in patients with leprosy and 20.19% (95%CI: 11.28% to 33.49%) in schistosomiasis patients. The overall pooled prevalence of HCV was estimated as 4.84% (95%CI: 3.02% to 7.67%). The prevalence rates among blood donors, risk groups, children and patients chronic liver disease (including HCC) was 0.87% (95%CI: 0.33% to 2.30%), 2.43% (95%CI: 1.21% to 4.8%), 1.37% (95%CI: 0.76% to 2.46%) and 29.82% (95%CI: 15.84% to 48.98%), respectively. The prevalence among genotypes of HCV was 21.9% (95%CI: 15.36% to 30.23%) in genotype 1, 0.87% (95%CI: 0.12% to 5.9%) in genotype 2, 25.21% (95%CI: 18.23% to 33.77%) in genotype 3, 46.24% (95%CI: 37.48% to 55.25%) in genotype 4, 2.52% (95%CI: 0.82% to 7.53%) in genotype 5, and 1.19% (95%CI: 0.07% to 16.38%) in genotype 6. The overall pooled prevalence of HDV was 28.99% (95%CI: 16.38% to 45.96%). The HDV prevalence rate among patients with chronic liver disease, including HCC, was 43.77% (95%CI: 35.09% to 52.84%). The overall pooled prevalence of HEV was 46.86% (95%CI: 5.31% to 93.28%).

## CONCLUSION

Our study demonstrates a high prevalence of all forms of viral hepatitis in Somalia and it also indicates that chronic HBV was the commonest cause of chronic liver disease. This highlights needs for urgent public health interventions and strategic policy directions to controlling the burden of the disease.

**Key words:** Viral hepatitis; Hepatitis A virus; Hepatitis B virus; Hepatitis C virus; Hepatitis D virus; Hepatitis E virus; Systematic review; Meta-analysis, Somalia

© **The Author(s) 2018.** Published by Baishideng Publishing Group Inc. All rights reserved.

**Core tip:** This is the first article reviewing epidemiology of viral hepatitis in Somalia with systematic review and meta-analysis of the published and unpublished reports from 1977 to 2016 among prevalence of all types' viral hepatitis in Somalia.

Hassan-Kadle MA, Mugtaba SO, Ogurtsov PP. Epidemiology of viral hepatitis in Somalia: Systematic review and meta-analysis study. *World J Gastroenterol* 2018; 24(34): 3927-3957 Available from: URL: <http://www.wjgnet.com/1007-9327/full/v24/i34/3927.htm> DOI: <http://dx.doi.org/10.3748/wjg.v24.i34.3927>

## INTRODUCTION

Viral hepatitis is a major public health problem affecting several hundred million people globally. The most common types of viral hepatitis are six distinct types that have been identified as hepatitis A, B, C, D, E, G viruses, and they may present in acute form or chronic form, which causes substantial morbidity and mortality, including chronic hepatitis, cirrhosis and hepatocellular carcinoma. The World Health Organization (WHO) estimates that 257 million people worldwide are infected with hepatitis B virus (HBV), which constitutes 3.5% of the population: Africa has the second-largest number of chronic HBV carriers after the Western Pacific regions. Both regions are considered of high endemicity, and the prevalence rate of each region was 6.1% and 6.2%, respectively. It is estimated that approximately 71 million people were living with hepatitis C virus (HCV) infection, which accounts for 1% of the world's population, in 2015. The regions with the highest prevalence of HCV were the eastern Mediterranean region (2.3%) and the European region (1.5%), while the prevalence in Africa was 1.0%. Hepatitis D virus (HDV) affects nearly 15 million people, and hepatitis E virus (HEV) annually infects 20 million people, with over 3.3 million symptomatic cases of hepatitis E and 44600 hepatitis E-related deaths being recorded<sup>[1]</sup> Hepatitis E is one of the leading causes of major outbreaks of acute viral hepatitis worldwide, especially in developing nations<sup>[2]</sup>. Hepatitis A virus (HAV) infection spans the entire world, with specifically high prevalence rates in older children and adults<sup>[3]</sup>. The number of deaths from viral hepatitis has increased from 1.10 million deaths in 2000 to 1.34 million deaths in 2015 compared to deaths from tuberculosis (from 1.67 to 1.37 million deaths), malaria (from 0.86 to 0.44 million deaths) and human Immunodeficiency virus (from 1.46 to 1.06 million deaths) between 2000 and 2015. Approximately 96% of these deaths resulted from complications of chronic HBV (66%) and HCV (30%), although hepatitis E and HAV infections accounted for 3.3% and 0.8% of these deaths, respectively<sup>[1]</sup>. HBV (887000 deaths) accounts for more deaths than HCV infection (399000 deaths),

while complications of both viruses (HBV and HCV) and cirrhosis (720000 deaths) account for more deaths than hepatocellular carcinoma (470000 deaths)<sup>[1]</sup>.

In Somalia, viral hepatitis, especially HBV, is of significant public health importance. Somalia is an area of the world with a high prevalence HBV infection of > 8. There are several studies of the prevalence of HAV, HBV, HCV, HDV, and HEV in Somalia; however, to the best of our knowledge, there is no meta-analysis to provide an overall estimation of the prevalence of all viral hepatitis infections in this country. A recent report explored the reasons for such a dearth of data<sup>[4]</sup>. In Somalia, largely due to the unsettling decades-long civil war, medical staffs are underqualified and undertrained, and limited access to modern laboratory facilities poses substantial diagnostic challenges<sup>[4]</sup>. Somalia is considered to be a country that has no national strategy for the surveillance, prevention and control of viral hepatitis<sup>[5]</sup>. The provision of high-quality epidemiological data for viral hepatitis in Somalia could help motivate the drafting of action at the policy level. To synthesize such high-quality epidemiological estimates, we decided to undertake this systematic review and meta-analysis of studies that reported the population-level prevalence of each type of viral hepatitis (A, B, C, D and E) in Somalia. We also aimed to understand the burden of viral hepatitis in Somalia, especially HBV and HCV, and to inform public health practitioners, researchers and policy makers.

## MATERIALS AND METHODS

### Study area

Somalia is located along the Gulf of Aden and Indian Ocean in the sub-region of East Africa, and it is bordered by Djibouti and the Gulf of Aden to the north, the Indian Ocean to the east, Kenya to the southwest and Ethiopia to the west. Somalia has the longest coastline in Africa. It has a land mass of 637657 km<sup>2</sup> and a population of approximately 12 million, 61% of whom live in rural areas. Life expectancy at birth in 2015 was 49 years for males and 54 years for females<sup>[6]</sup>. The country is divided into eighteen administrative regions and 92 districts. Seventy-one percent of the Somali labor force is in agriculture, and almost everyone living in rural areas is involved in livestock or in farming. The per-capita total health expenditure, as a percentage of gross domestic products, was not mentioned in World Bank estimations. The proportion of Somalia's population living below the poverty line has not been published to date, but the country was the fifth-poorest country of the world according the World Bank in 2016<sup>[7]</sup>. According to the United Nation, the Human Development Index of Somalia stood at 0.285, and the country ranked 165<sup>th</sup> out of 170 countries in 2012<sup>[8]</sup>. The country entered into a civil war in 1991, and many major structures collapsed, including the health sector and especially the public sectors.

### Data sources and search strategy

We conducted an electronic literature search in several biomedical databases, including PubMed, Google Scholar, African Journal Online, WHO African Index Medicus and Science Direct, using different combinations of key words. The search encompassed published and unpublished studies from 1977 to 2016 with epidemiological and/or clinical data on the seroprevalence of viral hepatitis in Somalia. The key words used were as follows: ["hepatitis A" AND (seroprevalence OR prevalence) AND "Somalia"], ["hepatitis B" AND (seroprevalence OR prevalence) AND "Somalia"], ["hepatitis C" AND seroprevalence OR prevalence) AND "Somalia"], ["hepatitis D" AND (seroprevalence OR prevalence) AND "Somalia"], ["hepatitis E" AND (seroprevalence OR prevalence) AND "Somalia"]. We also reconducted the search using full written phrases, such as "Viral hepatitis", "hepatitis A", "hepatitis B" or "hepatitis B surface antigen", "hepatitis C", "hepatitis D", "hepatitis E", "epidemiology", and "Somali immigrants." We searched unpublished studies from other sources, such as universities and the website of the Ministry of Health (<http://www.moh.gov.so/en/>) for non-indexed studies or reports on the topic. The statement was reviewed systemically following the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines<sup>[9]</sup>. If the full text of studies were not reachable, we contacted the authors by email.

### Study selection, inclusion and exclusion criteria

In this systematic review, we considered all studies published in peer-reviewed journals between 1977 and 2016, and unpublished primary data of each type of viral hepatitis in Somalia qualified for inclusion in the review, according to the PRISMA flow diagram<sup>[9]</sup>, which excludes case reports, systematic reviews, case series editorials, letters to the editor, commentaries, magazines, newspaper reports/articles and studies of other Somali ethnicities living in neighboring countries published in peer-reviewed journals. The titles and abstracts were screened for relevance, and full-text papers considered relevant for further screening were obtained wherever possible. The references of all identified full-text articles and reviews of the literature were also checked to identify whether there were any additional articles that were missed during screening. For the study selection of this systematic review, we sub-analyzed all articles according to population sub-groups, such as blood donors, pregnant women, children, and patients with chronic liver disease and in the general population. These subgroups were examined with no age restriction. For other populations, the age restrictions imposed were children, defined as those 12 years of age and below, and adults, defined those 18 years of age and above. The search language was restricted to English. The studies also included Somali people who immigrated to Italy, the United States, United Kingdom and Libya and were screened for hepatitis viruses.

**Data extraction**

For the data extraction, the two investigators, Hassan-Kadle MA and Mugtaba SO, independently applied the inclusion criteria selected the studies and extracted the data. The extracted data included the following descriptive information: author(s), publication year, country of study (because some studies were conducted among Somali populations outside of Somalia), total sample size, and cases of each type of viral hepatitis in each study.

**Statistical analysis**

**Calculation of pooled prevalence:** The prevalence of viral hepatitis was extracted from every individual observational study by dividing the number of patients who tested positive over the total number included in the study. The prevalence in each study was multiplied by a weight inversely proportional to the total size of the study sample. The pooled prevalence is the sum of these weighted prevalence estimates in each individual study. We used the R software statistical packages “meta” and “metaphor” to calculate the pooled prevalence (Viechtbauer *et al.*<sup>[10]</sup>; Schwarzer *et al.*<sup>[11]</sup>). We chose to use a random effects model in the meta-analysis because we expected considerable heterogeneity among studies, due largely to the different settings in which studies were conducted. To compare the results, we included the results for a fixed effects meta-analysis. We attempted a meta-regression model by adjusting for the potential effect of the country where the study was conducted<sup>[12]</sup>.

**Testing heterogeneity:** Heterogeneity testing was performed using the  $I^2$  and the Q statistic methods. We interpreted the  $I^2$  statistic results as follows: 0%, no observed heterogeneity; 25%, low; 50%, moderate; and 75%, high heterogeneity. We choose the  $P$  value = 0.05 as a cutoff for significant heterogeneity in interpreting the results of the Q statistic.

**Publication bias:** We used the funnel plot method to visually assess the studies by plotting the sample size against the observed prevalence. We assumed that smaller studies would vary more considerably around the pooled estimate than the larger studies would. Objective measures of the funnel plot symmetry were performed using Duval and Tweedie's trim and fill procedure (Duval *et al.*<sup>[13]</sup> and Egger's test (Egger *et al.*<sup>[14]</sup>).

**RESULTS****Search results**

Our search yielded 504 citations, which were retrieved in the literature review. After reviewing the abstracts and titles, we excluded duplicates and irrelevant studies after applying the exclusion criteria. A total of 29 articles were included in this study and were analyzed. The 29 studies were conducted in Somalia and outside of

Somalia (among Somali immigrants, Figure 1).

**Epidemiology of HAV**

Hepatitis A is a liver disease caused by HAV, and it occurs worldwide. This virus thus creates a public health concern, primarily in developing countries, due to its persistent circulation in the environment. Among the studies presented in Somalia, more than 90% of children had the HAV antibody by the age of 4 years<sup>[15-19]</sup>. In 1992, Mohamud KB and his colleagues studied a Somali sample of 593 subjects who were healthy rural and urban volunteers and child outpatients ages 0-83 years in three villages in Somalia (Mogadishu area: Buur-Ful village; Jowhar District: Mooda Moode; and Bur-Hakaba District and Bajuni Islands: Kismaio District). This sample showed a very high rate of HAV exposure of approximately 90%<sup>[15]</sup>. Another study by Bile *et al.*<sup>[16]</sup> conducted at two institutions for children in Somalia (Shebeli: 596 subjects and Societe Organization Sociale, SOS: 76 subjects) showed a very high rate of HAV in the two samples of 96% and 59%, respectively. One study indicated that HAV in Somalia occurs primarily between 4 mo to 4 years of age, because the child has passive immunity from maternal antibodies during the first 3 mo of life<sup>[17]</sup>. Sebastiani *et al.*<sup>[18]</sup> presented a result of 90.6%. Another study conducted in Italy for immigrant communities, which included 213 subjects, mostly originating from Somalia (177 or 83%), Ethiopia (21 or 10%), and Djibouti, Egypt and Saudi Arabia, showed a very high prevalence of HAV of 96%, including children (87.5% of children were under 12 years)<sup>[19]</sup>. These reports of anti-HAV prevalence rates across the 4 studies ranged from 59.2% to 96%<sup>[15-19]</sup>, as shown in Table 1. Four studies met the inclusion criteria for the meta-analysis of the prevalence of HAV infection. The four studies included examined the prevalence of HAV infection in the total of 1564 Somali participants, and the quality of the included studies also varied. We used the extracted data of the 4 studies to quantify the overall pooled prevalence of HAV infection. The pooled effect size for the prevalence of HAV infection among Somali people was 90.2% (95%CI: 77.8% to 96%). The heterogeneity was high ( $I^2$  = 95.6%, 95%CI: 92.2% to 97.5%). Another indication of high heterogeneity was the Q-statistic [Q (degrees of freedom = 4) = 90.31,  $P$  value < 0.001]. The result of this analysis is presented in the forest plot showing the effect sizes for the individual original studies and their 95%CIs, as shown in Figure 2.

Despite the significant heterogeneity, the funnel plot displayed a symmetric spread of studies in terms of relative weight and effect size, thereby indicating little evidence of publication bias (Figure 3). Notably, the total number of studies was small, and the individual studies were of variable sample size.

Moreover, Duval and Tweedie's trim and fill procedure for the detection of publication bias did not support the possibility of missing studies from the analysis. The






Figure 1 Schematic flow diagram of the studies reviewed for inclusion in analysis.

Table 1 Hepatitis A virus of overall prevalence at Somali population in Somalia and Somali immigrants *n* (%)

Author	Publication year	Total	Hepatitis A virus	Setting	Population
Faustini <i>et al</i> <sup>[19]</sup>	1994	213	204 (96)	Italy	Immigrants
Bile <i>et al</i> <sup>[16]</sup>	1992	596	572 (96)	Somalia	Local
Bile <i>et al</i> <sup>[16]</sup>	1992	76	45 (59.2)	Somalia	Local
Mohamud <i>et al</i> <sup>[15]</sup>	1992	593	534 (90)	Somalia	Local
Sebastiani <i>et al</i> <sup>[18]</sup>	1984	86	78 (90.6)	Somalia	Local

effect size imputed was 87.6%, which was close to the observed effect size. Notably, we could not perform Egger's test for the assessment of symmetry of the funnel plot due to the small number of included studies.

### Sub-analysis according to country/setting

The majority of the studies (*n* = 3) were conducted in Somalia with a pooled prevalence of HAV of 88.1% (95%CI: 71.1% to 95.7%) and high heterogeneity ( $I^2$  = 96.3%). These studies are followed by 1 study conducted in Italy, with a prevalence of HAV of 95.8% (95%CI: 92.1% to 97.8%, Figure 4).

### Age pattern of HAV infection

**Characteristics of the studies included:** All studies met the inclusion criteria for the meta-analysis of

the prevalence of HAV infection and examined the prevalence of HAV infection in a total of 1488 Somali participants belonging to 4 major age groups. The quality of the included studies varied (Table 2).

**Pooled prevalence of HAV infection per age group:** We used the extracted data of the 4 studies to quantify the overall pooled prevalence of HAV infection in each age group.

One study was conducted on patients younger than one year old. The pooled effect size for the prevalence of HAV infection among Somali people aged less than one year was 61.54% (95%CI: 40.14% to 79.24%).

The pooled effect size (out of 4 studies) for the prevalence of HAV infection among Somali people aged 1-10 years old was 91.91% (95%CI: 87.76% to 94.73%). The heterogeneity was moderate ( $I^2$  = 59.9%). Another indication of moderate heterogeneity was the Q-statistic [Q (degrees of freedom = 3) = 7.49].

The pooled effect size (out of 3 studies) for the prevalence of HAV infection among Somali people aged 11-19 years old was 96.31% (95%CI: 92.84% to 98.14%). The heterogeneity was low ( $I^2$  = 26.5%). Another indication of low heterogeneity was the Q-statistic [Q (degrees of freedom = 2) = 2.72].

The pooled effect size (out of 2 studies) for the prevalence of HAV infection among Somali people




Figure 2 Meta-analysis and forest plot presentation of the anti-hepatitis A virus antibody from 1984 to 1994.




Figure 3 Bias assessment plot of studies reporting of hepatitis A virus prevalence in Somalia from 1984 to 1994.

aged 20-39 years old was 91.3% (95%CI: 83.07% to 95.73%). The heterogeneity was moderate ( $I^2 = 46.3\%$ ). Another indication of heterogeneity was the Q-statistic [Q (degrees of freedom = 1) = 1.86].

The pooled effect size (out of 2 studies) for the prevalence of HAV infection among Somali people aged over 40 years old was 86.96% (95%CI: 75.68% to 93.47%). The heterogeneity was low ( $I^2 = 0\%$ ). Another indication of low heterogeneity was the Q-statistic [Q (degrees of freedom = 1) = 0.36]. The result of this analysis is presented in the forest plot showing the effect sizes for the individual original studies and their 95%CI, as shown in Figure 5.

Despite the significant heterogeneity, the funnel plot displayed a symmetric spread of studies in terms of relative weight and effect size, thereby indicating little evidence of publication bias (Figure 6). Notably, the total number of studies was small, and the individual studies were of variable sample size.

Moreover, the Duval and Tweedie's trim and fill procedure for the detection of publication bias did not support the possibility of missing studies from the analysis. The effect size imputed was 88.56%, which was not widely different from the observed effect size.

### Epidemiology of HBV

Somalia is classified among countries as having a high hepatitis B surface antigen (HBsAg) endemicity of more

than 8%<sup>[20,21]</sup>. The first study of HBV in the country was conducted in 1977, in which Delia S and his colleagues were presented with a higher frequency of HBsAg among patients with ancylostomiasis (33.33%) and with urinary schistosomiasis (25.92%) than among leprosy patients (9.67% in the L type and 6.89% in the T type), and the overall prevalence among these patients was shown to be high, with 76.1% (118/155) being observed among patients who were HBsAg positive and 11.11% of the controls. In the leprosy patients with schistosomiasis, the frequency was 40.0%<sup>[22]</sup>. Another study conducted in 1978 showed that HBsAg was found in 14.8% of these patients (54 cases), while the frequency was 34.0% among controls (47 cases). However, the overall prevalence in this study was 23.7% (77/101)<sup>[23]</sup>.


In 1979, Nuti *et al*<sup>[24]</sup> studied 222 Somali patients with the lepromatous form of leprosy (LL;  $n = 135$  patients) and the tuberculoid form of the disease (TT;  $n = 87$  patients) for HBV markers. The results showed that the proportion of leprosy and tuberculoid patients presenting with HBsAg was 24.4% (54/222) and 11.5% (26/222), respectively, but the overall prevalence of HBsAg-positive patients in this article was 36% (80/222)<sup>[24]</sup>. Another study published in this year revealed that among patients with acute viral hepatitis who were tested for the presence of HBsAg and the e-antigen and its corresponding antibodies, HBsAg was found in 60% of patients with hepatitis and 34% of controls, and the overall prevalence of HBsAg-positive patients was 48% (49/102)<sup>[25]</sup>. Nuti M and his colleagues also found that 14% (22/157) of patients were HBsAg-positive for the overall prevalence of their study, but in patients with bladder schistosomiasis and in controls, the prevalence was 19.4% (13/67) and 10% (9/90), respectively<sup>[26]</sup>.

In 1985, a study conducted in three different villages of Somalia found that 12.08% (40/331) of subjects aged 1-83 years were HBsAg-positive, 29.9% were anti-HBs-positive, 43.8% were anti-HBc-positive, 21.4% were anti-HBe-positive, and the overall prevalence in this study was 12% (46/383) of subjects, including those under one year of age. Among the HBsAg-positive subjects, 34.7% were HBeAg-positive and 21.7% had anti-HBcAg-IgM<sup>[27]</sup>. A survey study in 1987 of HBV

**Table 2** Age pattern of hepatitis A infection

Age group	Author/Publication year	Total	Cases	Total	Healthy	Serology
0-11 mo	Mohamud <i>et al</i> 1992 <sup>[15]</sup>	52	32	52	20	Anti-HAV
1-11 yr	Mohamud <i>et al</i> 1992 <sup>[15]</sup>	189	176	189	13	Anti-HAV
	Sebastiani <i>et al</i> 1984 <sup>[18]</sup>	35	33	35	2	Anti-HAV
	Faustini <i>et al</i> 1994 <sup>[19]</sup>	213	186	213	27	Anti-HAV
	Bile <i>et al</i> 1992 <sup>[16]</sup>	234	220	234	14	Anti-HAV
	Total	723	647	723	76	Anti-HAV
11-19 yr	Mohamud <i>et al</i> 1992 <sup>[15]</sup>	62	58	62	4	Anti-HAV
	Sebastiani <i>et al</i> 1984 <sup>[18]</sup>	21	20	21	1	Anti-HAV
	Bile <i>et al</i> 1992 <sup>[16]</sup>	362	353	362	9	Anti-HAV
	Total	445	431	445	14	Anti-HAV
20-39 yr	Mohamud <i>et al</i> 1992 <sup>[15]</sup>	164	153	164	11	Anti-HAV
	Sebastiani <i>et al</i> 1984 <sup>[18]</sup>	19	16	19	3	Anti-HAV
	Total	183	169	183	14	Anti-HAV
40+ yr	Mohamud <i>et al</i> 1992 <sup>[15]</sup>	126	111	126	15	Anti-HAV
	Sebastiani <i>et al</i> 1984 <sup>[18]</sup>	11	9	11	2	Anti-HAV
	Total	137	120	137	17	Anti-HAV

HAV: Hepatitis A virus.

**Figure 4** Forest plot of hepatitis A virus prevalence rates for studies conducted in Somalia from 1984 to 1992.

epidemiology was carried out among 383 adults from different areas of Somalia and in 135 pregnant women and 428 children from Mogadishu. The study showed a high incidence of HBsAg among nomadic males 20/85; (23%) and a lower incidence among males from agricultural and coastal areas, *i.e.*, 16/93 (17%) and 14/98 (14%), respectively. Meanwhile, the lowest frequency of HBsAg was observed among women from coastal areas (6/72; 8%) and among pregnant women (14/135; 10.4%), none of whom had HBeAg. However, a low number of children were HBsAg-positive, both under 4 years old (3/94; 3%) and 4-13 years of age (5/128; 4%). In the 15-19 age group, 50% of subjects showed seroconversion from HBeAg to anti-HBe. A total of 7 out of 41 HBsAg carriers aged over 20 had HBeAg, while the overall prevalence of 8.2% (78/946) was HBsAg-positive<sup>[28]</sup>.

In 1987, Jama H and his colleagues conducted a study of sexual transmitted diseases among varied population groups, and these diseases were detected in 22.4% (49/218) of subjects in the overall populations, including 37% of pregnant women, 4% of neonates,

22% of educated women and 20% of prostitutes<sup>[29]</sup>. Another study in the country showed that 50% (52/104) of subjects was HBsAg-positive<sup>[30]</sup>. In 1989, a total of 1138 subjects with HBsAg were examined from different regions of Somalia; the results showed that 19.3% (220/1138) of subjects were HBsAg-reactive<sup>[31]</sup>. Bile KM and his colleagues conducted a case-control study that detected that 28.8% (67/232) of the overall prevalence was HBsAg subjects, including cases and control groups<sup>[32]</sup>.

A total of 256 serum samples collected from blood donors (157 subjects) and hospitalized children (42 subjects) and adults (57 subjects) in Mogadishu were examined. The results showed that among 198 samples tested, the HBsAg carrier rate was 19.1% (22/115), 5.6% (2/36) and 21.3% (10/47) among blood donors, hospitalized children and hospitalized adults, respectively, but in the overall prevalence, 17.1% (34/198) were positive for HBsAg<sup>[33]</sup>.

The prevalence calculation carried out by Mohamud KB and his colleagues indicated that 10.5% (134/1272) of subjects were positive for HBsAg<sup>[15]</sup>. Another study




Figure 5 Forest plot of hepatitis A virus infection prevalence rates according to age groups from 1984 to 1994.




Figure 6 Bias assessment plot of studies reporting hepatitis A virus and age groups.

published in 1992 showed that 15.9% (95/596) of subjects were HBsAg-positive in one institution at Shabeli residence<sup>[16]</sup>. In a case-control study published in 1993, it was detected that 13.5% (29/124) of subjects in the overall prevalence of both groups were HBsAg-positive<sup>[34]</sup>. In 1994, a study conducted in Italy among immigrant communities detected HBsAg among

3.2% (7/213) of subjects<sup>[19]</sup>. Another study carried out in the United Kingdom among Somali immigrant communities showed that 5.6% (25/439) of subjects were positive for HBsAg<sup>[35]</sup>. Shire AM and his colleagues conducted a study in the United States and found that 13.6% (151/1109) of their sample were positive for HBsAg<sup>[36]</sup>. Unpublished studies performed in the country in 2011 and 2012 showed that 40.1% (59/147)<sup>[37]</sup> and 39.1% (61/156)<sup>[38]</sup> of subjects, respectively, were HBsAg reactive. A study in an immigrant community mostly originating from Somalia showed that 6.2% (31/500) of subjects had been detected with HBsAg<sup>[39]</sup>.

### Pooled prevalence of HBV infection

The reported HBsAg prevalence rates across the 23 studies ranged from 1.9% to 76%, as shown in Table 3. Additionally, the percentage of studies that reported prevalence rates that exceeded 8% was 82.6% (19/23), so we define this endemicity level of an area as high. A total of 17.3% (4/23) of studies reported prevalence rates of at least 8%. The 23 studies met the inclusion criteria for the meta-analysis of the prevalence of hepatitis virus infection and examined the prevalence

**Table 3** Summary of studies of included studies on the prevalence of hepatitis B viral infection in a Somali population in Somalia and Somali immigrants (1977-2014) *n* (%)

No.	Author	Publication year	Total	HBsAg	Healthy	Setting	Population
1	Padovese <i>et al</i> <sup>[39]</sup>	2014	500	31 (6.2)	469	Italy	Immigrants
2	Kadle <i>et al</i> <sup>[38]</sup>	2012	156	61 (39.1)	95	Somalia	Local
3	Shire <i>et al</i> <sup>[36]</sup>	2012	1109	151 (13.6)	958	United States	Immigrants
4	Khadjio <i>et al</i> <sup>[37]</sup>	2011	147	59 (40.1)	88	Somalia	Local
5	Aweis <i>et al</i> <sup>[35]</sup>	2001	439	25 (5.6)	414	United Kingdom	Immigrants
6	Nur <i>et al</i> <sup>[33]</sup>	2000	198	34 (17.1)	164	Somalia	Local
7	Faustini <i>et al</i> <sup>[19]</sup>	1994	213	7 (3.2)	206	Italy	Immigrants
8	Bile <i>et al</i> <sup>[34]</sup>	1993	124	29 (13.5)	95	Somalia	Local
9	Bile <i>et al</i> <sup>[16]</sup>	1992	596	95 (15.9)	501	Somalia	Local
10	Mohamud <i>et al</i> <sup>[15]</sup>	1992	1272	134 (10.5)	1138	Somalia	Local
11	Bile <i>et al</i> <sup>[32]</sup>	1991	232	67 (28.8)	165	Somalia	Local
12	Aceti <i>et al</i> <sup>[31]</sup>	1989	1138	220 (19.3)	918	Somalia	Local
13	Bile <i>et al</i> <sup>[40]</sup>	1991	158	3 (1.9)	155	Somalia	Local
14	Aceti <i>et al</i> <sup>[30]</sup>	1991	104	52 (50)	52	Somalia	Local
15	Jama <i>et al</i> <sup>[29]</sup>	1987	218	49 (22.4)	169	Somalia	Local
16	Bile <i>et al</i> <sup>[28]</sup>	1987	946	78 (8.2)	868	Somalia	Local
17	Sebastiani <i>et al</i> <sup>[27]</sup>	1985	383	46 (12)	337	Somalia	Local
18	Nuti <i>et al</i> <sup>[26]</sup>	1979	102	49 (48)	58	Somalia	Local
19	Nuti <i>et al</i> <sup>[25]</sup>	1979	157	22 (14)	135	Somalia	Local
20	Nuti <i>et al</i> <sup>[24]</sup>	1978	101	24 (23.7)	77	Somalia	Local
21	Nuti <i>et al</i> <sup>[23]</sup>	1979	222	80 (36)	142	Somalia	Local
22	Delia <i>et al</i> <sup>[22]</sup>	1977	155	118 (76.1)	37	Somalia	Local
23	Sebastiani <i>et al</i> <sup>[18]</sup>	1984	86	11 (12.7)	75	Somalia	Local

HBsAg: Hepatitis B surface antigen.

of HBV infection in a total of 8756 Somali participants. The quality of the included studies also varied. We used the extracted data of the 23 studies to quantify the overall pooled prevalence of HBV infection. The pooled effect size for the prevalence of HBV infection among Somali people was 18.9% (95%CI: 14% to 29%). The heterogeneity was high ( $I^2 = 97.6\%$ , 95%CI: 96.2% to 97.6%). Another indication of high heterogeneity was the Q-statistic [Q (degrees of freedom = 22) = 726.17,  $P$  value < 0.001]. The result of this analysis is presented in the forest plot showing the effect sizes for the individual original studies and their 95%CI, as shown in Figure 7.

Despite the significant heterogeneity, the funnel plot displayed a symmetric spread of studies in terms of relative weight and effect size, thereby indicating little evidence of publication bias (Figure 8). Notably, the total number of studies was reasonable, and the individual studies were of variable sample size.

Moreover, Duval and Tweedie's trim and fill procedure for the detection of publication bias did not support the possibility of missing studies from the analysis. The effect size imputed was 18.9%, which was notably similar to the observed effect size. Furthermore, Egger's test for assessment of symmetry of the funnel plot was not statistically significant ( $t = 0.6158$ , degrees of freedom = 21,  $P$ -value = 0.5447), providing further support for the absence of publication bias.

#### Sub-analysis according to country/setting

The majority of the studies ( $n = 19$ ) were conducted in Somalia, with a pooled prevalence of HBV of 23% (95%CI: 16.9% to 30.6%) and high heterogeneity ( $I^2$

= 96.9%). These findings are followed by 2 studies conducted in Italy, with a pooled prevalence of HBV of 4.6% (95%CI: 1.4% to 14.2%) and moderate heterogeneity ( $I^2 = 58.9\%$ ). One study was conducted in the United Kingdom, with a prevalence of HBV of 5.7% (95%CI: 1.1% to 24.6%), and another study was conducted in the United States, with a prevalence of HBV of 13.6% (95%CI: 3.0% to 45.0%) (Figure 9).

#### Meta-regression analysis

Examining the effect of setting, significant variability could be explained by meta-regressing the meta-analysis over the four settings: Italy, Somalia, the United Kingdom, and the United States. Tests of moderators indicated the following: coefficient(s) 2, 3, 4: QM (df = 3) = 10.5687,  $P$ -value = 0.0143. For the prevalence results in Somalia compared to those of Italy, the  $P$  value was 0.0056, and the estimate was 1.8193. For the United Kingdom and the United States, the variability was not significantly different from that of Italy ( $P$  value=0.8373 and 0.2608, respectively).

#### Sub-analysis according to the population

The majority of the studies ( $n = 19$ ) were conducted on the local population in Somalia with a pooled prevalence of HBV of 23% (95%CI: 16.9% to 30.6%) and high heterogeneity ( $I^2 = 96.9\%$ ). The remaining 4 studies were conducted on immigrants with a pooled prevalence of HBV of 23.1% (95%CI: 17.0% to 30.5%) and high heterogeneity ( $I^2 = 92.8\%$ ) (Figure 10).

#### Meta-regression analysis

Examining the effect of population, significant variability






Figure 7 Forest plot of hepatitis B virus infection prevalence rates in Somalia in published and unpublished studies from 1977 to 2014.




Figure 8 Bias assessment plot of studies reporting hepatitis B virus prevalence rate in Somalia from 1977 to 2014.

could be explained by meta-regressing the meta-analysis over the two population types: immigrants and locals. Tests of moderators indicated the following: coefficient 2: QM (df = 1) = 9.5448,  $P$ -val = 0.0020. For the prevalence results of the local Somali population compared to the immigrant population, the  $P$  value was 0.002, and the estimate was 1.4498.

### Pregnant women and HBV infection

A total of 2 studies presented HBV prevalence rates among pregnant women, and these studies showed that 10.4% (14/135) of subjects were HBsAg-positive<sup>[28]</sup>, while the other study presented 37% (19/52) of subjects as positive for HBsAg<sup>[29]</sup>, as shown in Table

4. The two studies met the inclusion criteria for the meta-analysis of the prevalence of HBV infection during pregnancy. The pooled effect size for the prevalence of HBV infection during pregnancy among Somali people was 20.5% (95%CI: 5.1% to 55.4%). The heterogeneity was high ( $I^2 = 93.7\%$ ). See Figure 11.

### HBV infection in children

A total of six studies showed HBV prevalence rates between 0% to 16% among children<sup>[16,27-30,40]</sup>. See Table 5. Six studies met the inclusion criteria for the meta-analysis of the prevalence of HBV infection among children. The pooled effect size for the prevalence of HBV infection in Somali children was 5.7% (95%CI:




Figure 9 Forest plot of hepatitis B virus prevalence rates for studies conducted according to setting from 1977 to 2014.

2.7% to 11.5%). The heterogeneity was high ( $I^2 = 88.1\%$ , 95%CI: 77.8% to 93.6%, Figure 12).

### Hepatitis B in chronic liver disease (including hepatocellular carcinoma)

Six studies met the inclusion criteria for the meta-analysis of the prevalence of HBV infection among patients with chronic liver disease, including patients with hepatocellular carcinoma, in Somali people. Their prevalence ranges from 17.9% to 50%<sup>[30,32,34,36-38]</sup> (Table 6). The pooled effect size for the prevalence of hepatitis B among chronic liver disease patients in Somali people was 39.2% (95%CI: 33.4% to 45.4%). The heterogeneity was moderate ( $I^2 = 53.7\%$ , 95%CI: 0% to 81.5%, Figure 13).

Despite the moderate heterogeneity, the funnel plot displayed a symmetric spread of studies in terms of relative weight and effect size, thereby indicating little evidence of publication bias (Figure 14). Moreover, the Duval and Tweedie's trim and fill procedure for the detection of publication bias did not support the possibility of missing studies from the analysis. The effect size imputed was 41.5%, which was close to the observed effect size (*i.e.*, 39.2%). However, the heterogeneity, expectedly, was statistically significant ( $Q = 18.28$ ,  $df = 7$ ,  $P = 0.0108$ ).

### Sub-analysis according to the age group

The available studies ( $n = 5$ ) contained data on the prevalence of HBV infection among three age groups




Figure 10 Forest plot of hepatitis B virus prevalence rates for studies conducted according to population from 1977 to 2014.

Table 4 Hepatitis B virus infection prevalence rates among pregnant women

Author	Year	Total	HBsAg	Healthy	Setting	Population
Bile <i>et al</i> <sup>[28]</sup>	1987	135	14 (10.3%)	121	Mogadishu	Pregnant women
Jama <i>et al</i> <sup>[29]</sup>	1987	52	19 (37%)	33	Mogadishu	Pregnant women

HBsAg: Hepatitis B surface antigen.

(below 20 years, 20-40 years, and above 40 years) that were available for extraction. The reported prevalence rate of these studies according to age group was as follows: below 20 (1% to 16%)<sup>[16,18,28,40]</sup>; 20-39 (6% to 16%)<sup>[18,28,40]</sup>; and above 40 (0% to 13%)<sup>[18,28,40]</sup> (Table 7). The HBV infection pooled effect was highest among the 20-39 age group at 12.4% (95%CI: 6.3% to 23.0%) followed by the above-40 age group at 11.8% (95%CI: 5.3% to 24.5%). The lowest pooled

prevalence was in the below-20 age group at 7.7% (95%CI: 4.2% to 13.6%). The heterogeneity was very low for the 20-39 and the above-40 age groups ( $I^2 = 0\%$ ) but was very high for the below-20 age group ( $I^2 = 91.1\%$ ) (Figure 15).

The funnel plot indicated little evidence to support publication bias. Furthermore, Egger's test for the assessment of symmetry of the funnel plot was not statistically significant ( $t = -1.8748$ , degrees of freedom = 11,  $P$ -value = 0.0876), providing some support, at the 5% significance level, for the absence of publication bias (Figure 16).



### Risk groups and HBV

**Characteristics of the studies included:** Seven studies met the inclusion criteria for the meta-analysis of the prevalence of HBV infection and their risk groups, as shown in Table 8. The 7 included studies examined the prevalence of HBV infection in a total of 1488 Somali participants belonging to 7 major risk groups.

**Table 5** Hepatitis B infection prevalence rates among Somali children

Author	Year	Total	Cases	HBsAg	Healthy	Setting	Population
Sebastiani <i>et al</i> <sup>[29]</sup>	1985	219	25	11%	194	Jowhar, Bur-Hakaba, Kismaio and Afgoy	Children living in four villages
Bile <i>et al</i> <sup>[16]</sup>	1992	596	95	16%	501	Mogadishu area	Children in government-operated residence for abandoned children in Shebelle
Bile <i>et al</i> <sup>[16]</sup>	1992	76	3	4%	73	Mogadishu	Children in government-operated residences for abandoned children in SOS institution
Bile <i>et al</i> <sup>[40]</sup>	1991	106	0	0%	106	Lower Shabelle	Children living in Mukay Dumis
Nur <i>et al</i> <sup>[30]</sup>	2000	36	2	6%	34	Mogadishu	Hospitalized children
Bile <i>et al</i> <sup>[28]</sup>	1987	428	11	3%	417	Mogadishu	Children living in Mogadishu
Jama <i>et al</i> <sup>[29]</sup>	1987	26	1	4%	25	Mogadishu	Newborn babies

HBsAg: Hepatitis B surface antigen.

**Figure 11** Forest plot of hepatitis B virus infection prevalence rates among pregnant women.**Figure 12** Forest plot of hepatitis B virus infection prevalence rates among Somali children.**Table 6** Studies among patients with chronic liver disease (including hepatocellular carcinoma) in Somalia *n* (%)

Author	Year	Total	HBsAg	Healthy	Setting	Population
Shire <i>et al</i> <sup>[36]</sup>	2012	30	5 (17.9)	25	United States	Immigrants
Kadle <i>et al</i> <sup>[38]</sup>	2012	156	61 (39.1)	95	Somalia	Local
Khadija <i>et al</i> <sup>[37]</sup>	2011	147	59 (40.1)	88	Somalia	Local
Bile <i>et al</i> <sup>[34]</sup>	1993	62	23 (37.1)	39	Somalia	Local
Bile <i>et al</i> <sup>[32]</sup>	1991	116	44 (37.9)	72	Somalia	Local
Aceti <i>et al</i> <sup>[30]</sup>	1991	104	52 (50)	52	Somalia	Local

HBsAg: Hepatitis B surface antigen.

The quality of the included studies also varied.

**Pooled prevalence of HBV infection of Risk groups:** We used the extracted data of the 12 studies to quantify the overall pooled prevalence of HAV

infection in each group.

One study was conducted on female prostitutes. The pooled effect size for the prevalence of HBV infection among Somali prostitutes was 20% (95%CI: 7.19% to 44.64%).

One study was conducted on hospitalized adults. The pooled effect size for the prevalence of HBV infection among Somali hospitalized adults was 21.28% (95%CI: 7.15% to 48.69%).

One study was conducted on hospitalized children. The pooled effect size for the prevalence of HBV infection among Somali hospitalized children was 5.56% (95%CI: 0.99% to 25.62%).

One study was conducted on patients with acute hepatitis. The pooled effect size for the prevalence of HBV infection among Somali acute hepatitis patients was 60% (95%CI: 31.66% to 82.92%).

One study was conducted on patients with




Figure 13 Forest plot of hepatitis B virus infection prevalence rates among patients with chronic liver disease, including hepatocellular carcinoma.

Table 7 Age pattern of hepatitis B virus infection *n* (%)

Age group	Author	Publication year	Total	HBsAg	Healthy
0-20	Sebastiani <i>et al</i> <sup>[40]</sup>	1985	219	25 (11)	194
	Bile <i>et al</i> <sup>[28]</sup>	1987	428	11 (6)	417
	Bile <i>et al</i> <sup>[40]</sup>	1991	119	1 (1)	118
	Bile <i>et al</i> <sup>[16]</sup>	1992	596	95 (16)	501
	Sebastiani <i>et al</i> <sup>[18]</sup>	1984	56	7 (13)	49
	Total		1418	139 (9.8)	1279
20-39	Sebastiani <i>et al</i> <sup>[40]</sup>	1985	94	13 (14)	81
	Bile <i>et al</i> <sup>[28]</sup>	1987	442	62 (14)	380
	Bile <i>et al</i> <sup>[40]</sup>	1991	36	2 (6)	34
	Sebastiani <i>et al</i> <sup>[18]</sup>	1984	19	3 (16)	16
	Total		591	80 (14)	511
40+	Sebastiani <i>et al</i> <sup>[40]</sup>	1985	70	8 (11)	62
	Bile <i>et al</i> <sup>[28]</sup>	1987	76	10 (13)	66
	Bile <i>et al</i> <sup>[40]</sup>	1991	3	0 (0)	3
	Sebastiani <i>et al</i> <sup>[18]</sup>	1984	11	1 (9.1)	10
	Total		160	19 (11.8)	141

HBsAg: Hepatitis B surface antigen.




Figure 14 Bias assessment plot of studies reporting among patients with chronic liver disease, including hepatocellular carcinoma.

ancylostomiasis. The pooled effect size for the prevalence of HBV infection among Somali ancylostomiasis patients was 33.55% (95%CI: 14.44% to 60.16%).

Four studies were conducted on leprosy patients. The pooled effect size for the prevalence of HBV infection among Somali leprosy patients was 12.34% (95%CI: 7.24% to 20.26%). The heterogeneity was high ( $I^2 = 85.3\%$ ). Another indication of high heterogeneity was the Q-statistic [Q (degrees of freedom = 3) = 20.38].

Three studies were conducted on schistosomiasis patients. The pooled effect size for the prevalence of HBV infection among Somali schistosomiasis patients was 20.19% (95%CI: 11.28% to 33.49%). The heterogeneity was low ( $I^2 = 36\%$ ). Another indication of low heterogeneity was the Q-statistic [Q (degrees of freedom = 2) = 3.13].

The result of this analysis is presented in the forest plot showing the effect sizes for the individual original studies and their 95%CI, as shown in Figure 17.

Despite the significant heterogeneity, the funnel plot displayed a symmetric spread of studies in terms of relative weight and effect size, thereby indicating little evidence of publication bias (Figure 18). Notably, the total number of studies was small, and the individual studies were of variable sample size.

Moreover, Duval and Tweedie's trim and fill procedure for the detection of publication bias did not support the possibility of missing studies from the analysis. The effect size imputed was 28.5% which was not widely different from the observed effect size.

### Epidemiology of HCV

There are few studies on HCV infection in Somalia. Watts DM and his colleagues studied 438 subjects, including female prostitutes (1.7% or 4/236), patients from a sexually transmitted disease clinic (2% or 2/80), male soldiers (1.3% or 1/79), and tuberculosis patients (2.3% or 1/43), while the overall prevalence showed that 1.8% (8/438) were anti-HCV positive<sup>[41]</sup>. Another study investigated children from two institutions for children in Somalia, showing that 1.5% of children were positive for anti-HCV in one residence of 596 children, including boys (1.9% or 6/309) and girls (1% or 3/287). However, in corresponding individuals, 76 children (boys and girls) showed a result of 0%<sup>[16]</sup>. A study conducted among blood donors revealed that (0.6% or 1/157) were anti-HCV-positive, while the overall presence of anti-HCV in this study was (2.4% or 6/256)<sup>[33]</sup>. A case-control study of HCV in chronic liver disease patients showed an estimate of 4% (35/885) in overall subjects<sup>[42]</sup>. Bile K and his colleagues found that 23.3% (29/124) of subjects had antibodies for HCV in the overall prevalence of cases and controls<sup>[34]</sup>. In 2014, a study conducted in Malta






Figure 15 Forest plot of hepatitis B virus infection prevalence rates among age groups.




Figure 16 Bias assessment plot of studies reporting among age groups.

among asylum seekers showed that 0.6 % (3/500) of subjects were anti-HCV-positive<sup>[39]</sup>. Another study in Italy among immigrant communities that mostly came from Somalia indicated that 2.3% (5/213) had HCV-positive antibodies<sup>[19]</sup>, and a study conducted in immigrant Somali communities in the United States showed that 9.1% (78/854) of subjects were anti-HCV-positive<sup>[36]</sup> (Table 9). Among population groups at risk for HCV in Somalia,

studies have shown a result of 0% to 7%<sup>[33,41,42]</sup>, shown in Table 10. Eleven studies met the inclusion criteria for the meta-analysis of the prevalence of HCV infection (Table 9).

The 11 included studies examined the prevalence of HCV infection in a total of 6257 Somali participants. The quality of the included studies also varied.

### Pooled prevalence of HCV infection

We used the extracted data of the 12 studies to quantify the overall pooled prevalence of HCV infection. The pooled effect size for the prevalence of HCV infection among Somali people was 4.84% (95%CI: 3.02% to 7.67%). The heterogeneity was high ( $I^2 = 93.5\%$ , 95%CI: 90.4% to 95.6%). Another indication of high heterogeneity was the Q-statistic [Q (degrees of freedom = 10) = 168.5,  $P$  value < 0.001]. The result of this analysis is presented in the forest plot showing the effect sizes for the individual original studies and their 95%CI, as shown in Figure 19.

Despite the significant heterogeneity, the funnel plot displayed a symmetric spread of studies in terms of relative weight and effect size, thereby indicating little




Figure 17 Forest plot of hepatitis B virus infection prevalence rates among risk groups.

Table 8 Risk groups and hepatitis B virus

Author	Year	Total	Cases	Healthy	Population
Nuti <i>et al</i> <sup>[24]</sup>	1979	135	33	102	Patients with leprosy
Nuti <i>et al</i> <sup>[24]</sup>	1979	87	10	77	Patients with leprosy
Nuti <i>et al</i> <sup>[23]</sup>	1978	54	8	46	Patients with schistosomiasis
Nuti <i>et al</i> <sup>[26]</sup>	1979	67	13	54	Patients with schistosomiasis
Delia <i>et al</i> <sup>[22]</sup>	1977	155	52	103	Patients with ancylostomiasis
Delia <i>et al</i> <sup>[22]</sup>	1977	155	40	115	Patients with schistosomiasis
Delia <i>et al</i> <sup>[22]</sup>	1977	155	15	140	Patients with leprosy
Delia <i>et al</i> <sup>[22]</sup>	1977	155	11	144	Patients with leprosy
Nuti <i>et al</i> <sup>[25]</sup>	1979	55	33	22	Patients with acute viral hepatitis
Jama <i>et al</i> <sup>[29]</sup>	1987	85	17	68	Female Prostitutes
Nur <i>et al</i> <sup>[33]</sup>	2000	36	2	34	Hospitalized children
Nur <i>et al</i> <sup>[33]</sup>	2000	47	10	37	Hospitalized adult

evidence of publication bias (Figure 20). Notably, the total number of studies was small, and the individual studies were of variable sample size.

Moreover, Duval and Tweedie's trim and fill procedure for the detection of publication bias did not support the possibility of missing studies from the analysis. The effect size imputed was 8.14%, which was not widely different from the observed effect size. Furthermore, Egger's test for the assessment of symmetry of the funnel plot was not statistically significant ( $t = -1.6334$ , degrees of freedom = 10,  $P$ -value = 0.1334), providing further support for the absence of publication bias.

#### Sub-analysis according to country/setting


The majority of the studies ( $n = 8$ ) were conducted in Somalia, with a pooled prevalence of HCV of 5.02% (95%CI: 2.18% to 11.13%) and high heterogeneity ( $I^2 = 94.9\%$ ). These findings are followed by 2 studies conducted in Italy, with a pooled prevalence of HCV of 1.22% (95%CI: 0.21% to 6.74%) and moderate heterogeneity ( $I^2 = 71.7\%$ ). There was 1 study from the United States with a prevalence of HCV of 9.13%

**Table 9** Summary of studies on the overall prevalence of hepatitis C viral infection in a Somali population in Somalia and Somali immigrants (1970-2016)

	Year	Total	Cases	Total	Healthy	Setting	Population
Daw <i>et al</i> <sup>[43]</sup>	2016	2012	164	2012	1848	Libiya	Immigrants
Padovese <i>et al</i> <sup>[39]</sup>	2014	500	3	500	497	Italy	Immigrants
Kadle <i>et al</i> <sup>[38]</sup>	2012	156	30	156	126	Somalia	Local
Shire <i>et al</i> <sup>[36]</sup>	2012	854	78	854	776	United States	Immigrants
Khadjio <i>et al</i> <sup>[37]</sup>	2011	147	15	147	132	Somalia	Local
Nur <i>et al</i> <sup>[33]</sup>	2000	256	6	256	250	Somalia	Local
Faustini <i>et al</i> <sup>[19]</sup>	1994	213	5	213	208	Italy	Immigrants
Watts <i>et al</i> <sup>[41]</sup>	1994	438	8	430	90	Somalia	Local
Aceti <i>et al</i> <sup>[42]</sup>	1993	885	35	885	850	Somalia	Local
Bile <i>et al</i> <sup>[34]</sup>	1993	124	29	124	95	Somalia	Local
Bile <i>et al</i> <sup>[16]</sup>	1992	596	9	596	587	Somalia	Local
Bile <i>et al</i> <sup>[16]</sup>	1992	76	0	76	76	Somalia	Local

**Table 10** Studies on hepatitis C virus among risk groups in Somalia

Author	Year	Total	Cases	Total	Healthy	Setting	Population
Aceti <i>et al</i> <sup>[42]</sup>	1993	287	0	287	278	Mogadishu	Hospitalized children with diseases other than hepatitis
Watts <i>et al</i> <sup>[41]</sup>	1994	236	4	236	232	Mogadishu, Marka, Kismayo	Female prostitutes
Nur <i>et al</i> <sup>[33]</sup>	2000	42	1	42	41	Mogadishu	Hospitalized children with measles, tuberculosis, anemia and other febrile illnesses
Watts <i>et al</i> <sup>[41]</sup>	1994	80	2	80	78	Mogadishu, Marka, Kismayo	Sexually transmitted disease patients
Nur <i>et al</i> <sup>[33]</sup>	2000	57	4	57	53	Mogadishu	Hospitalized adults with tuberculosis, malaria, acute respiratory infections, and unknown diagnosis (no clinically evident case of hepatitis)
Aceti <i>et al</i> <sup>[42]</sup>	1993	179	4	179	175	Mogadishu	Mixed populations (98 prisoners and 81 psychiatric patients)

**Figure 18** Bias assessment plot of studies reporting among risk groups.

(95%CI: 1.01% to 49.87%). There was 1 study from Libya with a prevalence of HCV of 8.15% (95%CI: 0.89% to 46.60%) (Figure 21).

### Meta-regression analysis

Examining the effect of setting, insignificant variability could be explained by meta-regressing the meta-analysis over the three settings: Italy, Somalia, and the United States. Tests of moderators indicated the following: coefficient(s) 2,3: QM (df = 2) = 2.6557,  $P = 0.2651$ . For the prevalence results in Somalia compared to Italy, the  $P$  value was 0.1476, and the estimate was 1.4512. For the United States, the variability was not significantly different from that of Italy ( $P$  value=0.1560

and estimate = 2.0938).

### Sub-analysis according to refugee status

The majority of the studies ( $n = 8$ ) were conducted in Somalia among non-immigrant populations, with a pooled prevalence of HCV of 4.99% (95%CI: 2.11% to 11.36%) and high heterogeneity ( $I^2 = 94.9\%$ ). The remaining 4 studies were conducted in Italy, Libya and the United States among refugees, showing a pooled prevalence of HCV of 3.81% (95%CI: 1.54% to 9.13%) and high heterogeneity ( $I^2 = 90.3\%$ ) (Figure 22).

### Meta-regression analysis

When examining the effect of refugee status type,




Figure 19 Forest plot of studies reporting chronic hepatitis C virus infection prevalence in Somalia.




Figure 20 Funnel plot of studies reporting chronic hepatitis C virus infection prevalence in Somalia.

insignificant variability could be explained by meta-regressing the meta-analysis over the two statuses: immigrants vs the local population. Tests of moderators indicated the following: coefficient(s) 2: QM (df = 1) = 0.3335,  $P = 0.5636$ . For the prevalence in the local population compared to that in immigrants, the  $P$  value was 0.5636, and the estimate was 0.3383.

### HCV infection in blood donors

**Characteristics of the studies included:** There are 2 studies that met the inclusion criteria for the meta-analysis of the prevalence of HCV infection among blood donors (Table 11). The included 2 studies examined the prevalence of HCV infection in a total of 466 Somali blood donors. The quality of the included studies also varied.

### Pooled prevalence of HCV infection in blood donors

We used the extracted data of the 2 studies to quantify the overall pooled prevalence of HCV infection in Somali blood donors. The pooled effect size for the prevalence of HCV infection among Somali blood donors was 0.87%

(95%CI: 0.33% to 2.30%). The heterogeneity was low ( $I^2 = 0\%$ ). Another indication of low heterogeneity was the Q-statistic [Q (degrees of freedom = 1) = 0.13,  $P = 0.7139$ ]. The result of this analysis is presented in the forest plot showing the effect sizes for the individual original studies and their 95%CI, as shown in Figure 23.

### HCV infection in different genotypes

**Characteristics of the studies included:** Two studies met the inclusion criteria for the meta-analysis of the prevalence of HCV infection among different genotypes (Table 12)<sup>[36,43]</sup>.

### Pooled prevalence of HCV infection per genotype group:

We used the extracted data of the 12 studies to quantify the overall pooled prevalence of HCV infection in each genotype group.

The pooled effect size (out of 2 studies) for the prevalence of HCV infection among Somali people of genotype 1 was 21.9% (95%CI: 15.36% to 30.23%). The heterogeneity was low ( $I^2 = 0\%$ ). Another indication of low heterogeneity was the Q-statistic [Q (degrees of freedom = 1) = 0.20].

The pooled effect size (out of 2 studies) for the prevalence of HCV infection among Somali people of genotype 2 was 0.87% (95%CI: 0.12% to 5.9%). The heterogeneity was low ( $I^2 = 0\%$ ). Another indication of low heterogeneity was the Q-statistic [Q (degrees of freedom = 1) = 0.10].

The pooled effect size (out of 2 studies) for the prevalence of HCV infection among Somali people of genotype 3 was 25.21% (95%CI: 18.23% to 33.77%). The heterogeneity was low ( $I^2 = 0\%$ ). Another indication of low heterogeneity was the Q-statistic [Q (degrees of freedom = 1) = 0.02].

The pooled effect size (out of 2 studies) for the prevalence of HCV infection among Somali people of genotype 4 was 46.24% (95%CI: 37.48% to 55.25%).




Figure 21 Forest plot of hepatitis C virus infection for studies conducted in Somalia and Outside of Somalia.

The heterogeneity was low ( $I^2 = 0\%$ ). Another indication of low heterogeneity was the Q-statistic [Q (degrees of freedom = 1) = 0.57].

The pooled effect size (out of 2 studies) for the prevalence of HCV infection among Somali people of genotype 5 was 2.52% (95%CI: 0.82% to 7.53%). The heterogeneity was low ( $I^2 = 0\%$ ). Another indication of low heterogeneity was the Q-statistic [Q (degrees of freedom = 1) = 0.00].

The pooled effect size (out of 1 studies) for the prevalence of HCV infection among Somali people of genotype 6 was 1.19% (95%CI: 0.07% to 16.38%).

The result of this analysis is presented in the forest plot showing the effect sizes for the individual original studies and their 95%CIs, as shown in Figure 24.

Despite the significant heterogeneity, the funnel plot displayed a symmetric spread of studies in terms of relative weight and effect size, thereby indicating little evidence of publication bias (Figure 25). Notably, the total number of studies was small, and the individual studies were of variable sample size.

Moreover, Duval and Tweedie's trim and fill procedure for the detection of publication bias did not support the possibility of missing studies from the analysis. The effect size imputed was 25.84%, which was not widely different from the observed effect size.

### Risk groups and HCV

**Characteristics of the studies included:** Three studies met the inclusion criteria for the meta-analysis of prevalence of HCV infection (Table 10). The 3 included studies examined the prevalence of HCV infection in the total of 881 Somali participants. The quality of the included studies also varied.

### Pooled prevalence of HCV infection in risk groups:

We used the extracted data of the 6 studies to quantify the overall pooled prevalence of HCV infection in Somali risk groups. The pooled effect size for the prevalence of HCV infection among Somali risk groups was 2.43% (95%CI: 1.21% to 4.8%). The heterogeneity was low ( $I^2 = 41.8\%$ , 95%CI: 0% to 77%). Another indicator of low heterogeneity was the Q-statistic [Q (degrees of freedom = 5) = 8.6,  $P$  value = 0.1263]. The result of this analysis is presented in the forest plot showing the effect sizes for the individual original studies and their 95%CIs, as shown in Figure 26.

As expected from the low heterogeneity, the funnel plot displayed a symmetric spread of studies in terms of relative weight and effect size, thereby indicating little evidence of publication bias (Figure 27). Notably, the total number of studies was small, and the individual studies were of variable sample size.






Figure 22 Forest plot of studies reporting chronic hepatitis C virus prevalence amongst the local population and Somali immigrants.




Figure 23 Forest plot of studies reporting chronic hepatitis C virus prevalence among blood donors in Somalia.

Moreover, Duval and Tweedie's trim and fill procedure for the detection of publication bias did not support the possibility of missing studies from the analysis. The effect size imputed was 3.36%, which was close to the observed effect size.

### HCV infection in Somali children

**Characteristics of the studies included:** Three studies met the inclusion criteria for the meta-analysis of the prevalence of HCV infection (Table 13). The 3 included studies examined the prevalence of HCV infection in a total of 1001 Somali participants. The quality of the included studies also varied.

### Pooled prevalence of HCV infection in children:

We used the extracted data of the 4 studies to quantify the overall pooled prevalence of HCV infection in Somali children. The pooled effect size for the prevalence of HCV infection among Somali children was 1.37% (95%CI: 0.76% to 2.46%). The heterogeneity was low ( $I^2 = 0\%$ , 95%CI: 0% to 83.8%). Another indicator of low heterogeneity was the Q-statistic [Q (degrees of

freedom = 3) = 2.83,  $P$  value = 0.4181]. The result of this analysis is presented in the forest plot showing the effect sizes for the individual original studies and their 95%CIs, as shown in Figure 28.

As expected from the low heterogeneity, the funnel plot displayed a symmetric spread of studies in terms of relative weight and effect size, thereby indicating little evidence of publication bias (Figure 29). Notably, the total number of studies was small, and the individual studies were of variable sample size.

Moreover, Duval and Tweedie's trim and fill procedure for the detection of publication bias did not support the possibility of missing studies from the analysis. The effect size imputed was 1.63%, which was close to the observed effect size.

### HCV infection in Somali chronic liver disease patients

**Characteristics of the studies included:** Five studies met the inclusion criteria for the meta-analysis of the prevalence of HCV infection (Table 14). The 5 included studies examined the prevalence of HCV infection in a total of 505 Somali participants. The quality of the

**Table 11 Studies on hepatitis C among blood donors**

Author	Year	Total	Cases	Healthy	Town	Group
Nur <i>et al</i> <sup>[33]</sup>	2000	157	1	156	Mogadishu	Blood donors
Aceti <i>et al</i> <sup>[42]</sup>	1993	309	3	306	Mogadishu	Blood donors

**Table 12 Studies on distribution genotypes of hepatitis C virus infection in Somalia**

Type of genotype	Author/Publication	Year	Total	Cases	Healthy	Serology
Genotype 1	Daw <i>et al</i> <sup>[43]</sup>	/2016	78	18	60	Anti-HCV
Genotype 1	Shire <i>et al</i> <sup>[36]</sup>	/2012	41	8	33	Anti-HCV
Genotype 2	Daw <i>et al</i> <sup>[43]</sup>	/2016	78	0	78	Anti-HCV
Genotype 2	Shire <i>et al</i> <sup>[36]</sup>	/2012	41	0	41	Anti-HCV
Genotype 3	Daw <i>et al</i> <sup>[43]</sup>	/2016	78	20	58	Anti-HCV
Genotype 3	Shire <i>et al</i> <sup>[36]</sup>	/2012	41	10	31	Anti-HCV
Genotype 4	Daw <i>et al</i> <sup>[43]</sup>	/2016	78	38	40	Anti-HCV
Genotype 4	Shire <i>et al</i> <sup>[36]</sup>	/2012	41	17	24	Anti-HCV
Genotype 5	Daw <i>et al</i> <sup>[43]</sup>	/2016	78	2	76	Anti-HCV
Genotype 5	Shire <i>et al</i> <sup>[36]</sup>	/2012	41	1	10	Anti-HCV
Genotype 6	Shire <i>et al</i> <sup>[36]</sup>	/2012	41	0	40	Anti-HCV

included studies also varied.

**Pooled prevalence of HCV infection in chronic liver disease groups:** We used the extracted data of the 6 studies to quantify the overall pooled prevalence of HCV infection in Somali chronic liver disease groups. The pooled effect size for the prevalence of HCV infection among Somali chronic liver disease groups was 29.82% (95%CI: 15.84% to 48.98%). The heterogeneity was high ( $I^2 = 92.3\%$ , 95%CI: 85% to 96%). Another indicator of high heterogeneity was the Q-statistic [Q (degrees of freedom = 4) = 51.92,  $P$  value < 0.0001]. The result of this analysis is presented in the forest plot showing the effect sizes for the individual original studies and their 95%CIs, as shown in Figure 30.

Despite the high heterogeneity, the funnel plot displayed a symmetric spread of studies in terms of relative weight and effect size, thereby indicating little evidence of publication bias (Figure 31). Notably, the total number of studies was small, and the individual studies were of variable sample size.

Moreover, Duval and Tweedie's trim and fill procedure for the detection of publication bias did not support the possibility of missing studies from the analysis. The effect size imputed was 22.2%, which was close to the observed effect size.

### Epidemiology of HDV infection

Hepatitis D, or delta hepatitis, is caused by the HDV, a unique RNA pathogen. It requires the HBV for its own replication and to infect as co-infection. HDV is transmitted by percutaneous or sexual contact with infected blood or blood products. The main vulnerable group consists of patients with chronic HBsAg infection who become super-infected with the virus. It is estimated that 5% of HBV-infected persons are also

co-infected with HDV<sup>[1]</sup>. Aceti A and his colleagues conducted a study in 1989 showing that among 220 asymptomatic HBsAg-positive carriers, 16.80% (37) had the antibody for HDV<sup>[31]</sup>. In a study of the prevalence of HDV infection in patients with chronic liver disease in Somalia, 52 of the 104 patients studied were positive for HBsAg, and 26 (50%) had anti-delta antibodies<sup>[30]</sup>. In a 1993 study by Khalif Bile and his colleagues with 62 Somali patients with chronic liver disease, including hepatocellular carcinoma, 37.1% (23) of cases with chronic liver disease were positive for HBsAg, and 34.6% (8) of chronic liver disease patients had anti-HDV. Meanwhile, in the control group, 14.3% of those who were positive for HBsAg had anti-HDV<sup>[32]</sup>. Another study conducted among 67 HBsAg-positive patients showed that 30% (20) of these patients were anti-HDV-positive. Additionally, this study showed that 38.6% (17) of the 44 cases of chronic liver disease patients were HBsAg-positive<sup>[32]</sup>. One study examining the prevalence of HDV showed that 34.4% (10/29) of those who were HBsAg-positive and 39.1% (9/23) of chronic liver disease patients who were HBsAg-positive had antibodies for HDV<sup>[34]</sup>. Additionally, 0% of community immigrants from Somalia and neighboring countries who were positive for HBsAg were shown to have anti-HDV<sup>[19]</sup>. Five studies met the inclusion criteria for the meta-analysis of the prevalence of HDV infection, and the five included studies examined the prevalence of HDV infection in a total of 375 Somali participants. The quality of the included studies also varied (Table 15).

### Pooled prevalence of HDV infection

We used the extracted data of the 5 studies to quantify the overall pooled prevalence of HDV infection. The pooled effect size for the prevalence of HDV infection among Somali people was 28.99% (95%CI: 16.38% to 45.96%). The heterogeneity was high ( $I^2 = 84.9\%$ , 95%CI: 66% to 93%). Another indicator of high heterogeneity was the Q-statistic [Q (degrees of freedom = 4) = 26.4,  $P$  value < 0.001]. The result of this analysis is presented in the forest plot showing the effect sizes for the individual original studies and their 95%CIs, as shown in Figure 32.

Despite the significant heterogeneity, the funnel plot displayed a symmetric spread of studies in terms of relative weight and effect size, thereby indicating little evidence of publication bias (Figure 33). Notably, the total number of studies was small, and the individual studies were of variable sample size.

Moreover, Duval and Tweedie's trim and fill procedure for the detection of publication bias did not support the possibility of missing studies from the analysis. The effect size imputed was 28.99%, which was identical to the observed effect size.

### Epidemiology of chronic liver disease and HDV infection

Three studies met the inclusion criteria for the meta-analysis of prevalence of chronic liver disease and HDV




Figure 24 Forest plot of studies reporting on distribution genotypes of hepatitis C virus infection in Somalia.




Figure 25 Bias assessment plot of studies reporting among the distribution of genotypes of hepatitis C virus.

infection. These three included studies had a total of 119 Somali participants, and the quality of the included studies varied (Table 16).

#### **Pooled prevalence of chronic liver disease and HDV infection**

We used the extracted data of the 3 studies to quantify

the overall pooled prevalence of chronic liver disease and HDV infection. The pooled effect size for the prevalence of chronic liver disease and HDV infection among Somali people was 43.77% (95%CI: 35.09% to 52.84%). The heterogeneity was low but had a wide confidence interval ( $I^2 = 0\%$ , 95%CI: 0% to 86%). Another indicator of low heterogeneity was the




Figure 26 Forest plot of studies reporting chronic hepatitis C virus prevalence among risk groups in Somalia.




Figure 27 Bias assessment plot of studies reporting among risk groups in Somalia.

Q-statistic [ $Q$  (degrees of freedom = 2) = 1.49,  $P$  value = 0.4758]. The result of this analysis is presented in the forest plot showing the effect sizes for the individual original studies and their 95% CIs, as shown in Figure 34.

The funnel plot displayed a symmetric spread of the three studies in terms of relative weight and effect size, thereby indicating little evidence of publication bias (Figure 35). Notably, the total number of studies was small, and the individual studies were of variable sample size.

Moreover, Duval and Tweedie's trim and fill procedure for the detection of publication bias did not support the possibility of missing studies from the analysis. The effect size imputed was 43.77%, which was identical to the observed effect size.

### Epidemiology of HEV infection

It is believed that more than 50% of hepatitis E cases in developing countries are unrelated to HAV or HBV infection, and a high proportion of these cases seems to be enterically transmitted<sup>[44]</sup>. Studies on the outbreak of an acute viral hepatitis occurring in three villages in the lower Shabeli region of southern Somalia examined the presence of antibodies to the hepatitis E antigen, and it was found that the prevalence of anti-HEV ranged from 77.8% to 94.0% among the three villages and was widely distributed among all age groups, without a preponderance in any specific age. During the onset of

the outbreak, 111 samples were collected in one of the three villages, which showed that a very low prevalence (4.5% or 5/111) was positive for IgG anti-HEV<sup>[45]</sup>. In another survey of 142 villages with a population of 245312 individuals, 11413 icteric cases were recorded, 346 of whom had died, corresponding to an attack rate and a case fatality rate of 4.6% and 3.0%, respectively. The role of HEV in this epidemic was proven by documenting anti-HEV in 88.2% (128 of 154) of the sampled cases, which served as a sign of recent infection with HEV. Further, the attack rate was found to be higher with increasing age, from 5% in the 1- to 4-year-old age group to 13% in the 5- to 15-year-old age group, and to 20% for persons older than 15 years of age. A high fatality rate of 13.8% was estimated among pregnant females. Therefore, the attack rate was higher (6.0%) in villages supplied with river water, while fewer cases were recorded in those relying on wells (1.7%) or ponds (1.2%) for their water supply<sup>[46]</sup>. Burans and his colleagues conducted a study during the operation of Restore of Hobe in Somalia in 1994, finding that among 31 Somalis with acute hepatitis, 20 (65%) had IgM anti-HEV<sup>[47]</sup>. Three studies met the inclusion criteria for the meta-analysis of the prevalence of HEV infection, and a total of 287 Somali participants were examined. The quality of included studies also varied (Table 17).



### Pooled prevalence of HEV infection

We used the extracted data of the 3 studies to quantify the overall pooled prevalence of HEV infection. The pooled effect size for the prevalence of HEV infection among Somali people was 46.86% (95%CI: 5.31% to 93.28%). The heterogeneity was high ( $I^2 = 97.9\%$ , 95%CI: 96% to 98.9%). Another indicator of high heterogeneity was the Q-statistic [ $Q$  (degrees of freedom = 2) = 93.4,  $P$  value < 0.001]. The result of this analysis is presented in the forest plot showing the effect sizes for the individual original studies and their 95% CIs, as shown in Figure 36.

Because the significant heterogeneity, the funnel plot displayed a somewhat asymmetric spread of studies in terms of relative weight and effect size, thereby indicating evidence of publication bias (Figure 37). Notably, the total number of studies was small, and

**Table 13** Studies on hepatitis C virus infection among Somali children

Author	Year	Total	Cases	Total	Healthy	Setting	Population
Bile <i>et al</i> <sup>[16]</sup>	1992	596	9	596	587	Mogadishu area	Children in government-operated residences for abandoned children in Shebeli
Bile <i>et al</i> <sup>[16]</sup>	1992	76	0	76	76	Mogadishu area	Children in government-operated residence for abandoned children in SOS institution
Nur <i>et al</i> <sup>[33]</sup>	2000	42	1	42	41	Mogadishu	Hospitalized children
Aceti <i>et al</i> <sup>[42]</sup>	1993	287	0	287	278	Mogadishu	Hospitalized children with diseases other than hepatitis

**Figure 28** Forest plot of studies reporting chronic hepatitis C virus prevalence among Somali children.**Figure 29** Bias assessment plot of studies reporting among children in Somalia.

the individual studies were of variable sample size. The study by Mushahwar *et al*<sup>[45]</sup> is clearly an outlier.

Moreover, Duval and Tweedie's trim and fill procedure for the detection of publication bias did support the possibility of missing studies from the analysis. The effect size imputed was 88.28%, which was well above the observed effect size. The funnel plot generated *via* Duval and Tweedie's trim and fill procedure indicated the absence of studies with a high prevalence if the balance was to be restored (Figure 38).

## DISCUSSION

Viral hepatitis is major public health problem in the world, especially in developing countries. To our knowledge, this review is the first systematic review and meta-analysis study that has attempted to thoroughly summarize evidence on the different types of viral hepatitis infection in Somalia. The aim of this review was to understand the burden of viral hepatitis, especially HBV and HCV, in Somalia and to inform public health practitioners, researchers and policy makers in the


development of national strategies for the prevention and control of viral hepatitis. The findings of this review clearly show the burden of the viral hepatitis in the country, particularly HBV. The estimation prevalence of HBV in most African countries was 5%-20%<sup>[48]</sup>, and the WHO HBV endemic definition of HBsAg prevalence is 5%-7%<sup>[5]</sup>. In our review, we identified the prevalence of HBV in Somalia, as detected by HBsAg, to be high (18.9%). Somalia was categorized among countries with a high endemicity of HBV (prevalence  $\geq 8\%$ )<sup>[20,21]</sup>. The rate of global prevalence (3.61%) reported by Sweitzer and his colleagues, as well as the rate of 8.83% of the WHO African region, was significantly less than our study result<sup>[49]</sup>. In countries such the United States, Iran and Kosovo, the HBV prevalence rate has been estimated to be approximately < 0.27%, 2.14%, and 4.2%, respectively<sup>[50-52]</sup>. Therefore, this comparative information highlighted the extent of the HBV burden in Somalia. The pooled prevalence in this study was higher than that of a recent report from African countries, such as Ethiopia, Cameroon, Ghana, and Nigeria, the pooled prevalence of which was 7.4%, 11.2%, 12.3%, and 13.6%, respectively<sup>[48,53-55]</sup>.

Additionally, HBV infection among blood donors (19.1%)<sup>[33]</sup> and the pooled prevalence of pregnant women (20.5%) also remain high in Somalia, which justifies the requirements of a national HBV screening program for all pregnant women in antenatal clinics and a national policy to vaccinate all pregnant women and adults who test negative for HBV, so as to prevent and reduce the risk of HBV of mother-to-child transmission. We also need to establish a national blood policy throughout Somalia for the prevention or reduction of receiving contaminated blood, the occurrence of which remains high within this HBV population. According to studies of HBV infection among children, the prevalence (5.7%) is lower than that of other groups, while that




**Table 14** Studies on hepatitis C virus infection among patients with chronic liver disease, including hepatocellular carcinoma, in Somalia

Author	Year	Total	Cases	Total	Healthy	Setting	Population
Shire <i>et al</i> <sup>[36]</sup>	2012	30	22	30	8	United States	Immigrants
Kadle <i>et al</i> <sup>[38]</sup>	2012	156	30	156	126	Somalia	Local
Khadija <i>et al</i> <sup>[37]</sup>	2011	147	15	147	132	Somalia	Local
Bile <i>et al</i> <sup>[34]</sup>	1993	62	25	62	37	Somalia	Local
Aceti <i>et al</i> <sup>[42]</sup>	1993	110	28	110	82	Somalia	Local

**Figure 30** Forest plot of studies reporting chronic hepatitis C virus prevalence among patients with chronic liver disease, including hepatocellular carcinoma, in Somalia.**Table 15** Summary of studies on overall prevalence of hepatitis D viral infection of HBsAg-positive carriers in the Somali population in Somalia and Somali immigrants (1970-2016) n (%)

Author	Year	Total	Hepatitis D virus	Healthy	Setting	Population
Aceti <i>et al</i> <sup>[31]</sup>	1989	220	37 (16.8)	138	Somalia	Local
Bile <i>et al</i> <sup>[32]</sup>	1991	67	20 (30)	47	Somalia	Local
Aceti <i>et al</i> <sup>[30]</sup>	1991	52	26 (50)	26	Somalia	Local
Bile <i>et al</i> <sup>[34]</sup>	1993	29	10 (34.4)	19	Somalia	Local
Faustini <i>et al</i> <sup>[19]</sup>	1994	7	0 (0)	7	Italy	Immigrants
Total		375	93 (24.8)	237		

**Figure 31** Bias assessment plot of studies reporting chronic liver disease.

of Somali immigrants around the world and of chronic liver disease patients was 23% and 39.2%, respectively. These findings therefore demonstrate the true epidemiological burden of HBV in Somalia.

Regarding HCV, we found a high pooled prevalence of 4.84% in Somalia. This prevalence rate is considerably

higher than the reported pooled prevalence in Somalia of 0.9% by Chaabna *et al*<sup>[56]</sup> and 1.5% by Karoney *et al*<sup>[57]</sup>. Meanwhile, in Djibouti and Sudan, the pooled prevalence rate was 0.3% and 1.0%, respectively<sup>[56]</sup>. We also found close prevalence rates in our study and found that the pooled prevalence rate in Ethiopia, Ghana, and the Democratic Republic of the Congo was 3.1%<sup>[48]</sup>, 3.0%<sup>[58]</sup> and 2.9%<sup>[59]</sup>, respectively, but the overall prevalence rate is 3.0% in all sub-Saharan African countries<sup>[60]</sup>. However, in Cameroon, the estimated prevalence of anti-HCV is 11.6%<sup>[61]</sup> and the reported prevalence rate for Egypt is 14.7%<sup>[61,62]</sup>; thus, the estimated prevalence of anti-HCV is much lower in Somalia than in these two countries. The pooled prevalence rate of sub-groups examined in this study, such as blood donors, Somali immigrants, risk groups, children and patients with chronic liver disease, was 0.8%, 3.81%, 2.43%, 1.37%, and 29.82%, respectively. These findings may emphasize that the level of chronic HCV in Somalia may be significantly high. Genotypes commonly found in Africa are 1, 4 and 5<sup>[57]</sup>. Therefore, in this review, we also present the most frequent genotype among the Somali population, which is genotype 4. Most African countries, such as Uganda, Sudan, Rwanda, Burundi, Cameroon and Egypt, also show a predominance of HCV genotype 4<sup>[57]</sup>. The next most prevalent genotype in our country is genotype 3, which is also similar to our neighboring countries, such as Ethiopia, Kenya, and Eritrea<sup>[57]</sup>. Meanwhile, the next most prevalent genotype in Somalia is genotype 1.

The number of studies of HDV in Somalia among Somali patients and chronic liver disease patients was very low, but the data available thus far demonstrated a considerable prevalence of HDV infection. Our results




Figure 32 Forest plot of studies reporting hepatitis D virus infection prevalence in Somalia.

Table 16 Studies among patients with chronic liver disease (including hepatocellular carcinoma) in Somalia  $n$  (%)

Author	Year	Total	Hepatitis D virus	Total	Healthy	Setting	Population
Bile <i>et al</i> <sup>[34]</sup>	1993	23	9 (39.1)	23	14	Somalia	Local
Bile <i>et al</i> <sup>[32]</sup>	1991	44	17 (38.6)	44	27	Somalia	Local
Aceti <i>et al</i> <sup>[30]</sup>	1991	52	26 (50)	52	26	Somalia	Local
Total		119	52 (43.6)	119	67		




Figure 33 Bias assessment plot of studies reporting of hepatitis D virus infection in Somalia.



showed that the pooled prevalence rate of HDV in Somalia was 28.99%. The pooled prevalence of Sudan and Uganda, which are other sub-Saharan African countries, was 27.8%<sup>[63]</sup> and 30.6%<sup>[63]</sup>, respectively, and the pooled prevalence rate of the EMRO region is 14.74%<sup>[64]</sup>. Therefore, our result is similar to that of other countries in our region. The worldwide estimation rate also indicated that 5% of HBsAg carriers were infected with HDV<sup>[1]</sup>. The pooled prevalence rate in this review for patients with chronic liver disease, including hepatocellular carcinoma and positivity for anti-HDV, was shown to be 43.77%, and this result is similar to the prevalence rates among patients in the EMRO region with chronic liver disease (chronic hepatitis disease 27.8% and cirrhosis/hepatocellular carcinoma 36.57%)<sup>[64]</sup>. However, the previous pooled estimation in Somalia showed a similar prevalence rate as chronic hepatitis disease (47.36%) and cirrhosis/hepatocellular carcinoma (33.20%)<sup>[64]</sup>. Other countries in the EMRO region also showed results similar to our findings. For

example, the prevalence of chronic hepatitis disease and cirrhosis/hepatocellular carcinoma in Pakistan was 37.38% and 53.77%, respectively, and the prevalence of the same conditions in Egypt was 24.37% and 29.6%, respectively<sup>[64]</sup>. These results indicate that HDV infection is endemic in Somalia.

Although the number of prevalence studies of hepatitis A and E virus infections among Somali patients were also very low in Somalia due to the lack of government support in the country in the last three decades, the available data for HAV and HEV still illustrated a high prevalence rate in the country. These viruses were transmitted through the fecal-oral route, and many environmental and socio-economic factors promoted the transmission routes. This systematic review showed a high pooled prevalence rate of HAV, which is 90.2%, and this prevalence rate was close to the individual estimation rates of old reports in Somalia<sup>[15,16,18,19]</sup>. We also indicate that the estimates of HEV infection in previous reports in Somalia may quite high, ranging between 4.5% and 88.2%<sup>[45-47]</sup>, which is similar to estimates for the Ghanaian population of anti-HEV, with their range of 5.8% and 71.55%<sup>[65]</sup>. However, this review showed a pooled prevalence rate of HEV of 46.86%, which is also high among Somali patients. HEV infection in Africa is widespread and is predicted to be a hazard to numerous lives, in particular for pregnant women and their fetuses<sup>[2]</sup>. Serological studies from Egypt have shown that the seroprevalence of anti-HEV can reach close to 100% in the general population<sup>[2]</sup>. Another serological study from Ghana showed that anti-HEV can also reach close to 50%, and a study in Nigeria showed nearly 100% of anti-HEV in the general population<sup>[2]</sup>. Meanwhile, a number of large outbreaks occurred and were reported at different times in many countries in Africa, such as Kenya, Sudan, Ethiopia, Somalia, Uganda and Chad<sup>[2,45]</sup>. The outbreak that occurred in Somalia showed a range of 77.8% to 94.0% among three villages in the country<sup>[45]</sup>. A survey from 142 villages showed 11413 jaundice patients in a population of 245312 individuals, of whom 346 persons died, with a corresponding attack rate of 4.6% and a case fatality rate of 3.0%<sup>[46]</sup>. An older report demonstrated that the attack rate was higher with increasing age, from 5%, 13% and 20% for the

**Table 17** Studies on hepatitis E virus in Somalia *n* (%)

Author	Year	Total	Hepatitis E virus	Healthy	Setting	Population
Burans <i>et al</i> <sup>[47]</sup>	1994	31	20 (64.5)	11	Somalia	Local
Mushahwar <i>et al</i> <sup>[45]</sup>	1993	111	5 (4.5)	106	Somalia	Local
Bile <i>et al</i> <sup>[46]</sup>	1994	145	128 (88.2)	17	Somalia	Local
Total		287	153 (53.3)	134		

**Figure 34** Forest plot of hepatitis D virus infection prevalence rates among patients with chronic liver disease in Somalia.**Figure 35** Bias assessment plot of studies reporting of hepatitis D virus infection among chronic liver disease in Somalia.

age groups of 1-4 years, 5-15 years and above 15 years old, respectively. Meanwhile, pregnant women had a high fatality rate, estimated to be 13.8%<sup>[46]</sup>, in comparison to the rates in other African countries, such as Ghana, (66.7%)<sup>[65]</sup>, Sudan (31.1%)<sup>[66]</sup> and the Central African Republic (14.3%)<sup>[67]</sup>. However, the fatality rate of pregnant women was higher than that of our older report in the country<sup>[46]</sup>. This systematic review and meta-analysis study indicates that the two viruses of HAV and HEV are endemic in the country.

This systematic review and meta-analysis study had several limitations. The main limitation is that to the best of our knowledge, this review is the first that focuses on all types of viral hepatitis infection in Somalia. The second limitation of this review is the relatively small number of studies identified on all types of viral hepatitis in the country before and after the civil war. This lack of studies highlights the fact that research and knowledge of the viral hepatitis infection burden in Somalia are less developed than in other African and Arab countries. The third limitation is that the majority of our studies involved samples from only four or five regions of Somalia, especially in South Somalia, and samples of Somali immigrants around the world. Subsequently, we did not have samples

from the southwestern, central, and northern regions of the country, such as Gedo, Bakol, Hiran, Galgaduud, Mudug, Nugal, East, Sol, Sanag, Togder, Northwest and Awdal regions, which were included in the studies reviewed, while most studies were completed before the government collapsed, and few studies were done during and after the civil war.

In conclusions, although this systematic review and meta-analysis study is based on a limited number of studies and a small sample size, the study effects were a potential limitation of the review. However, the results of this review indicate that all types of viral hepatitis infection are common among Somalis, particularly HBV, which is more prevalent and endemic. This study has also documented a high prevalence rate of HCV infection in Somalia. The prevalence of these viruses could be understood as one of the nation's public health problems, and urgent public health interventions are needed to reduce the infection rate of the country, because these preventable diseases, particularly HBV and HAV, can be eradicated by vaccination. Preventive measures include increasing awareness and knowledge about the transmission of all forms of viral hepatitis, screening all donated blood, targeting high-risk groups, providing treatment for affected persons, and generally improving




Figure 36 Forest plot of studies reporting hepatitis E virus infection prevalence in Somalia.




Figure 37 Funnel plot of studies reporting hepatitis E virus infection prevalence in Somalia.




Figure 38 Bias assessment plot of studies reporting hepatitis E virus prevalence in Somalia.

the sanitary and living conditions. Other efforts needed to reduce the burden of the disease and strengthen the Somali health system include the establishment of a national blood policy and prevention and control viral hepatitis policy. Additionally, further studies are needed to fully understand the population factors underlying the high prevalence of all forms viral hepatitis, particularly in underrepresented regions and among adults, children, blood donors and high-risk groups, to offer better perspectives on all forms of the viral hepatitis burden in Somalia. The generation of up-to-date data is recommended, especially regarding the magnitude of HAV, HCV, and their genotypes, HDV and HEV.

## ARTICLE HIGHLIGHTS

### Research background

Viral hepatitis is a major public health problem affecting several hundred million people globally. The most common types of viral hepatitis are six distinct

types (hepatitis A, B, C, D, E, G viruses), and they may present in acute form or chronic form causes substantial morbidity and mortality (including chronic hepatitis, cirrhosis and hepatocellular carcinoma).

### Research motivation

In the field of viral hepatitis, there is a lack of researches last three decades in country and all articles about viral hepatitis published and unpublished are scattered. In this systematic review and meta-analysis the authors aim to provide a clear understanding of viral hepatitis epidemiology and their clinical burdens in Somalia according to the related documents published and unpublished articles during the last decades.

### Research objectives

The main objective of this study is to determine the prevalence of all viral hepatitis in Somalia especially hepatitis B virus (HBV) and hepatitis C virus (HCV), and to inform public health practitioners, researchers and policy makers, and to be a baseline data for future Hepatology researches in the country.

### Research methods

A systematic review and meta-analysis was conducted as Preferred Reporting Items for Systematic Reviews and Meta-Analyses guidelines. A comprehensive

literature search of published studies on viral hepatitis was performed from 1977-2016 in PubMed, Google Scholar, Science Direct, World Health Organization African Index Medicus and the Africa Journals Online databases, as well as on the Ministry of Health website. We also captured unpublished articles that were not available on online systems.

### Research results

Twenty-nine studies from Somalia and Somali immigrants (United Kingdom, United States, Italy, Libya) with a combined sample size for each type of viral hepatitis were analyzed. The overall pooled prevalence rate of hepatitis A virus was 90.2%. The overall pooled prevalence of HBV was 18.9%. The overall pooled prevalence of HCV was estimated as 4.84%. The overall pooled prevalence of hepatitis D virus was 28.99%. The overall pooled prevalence of hepatitis E virus was 46.86%.

### Research conclusions

This study demonstrates a high prevalence of all forms of viral hepatitis in Somalia and it also indicates that chronic HBV was the commonest cause of chronic liver disease.

### Research perspectives

Viral hepatitis in Somalia demonstrated a high rate in its all types of hepatitis especially hepatitis B virus and C, while hepatitis B determined the common cause of chronic liver disease in Somalia. According to this systematic review and meta-analysis, further studies are needed in order to search out data about viral hepatitis in different regions of the country, risk factors and its complications.

## REFERENCES

- 1 **World Health Organization.** Global Hepatitis Report 2017 Accessed March 16, 2018. Available from: URL: [apps.who.int/iris/bitstream/10665/255016/1/9789241565455-eng.pdf](https://apps.who.int/iris/bitstream/10665/255016/1/9789241565455-eng.pdf)
- 2 **Kim JH, Nelson KE, Panzner U, Kasture Y, Labrique AB, Wierzb TF.** A systematic review of the epidemiology of hepatitis E virus in Africa. *BMC Infect Dis* 2014; **14**: 308 [PMID: 24902967 DOI: 10.1186/1471-2334-14-308]
- 3 **Jacobsen KH, Wiersma ST.** Hepatitis A virus seroprevalence by age and world region, 1990 and 2005. *Vaccine* 2010; **28**: 6653-6657 [PMID: 20723630 DOI: 10.1016/j.vaccine.2010.08.037]
- 4 **Hassan-Kadle MA.** The Diagnostic Challenges, Possible Etiologies and Lack of Researches of Hepatocellular Carcinoma in Somalia. *Open J Gastroenterol* 2017; **7**: 115 [DOI: 10.4236/ojgas.2017.73013]
- 5 **World Health Organization.** Global policy report on the prevention and control of viral hepatitis, 2013. Accessed March 16, 2018. Available from: URL: [apps.who.int/iris/bitstream/10665/85397/1/9789241564632\\_eng.pdf](https://apps.who.int/iris/bitstream/10665/85397/1/9789241564632_eng.pdf)
- 6 **Central Intelligence Agency.** The world fact book. Accessed March 16, 2018. Available from: URL: (<https://www.cia.gov/library/publications/the-world-factbook/geos/so.html>)
- 7 **World Bank.** Somalia overview. Accessed March 16, 2018. Available from: URL: <http://www.worldbank.org/en/country/somalia/overview>
- 8 **African Development Bank Group.** Somalia Country Brief 2013-2015. Accessed March 16, 2018. Available from: URL: <http://www.afdb.org/fileadmin/uploads/afdb/Documents/Project-and-Operations/2013-2015%20-%20Somalia%20-%20Country%20Brief.pdf>
- 9 **Moher D, Liberati A, Tetzlaff J, Altman DG; PRISMA Group.** Reprint--preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *Phys Ther* 2009; **89**: 873-880 [PMID: 19723669 DOI: 10.1016/j.jclinepi.2009.06.005]
- 10 **Viechtbauer W.** Package 'metafor'. Meta-Analysis Package for R. Accessed March 16, 2018. Available from: URL: <https://cran.r-project.org/web/packages/metafor/metafor.pdf>
- 11 **Schwarzer G.** Package 'meta'. General Package for Meta-Analysis. Accessed March 16, 2018. Available from: URL: <https://cran.r-project.org/web/packages/meta/meta.pdf>
- 12 **Normand SL.** Meta-analysis: formulating, evaluating, combining, and reporting. *Stat Med* 1999; **18**: 321-359 [PMID: 10070677 DOI: 10.1002/(SICI)1097-0258(19990215)18:3<321::AID-SIM28>3.0.CO;2-P]
- 13 **Duval S, Tweedie R.** Trim and fill: A simple funnel-plot-based method of testing and adjusting for publication bias in meta-analysis. *Biometrics* 2000; **56**: 455-463 [PMID: 10877304 DOI: 10.1111/j.0006-341X.2000.00455.x]
- 14 **Egger M, Davey Smith G, Schneider M, Minder C.** Bias in meta-analysis detected by a simple, graphical test. *BMJ* 1997; **315**: 629-634 [PMID: 9310563 DOI: 10.1136/bmj.315.7109.629]
- 15 **Mohamud KB, Aceti A, Mohamed OM, Pennica A, Maalin KA, Biondi R, Hagi MA, Quaranta G, Paparo SB, Celestino D.** [The circulation of the hepatitis A and B viruses in the Somali population]. *Ann Ital Med Int* 1992; **7**: 78-83 [PMID: 1334688]
- 16 **Bile K, Mohamud O, Aden C, Isse A, Norder H, Nilsson L, Magnus L.** The risk for hepatitis A, B, and C at two institutions for children in Somalia with different socioeconomic conditions. *Am J Trop Med Hyg* 1992; **47**: 357-364 [PMID: 1524149 DOI: 10.4269/ajtmh.1992.47.357]
- 17 **Bile K, Aceti A, Hagi MA, Paparo BS, Pennica A, Celestino D, Sebastiani A.** Primary hepatitis A virus infection in developing countries: decline of maternal antibodies and time of infection. *Boll Ist Sieroter Milan* 1986; **65**: 464-466 [PMID: 3034305]
- 18 **Sebastiani A, Aceti A, Russo V, Paparo BS, Pennica A, Bile K, Farah AA, Farah AH.** Age-specific prevalence of hepatitis A and hepatitis B virus infection in fluvial Somalian village. *Estratto dal* 1984; **30**: 17-23
- 19 **Faustini A, Franco E, Saitto C, Cauletti M, Zaratti L, Papini P, Ali' Ahmed S, Zampieri F, Ierussi A, Panà A.** Hepatitis A, B, C and D in a community in Italy of immigrants from NE Africa. *J Public Health Med* 1994; **16**: 71-78 [PMID: 8037956 DOI: 10.1093/oxfordjournals.pubmed.a042938]
- 20 **Averhoff F.** Hepatitis B. Accessed March 16, 2018. Available from: URL: <http://wwwnc.cdc.gov/travel/yellowbook/2016/infectious-diseases-related-to-travel/hepatitis-b>
- 21 **Howell J, Ladep NG, Lemoine M, Thurst MR, Taylor-Robinson SD.** Hepatitis B in Sub-Saharan Africa. *South Sudan Medical J* 2014; **7**: 59-61
- 22 **Delia S, Nuti M, Soro S.** [Relations between hepatitis B surface antigen and parasitic diseases: observations in patients with ancylostomiasis and schistosomiasis]. *Quad Sclavo Diagn* 1977; **13**: 238-243 [PMID: 594309]
- 23 **Nuti M, Abdullahi SE, Thamer G.** [E antigen (HBeAg) and surface antigen (HBsAg) in bladder schistosomiasis]. *Parassitologia* 1978; **20**: 153-159 [PMID: 553266]
- 24 **Nuti M, Tarabini G, Palermo P, Tarabini GL, Thamer G.** Leprosy and hepatitis B virus markers: incidence of HBsAg and HBeAg in Somalian patients. *Int J Lepr Other Mycobact Dis* 1979; **47**: 580-584 [PMID: 122628]
- 25 **Nuti M, Harare O, Thamer G.** The surface antigen (HBsAg) and the e-antigen (HBeAg) in Somalian patients with acute viral hepatitis. *Trans R Soc Trop Med Hyg* 1979; **73**: 185-187 [PMID: 473307 DOI: 10.1016/0035-9203(79)90207-4]
- 26 **Nuti M, Abdullahi Elmi S, Alario C.** [Diffusion of hepatitis B surface antigens in subjects with bladder schistosomiasis]. *Boll Ist Sieroter Milan* 1979; **58**: 220-223 [PMID: 518754]
- 27 **Sebastiani A, Aceti A, Paparo BS, Pennica A, Ilardi I, Bile K, Mohamud OM.** Hepatitis B virus circulation in three different villages of Somalia. *Trans R Soc Trop Med Hyg* 1985; **79**: 162-164 [PMID: 4002285 DOI: 10.1016/0035-9203(85)90323-2]
- 28 **Bile KM, Aden A, Lindberg G, Nilsson L, Lidman K, Norder H, Magnus L.** Epidemiology of hepatitis B in Somalia: inference from a cross-sectional survey of serological markers. *Trans R Soc Trop Med Hyg* 1987; **81**: 824-828 [PMID: 3502433 DOI: 10.1016/0035-9203(87)90043-5]
- 29 **Kolmos HJ, Zimakoff J.** [DASPID--an interdisciplinary Danish study group for the elucidation of peritonitis among peritoneal dialysis patients]. *Ugeskr Laeger* 1991; **153**: 1068 [PMID: 2024336]



- 30 **Aceti A**, Mohamed OM, Paparo BS, Mohamud OM, Quaranta G, Maalin KA, Sebastiani A. High prevalence of anti-hepatitis delta virus antibody in chronic liver disease in Somalia. *Trans R Soc Trop Med Hyg* 1991; **85**: 541-542 [PMID: 1755067 DOI: 10.1016/0035-9203(91)90249-X]
- 31 **Aceti A**, Paparo BS, Celestino D, Pennica A, Caferro M, Grilli A, Sebastiani A, Mohamud OM, Abdirahman M, Bile K. Sero-epidemiology of hepatitis delta virus infection in Somalia. *Trans R Soc Trop Med Hyg* 1989; **83**: 399-400 [PMID: 2617589 DOI: 10.1016/0035-9203(89)90516-6]
- 32 **Bile KM**, Abdirahman M, Aden C, Norder H, Magnius L, Lindberg G, Nilsson LH. Minor role of hepatitis B virus in the causation of chronic liver disease in Somalia indicated by a case-control study. *Trans R Soc Trop Med Hyg* 1991; **85**: 104-108 [PMID: 1712517 DOI: 10.1016/0035-9203(91)90177-Z]
- 33 **Nur YA**, Groen J, Elmi AM, Ott A, Osterhaus AD. Prevalence of serum antibodies against bloodborne and sexually transmitted agents in selected groups in Somalia. *Epidemiol Infect* 2000; **124**: 137-141 [PMID: 10722141 DOI: 10.1017/S0950268899003441]
- 34 **Bile K**, Aden C, Norder H, Magnius L, Lindberg G, Nilsson L. Important role of hepatitis C virus infection as a cause of chronic liver disease in Somalia. *Scand J Infect Dis* 1993; **25**: 559-564 [PMID: 7506842 DOI: 10.3109/00365549309008543]
- 35 **Aweis D**, Brabin BJ, Beeching NJ, Bunn JE, Cooper C, Gardner K, Iriyagolle C, Hart CA. Hepatitis B prevalence and risk factors for HBsAg carriage amongst Somali households in Liverpool. *Commun Dis Public Health* 2001; **4**: 247-252 [PMID: 12109390]
- 36 **Shire AM**, Sandhu DS, Kaiya JK, Oseini AM, Yang JD, Chaiteerakij R, Mettler TA, Giana NH, Roberts RO, Therneau TM, Petersen GM, Knutson KL, Roberts LR. Viral hepatitis among Somali immigrants in Minnesota: association of hepatitis C with hepatocellular carcinoma. *Mayo Clin Proc* 2012; **87**: 17-24 [PMID: 22212964 DOI: 10.1016/j.mayocp.2011.08.001]
- 37 **Olow KH**, Warsame FS. Prevalence of Liver Cirrhosis Post - Hepatitis B and C Among Adults in Mogadishu Hospitals. 2011
- 38 **Kadle M**, Hassan AM, Yasin AM, Sheikh AH, Omar MS, Sayid AM. Frequency of Hepatocellular Carcinoma and Its Associated of Hepatitis B and C in Patients Attending Mogadishu Hospitals. Accessed March 16, 2018. Available from: URL: [https://www.researchgate.net/publication/309564485\\_Frequency\\_of\\_Hepatocellular\\_Carcinoma\\_and\\_its\\_associated\\_Hepatitis\\_B\\_and\\_C\\_in\\_Patients\\_attending\\_Mogadishu\\_hospitals](https://www.researchgate.net/publication/309564485_Frequency_of_Hepatocellular_Carcinoma_and_its_associated_Hepatitis_B_and_C_in_Patients_attending_Mogadishu_hospitals)
- 39 **Padovese V**, Egidio AM, Melillo TF, Farrugia B, Carabot P, Didero D, Costanzo G, Mirisola C. Prevalence of latent tuberculosis, syphilis, hepatitis B and C among asylum seekers in Malta. *J Public Health (Oxf)* 2014; **36**: 22-27 [PMID: 23559596 DOI: 10.1093/pubmed/fdt036]
- 40 **Bile K**, Abdirahman M, Mohamud O, Aden C, Isse A, Nilsson L, Norder H, Magnius L. Late seroconversion to hepatitis B in a Somali village indicates the important role of venereal transmission. *J Trop Med Hyg* 1991; **94**: 367-373 [PMID: 1758006]
- 41 **Watts DM**, Corwin AL, Omar MA, Hyams KC. Low risk of sexual transmission of hepatitis C virus in Somalia. *Trans R Soc Trop Med Hyg* 1994; **88**: 55-56 [PMID: 8154002 DOI: 10.1016/0035-9203(94)90495-2]
- 42 **Aceti A**, Taliani G, Bruni R, Sharif OS, Moallin KA, Celestino D, Quaranta G, Sebastiani A. Hepatitis C virus infection in chronic liver disease in Somalia. *Am J Trop Med Hyg* 1993; **48**: 581-584 [PMID: 7683179 DOI: 10.4269/ajtmh.1993.48.581]
- 43 **Daw MA**, El-Bouzedi A, Ahmed MO, Dau AA, Agnan MM. In association with the Libyan Study Group of Hepatitis & HIV. Epidemiology of hepatitis C virus and genotype distribution in immigrants crossing to Europe from North and sub-Saharan Africa. *Travel Med Infect Dis* 2016; **14**: 517-526 [PMID: 27502972 DOI: 10.1016/j.tmaid.2016.05.020]
- 44 **Bradley DW**. Enterically-transmitted non-A, non-B hepatitis. *Br Med Bull* 1990; **46**: 442-461 [PMID: 2116213 DOI: 10.1093/oxfordjournals.bmb.a072409]
- 45 **Mushahwar IK**, Dawson GJ, Bile KM, Magnius LO. Serological studies of an enterically transmitted non-A, non-B hepatitis in Somalia. *J Med Virol* 1993; **40**: 218-221 [PMID: 8355020 DOI: 10.1002/jmv.1890400309]
- 46 **Bile K**, Isse A, Mohamud O, Allebeck P, Nilsson L, Norder H, Mushahwar IK, Magnius LO. Contrasting roles of rivers and wells as sources of drinking water on attack and fatality rates in a hepatitis E epidemic in Somalia. *Am J Trop Med Hyg* 1994; **51**: 466-474 [PMID: 7943574 DOI: 10.4269/ajtmh.1994.51.466]
- 47 **Burans JP**, Sharp T, Wallace M, Longer C, Thornton S, Batchelor R, Clemens V, Hyams KC. Threat of hepatitis E virus infection in Somalia during Operation Restore Hope. *Clin Infect Dis* 1994; **18**: 100-102 [PMID: 8054416 DOI: 10.1093/clinids/18.1.100]
- 48 **Belyhun Y**, Maier M, Mulu A, Diro E, Liebert UG. Hepatitis viruses in Ethiopia: a systematic review and meta-analysis. *BMC Infect Dis* 2016; **16**: 761 [PMID: 27993129 DOI: 10.1186/s12879-016-2090-1]
- 49 **Schweitzer A**, Horn J, Mikolajczyk RT, Krause G, Ott JJ. Estimations of worldwide prevalence of chronic hepatitis B virus infection: a systematic review of data published between 1965 and 2013. *Lancet* 2015; **386**: 1546-1555 [PMID: 26231459 DOI: 10.1016/S0140-6736(15)61412-X]
- 50 **Wasley A**, Kruszon-Moran D, Kuhnert W, Simard EP, Finelli L, McQuillan G, Bell B. The prevalence of hepatitis B virus infection in the United States in the era of vaccination. *J Infect Dis* 2010; **202**: 192-201 [PMID: 20533878 DOI: 10.1086/653622]
- 51 **Alavian SM**, Hajarizadeh B, Ahmadzad-Asl M, Kabir A, Bagheri-Lankarani K. Hepatitis B Virus infection in Iran: A systematic review. *Hepatitis monthly* 2008; **8**: 281-294
- 52 **Fejza H**, Telaku S. Prevalence of HBV and HCV among blood donors in Kosovo. *Virol J* 2009; **6**: 21 [PMID: 19216773 DOI: 10.1186/1743-422X-6-21]
- 53 **Bigna JJ**, Amougou MA, Asangbeh SL, Kenne AM, Noumegni SRN, Ngo-Malabo ET, Noubiap JJ. Seroprevalence of hepatitis B virus infection in Cameroon: a systematic review and meta-analysis. *BMJ Open* 2017; **7**: e015298 [PMID: 28667212 DOI: 10.1136/bmjopen-2016-015298]
- 54 **Ofori-Asenso R**, Agyeman AA. Hepatitis B in Ghana: a systematic review & meta-analysis of prevalence studies (1995-2015). *BMC Infect Dis* 2016; **16**: 130 [PMID: 26987556 DOI: 10.1186/s12879-016-1467-5]
- 55 **Musa BM**, Bussell S, Borodo MM, Samaila AA, Femi OL. Prevalence of hepatitis B virus infection in Nigeria, 2000-2013: a systematic review and meta-analysis. *Niger J Clin Pract* 2015; **18**: 163-172 [PMID: 25665986 DOI: 10.4103/1119-3077.151035]
- 56 **Chaabna K**, Kouyoumjian SP, Abu-Raddad LJ. Hepatitis C Virus Epidemiology in Djibouti, Somalia, Sudan, and Yemen: Systematic Review and Meta-Analysis. *PLoS One* 2016; **11**: e0149966 [PMID: 26900839 DOI: 10.1371/journal.pone.0149966]
- 57 **Karone MJ**, Siika AM. Hepatitis C virus (HCV) infection in Africa: a review. *Pan Afr Med J* 2013; **14**: 44 [PMID: 23560127 DOI: 10.11604/pamj.2013.14.44.2199]
- 58 **Agyeman AA**, Ofori-Asenso R, Mprah A, Ashiagbor G. Epidemiology of hepatitis C virus in Ghana: a systematic review and meta-analysis. *BMC Infect Dis* 2016; **16**: 391 [PMID: 27507267 DOI: 10.1186/s12879-016-1708-7]
- 59 **Muzembo BA**, Akita T, Matsuoka T, Tanaka J. Systematic review and meta-analysis of hepatitis C virus infection in the Democratic Republic of Congo. *Public Health* 2016; **139**: 13-21 [PMID: 27450441 DOI: 10.1016/j.puhe.2016.06.017]
- 60 **Madhava V**, Burgess C, Drucker E. Epidemiology of chronic hepatitis C virus infection in sub-Saharan Africa. *Lancet Infect Dis* 2002; **2**: 293-302 [PMID: 12062995 DOI: 10.1016/S1473-3099(02)00264-5]
- 61 **Gower E**, Estes C, Blach S, Razavi-Shearer K, Razavi H. Global epidemiology and genotype distribution of the hepatitis C virus infection. *J Hepatol* 2014; **61**: S45-S57 [PMID: 25086286 DOI: 10.1016/j.jhep.2014.07.027]
- 62 **Mohamoud YA**, Mumtaz GR, Riome S, Miller D, Abu-Raddad LJ. The epidemiology of hepatitis C virus in Egypt: a systematic review and data synthesis. *BMC Infect Dis* 2013; **13**: 288 [PMID: 23799878 DOI: 10.1186/1471-2334-13-288]

- 63 **Mudawi HM.** Epidemiology of viral hepatitis in Sudan. *Clin Exp Gastroenterol* 2008; **1**: 9-13 [PMID: 21677820 DOI: 10.2147/CEG.S3887]
- 64 **Amini N,** Alavian SM, Kabir A, Aalaei-Andabili SH, Saiedi Hosseini SY, Rizzetto M. Prevalence of hepatitis d in the eastern mediterranean region: systematic review and meta analysis. *Hepat Mon* 2013; **13**: e8210 [PMID: 23554822 DOI: 10.5812/hepatmon.8210]
- 65 **Ofori-Asenso R,** Agyeman AA. Hepatitis E infection among Ghanaians: a systematic review. *Infect Dis Poverty* 2017; **6**: 29 [PMID: 28162095 DOI: 10.1186/s40249-017-0239-9]
- 66 **Boccia D,** Guthmann JP, Klovstad H, Hamid N, Tatay M, Ciglenecki I, Nizou JY, Nicand E, Guerin PJ. High mortality associated with an outbreak of hepatitis E among displaced persons in Darfur, Sudan. *Clin Infect Dis* 2006; **42**: 1679-1684 [PMID: 16705571 DOI: 10.1086/504322]
- 67 **Goumba CM,** Yandoko-Nakouné ER, Kommas NP. A fatal case of acute hepatitis E among pregnant women, Central African Republic. *BMC Res Notes* 2010; **3**: 103 [PMID: 20398305 DOI: 10.1186/1756-0500-3-103]

**P-Reviewer:** Esmat SM, Pokorska-Spiwak M, Wang K  
**S-Editor:** Wang JL **L-Editor:** A **E-Editor:** Bian YN





Published by **Baishideng Publishing Group Inc**  
7901 Stoneridge Drive, Suite 501, Pleasanton, CA 94588, USA  
Telephone: +1-925-223-8242  
Fax: +1-925-223-8243  
E-mail: [bpgoffice@wjgnet.com](mailto:bpgoffice@wjgnet.com)  
Help Desk: <http://www.f6publishing.com/helpdesk>  
<http://www.wjgnet.com>



ISSN 1007-9327

