

World Journal of *Gastroenterology*

World J Gastroenterol 2018 August 28; 24(32): 3567-3676



**REVIEW**

- 3567 Positioning of old and new biologicals and small molecules in the treatment of inflammatory bowel diseases
Reinglas J, Gonczi L, Kurt Z, Bessissow T, Lakatos PL
- 3583 Prognostic significance of tumor immune microenvironment and immunotherapy: Novel insights and future perspectives in gastric cancer
Lazăr DC, Avram MF, Romoșan I, Cornianu M, Tăban S, Goldiș A
- 3617 *Helicobacter pylori* infection and liver diseases: Epidemiology and insights into pathogenesis
Okushin K, Tsutsumi T, Ikeuchi K, Kado A, Enooku K, Fujinaga H, Moriya K, Yotsuyanagi H, Koike K

MINIREVIEWS

- 3626 Expansion of the hepatocellular carcinoma Milan criteria in liver transplantation: Future directions
Pavel MC, Fuster J
- 3637 Diagnosis and management of fibromuscular dysplasia and segmental arterial mediolysis in gastroenterology field: A mini-review
Ko M, Kamimura K, Ogawa K, Tominaga K, Sakamaki A, Kamimura H, Abe S, Mizuno K, Terai S

ORIGINAL ARTICLE**Basic Study**

- 3650 Abnormal expression of HMGB-3 is significantly associated with malignant transformation of hepatocytes
Zheng WJ, Yao M, Fang M, Wang L, Dong ZZ, Yao DF

Retrospective Cohort Study

- 3663 C-peptide as a key risk factor for non-alcoholic fatty liver disease in the United States population
Atsawarungruangkit A, Chenbhanich J, Dickstein G

Clinical Trials Study

- 3671 Vascular anatomy of inferior mesenteric artery in laparoscopic radical resection with the preservation of left colic artery for rectal cancer
Wang KX, Cheng ZQ, Liu Z, Wang XY, Bi DS

ABOUT COVER

Editorial board member of *World Journal of Gastroenterology*, Fabio Grizzi, PhD, Assistant Professor, Department of Immunology and Inflammation, Humanitas Clinical and Research Hospital, Rozzano 20089, Italy

AIMS AND SCOPE

World Journal of Gastroenterology (*World J Gastroenterol*, *WJG*, print ISSN 1007-9327, online ISSN 2219-2840, DOI: 10.3748) is a peer-reviewed open access journal. *WJG* was established on October 1, 1995. It is published weekly on the 7th, 14th, 21st, and 28th each month. The *WJG* Editorial Board consists of 642 experts in gastroenterology and hepatology from 59 countries.

The primary task of *WJG* is to rapidly publish high-quality original articles, reviews, and commentaries in the fields of gastroenterology, hepatology, gastrointestinal endoscopy, gastrointestinal surgery, hepatobiliary surgery, gastrointestinal oncology, gastrointestinal radiation oncology, gastrointestinal imaging, gastrointestinal interventional therapy, gastrointestinal infectious diseases, gastrointestinal pharmacology, gastrointestinal pathophysiology, gastrointestinal pathology, evidence-based medicine in gastroenterology, pancreatology, gastrointestinal laboratory medicine, gastrointestinal molecular biology, gastrointestinal immunology, gastrointestinal microbiology, gastrointestinal genetics, gastrointestinal translational medicine, gastrointestinal diagnostics, and gastrointestinal therapeutics. *WJG* is dedicated to become an influential and prestigious journal in gastroenterology and hepatology, to promote the development of above disciplines, and to improve the diagnostic and therapeutic skill and expertise of clinicians.

INDEXING/ABSTRACTING

World Journal of Gastroenterology (*WJG*) is now indexed in Current Contents®/Clinical Medicine, Science Citation Index Expanded (also known as SciSearch®), Journal Citation Reports®, Index Medicus, MEDLINE, PubMed, PubMed Central and Directory of Open Access Journals. The 2018 edition of Journal Citation Reports® cites the 2017 impact factor for *WJG* as 3.300 (5-year impact factor: 3.387), ranking *WJG* as 35th among 80 journals in gastroenterology and hepatology (quartile in category Q2).

EDITORS FOR THIS ISSUE

Responsible Assistant Editor: *Xiang Li*
Responsible Electronic Editor: *Shu-Yu Yin*
Proofing Editor-in-Chief: *Lian-Sheng Ma*

Responsible Science Editor: *Rao-Yu Ma*
Proofing Editorial Office Director: *Ze-Mao Gong*

NAME OF JOURNAL
World Journal of Gastroenterology

ISSN
ISSN 1007-9327 (print)
ISSN 2219-2840 (online)

LAUNCH DATE
October 1, 1995

FREQUENCY
Weekly

EDITORS-IN-CHIEF
Andrzej S Tarnawski, MD, PhD, DSc (Med),
Professor of Medicine, Chief Gastroenterology, VA
Long Beach Health Care System, University of California, Irvine, CA, 5901 E. Seventh Str., Long Beach, CA 90822, United States

EDITORIAL BOARD MEMBERS
All editorial board members resources online at <http://www.wjgnet.com/1007-9327/editorialboard.htm>

EDITORIAL OFFICE
Ze-Mao Gong, Director
World Journal of Gastroenterology
Baishideng Publishing Group Inc
7901 Stoneridge Drive, Suite 501,
Pleasanton, CA 94588, USA
Telephone: +1-925-2238242
Fax: +1-925-2238243
E-mail: editorialoffice@wjgnet.com
Help Desk: <http://www.f6publishing.com/helpdesk>
<http://www.wjgnet.com>

PUBLISHER
Baishideng Publishing Group Inc
7901 Stoneridge Drive, Suite 501,
Pleasanton, CA 94588, USA
Telephone: +1-925-2238242
Fax: +1-925-2238243
E-mail: bpgoffice@wjgnet.com
Help Desk: <http://www.f6publishing.com/helpdesk>
<http://www.wjgnet.com>

PUBLICATION DATE
August 28, 2018

COPYRIGHT
© 2018 Baishideng Publishing Group Inc. Articles published by this Open-Access journal are distributed under the terms of the Creative Commons Attribution Non-commercial License, which permits use, distribution, and reproduction in any medium, provided the original work is properly cited, the use is non commercial and is otherwise in compliance with the license.

SPECIAL STATEMENT
All articles published in journals owned by the Baishideng Publishing Group (BPG) represent the views and opinions of their authors, and not the views, opinions or policies of the BPG, except where otherwise explicitly indicated.

INSTRUCTIONS TO AUTHORS
Full instructions are available online at <http://www.wjgnet.com/bpg/gerinfo/204>

ONLINE SUBMISSION
<http://www.f6publishing.com>

Retrospective Cohort Study

C-peptide as a key risk factor for non-alcoholic fatty liver disease in the United States population

Amporn Atsawarungrangkit, Jirat Chenbhanich, George Dickstein

Amporn Atsawarungrangkit, Jirat Chenbhanich, George Dickstein, Department of Medicine, MetroWest Medical Center, Framingham, MA 01702, United States

Amporn Atsawarungrangkit, Jirat Chenbhanich, George Dickstein, Department of Medicine, Tufts University School of Medicine, Boston, MA 02111, United States

ORCID number: Amporn Atsawarungrangkit (0000-0003-0622-6839); Jirat Chenbhanich (0000-0002-2274-5965); George Dickstein (0000-0003-0100-5498).

Author contributions: Atsawarungrangkit A contributed to the study concept and design, analysis and interpretation of data, and preparation of manuscript; Chenbhanich J and Dickstein G contributed to interpretation of data and preparation and manuscript; all authors read and approved the final manuscript.

Institutional review board statement: This study uses the publicly available data from the Third National Health and Nutrition Examination Survey (NHANES III), which is conducted by the National Center for Health Statistics (NCHS). The NHANES protocol was approved by the NCHS Research Ethics Review Board.

Informed consent statement: In NHANES III, the consent form was signed by participants in the survey.

Conflict-of-interest statement: No conflict of interest exists.

Open-Access: This article is an open-access article which was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

Manuscript source: Unsolicited manuscript

Correspondence to: Amporn Atsawarungrangkit, BPharm, MD, Doctor, Research Fellow, Department of Medicine,

MetroWest Medical Center, 115 Lincoln St., Framingham, MA 01702, United States. a.atsawarungrangkit@mwmc.com
Telephone: +1-857-3126114

Received: May 30, 2018

Peer-review started: May 31, 2018

First decision: July 4, 2018

Revised: July 6, 2018

Accepted: July 16, 2018

Article in press: July 16, 2018

Published online: August 28, 2018

Abstract

AIM

To determine whether fasting C-peptide is an independent predictor for non-alcoholic fatty liver disease (NAFLD) in United States population.

METHODS

Using the National Health and Nutrition Examination Survey (NHANES) 1988-1994, NAFLD participants aged 20 or greater without any other liver diseases were included in this study. Excessive alcohol intake is defined as > 2 drinks per day for males and > 1 drink per day for females. C-peptide and 27 other factors known to be associated with NAFLD (*e.g.*, age, gender, body mass index, waist circumference, race/ethnicity, liver chemistries, and other diabetes tests) were tested in both univariate and multivariate level using logistic regression with a *P*-value 0.05.

RESULTS

Of 18825 participants aged ≥ 20 , 3235 participants ($n = 3235$) met inclusion criteria. There were 23 factors associated with NAFLD by univariate analysis. 9 factors, ranked by the highest change in pseudo R^2 , were found to be significant predictors of NAFLD in multivariate model: waist circumference, fasting C-peptide, natural log of alanine aminotransferase (ALT), total protein, being

Mexican American, natural log of glycated hemoglobin, triglyceride level, being non-Hispanic white, and ferritin level.

CONCLUSION

Together with waist circumference and ALT, fasting C-peptide is among three most important predictors of NAFLD in United States population in the NHANES data set. Further study is needed to validate the clinical utility of fasting C-peptide in diagnosis or monitoring insulin resistance in NAFLD patients.

Key words: Insulin resistance; Fatty liver; Hepatosteatosis; Metabolic syndrome; C-peptide

© The Author(s) 2018. Published by Baishideng Publishing Group Inc. All rights reserved.

Core tip: Non-alcoholic fatty liver disease (NAFLD) is a growing global epidemic and associated with many conditions and factors, including insulin resistance. However, C-peptide has not been used in practice to assess insulin resistance in NAFLD patients. Using a large national dataset, we demonstrated that three most important risk factors for NAFLD are waist circumference, fasting C-peptide, and alanine aminotransferase, respectively. Such results revealed that C-peptide superior to measurement of fasting insulin levels and can potentially be used for screening or monitoring the degree of insulin resistance in NAFLD.

Atsawarungrangkit A, Chenbhanich J, Dickstein G. C-peptide as a key risk factor for non-alcoholic fatty liver disease in the United States population. *World J Gastroenterol* 2018; 24(32): 3663-3670 Available from: URL: <http://www.wjgnet.com/1007-9327/full/v24/i32/3663.htm> DOI: <http://dx.doi.org/10.3748/wjg.v24.i32.3663>

INTRODUCTION

Non-alcoholic fatty liver disease (NAFLD) is a condition in which hepatic steatosis exists in the absence of excessive alcohol consumption. NAFLD is the most common cause of chronic liver disease in the United States with estimated prevalence around 30%-40%^[1-3]. Given the epidemic of obesity, NAFLD is increasingly prevalent and challenging^[4,5]. NAFLD can progress to more severe liver diseases, such as non-alcoholic steatohepatitis (NASH), cirrhosis, and hepatocellular carcinoma.

Obesity and insulin resistance are among important risk factors for NAFLD^[6,7]. Many studies found that indicators of obesity [*i.e.*, body mass index (BMI) and waist circumference] and insulin resistance [*i.e.*, glycated hemoglobin (HbA1c), insulin level, fasting glucose, and diabetes mellitus] are independently associated with NAFLD and/or severity of liver fibrosis in NAFLD^[8-11]. C-peptide levels can be used to measure insulin secretion^[12]. However, there is limited evidence

of the association between NAFLD and C-peptide at the multivariate level^[13,14].

Both C-peptide and insulin are produced and released in equimolar amounts. C-peptide can therefore be used to assess endogenous insulin secretion. However, the level of C-peptide and insulin level in blood are typically different deriving from the differences in clearance mechanisms and half-life^[15]. In addition to diabetes and insulin resistance, C-peptide has been associated with many risk factors for NAFLD including cardiovascular diseases and metabolic syndrome^[16-18].

Therefore, our primary objective was to determine if fasting C-peptide is independently associated with NAFLD using multivariate analysis in the United States general population.

MATERIALS AND METHODS

Study population and study design

The Third National Health and Nutrition Examination Survey (NHANES III) is a probability sample of 39695 persons aged 2 mo and older representing the United States population and conducted by the National Center for Health Statistics (NCHS) to evaluate health and nutritional status^[19]. The survey collected multiple data sets, including demographic, interviews, physical examinations, and laboratory testing of biologic samples. NHANES III was conducted from 1988 to 1994. The NHANES protocol was approved by the NCHS Research Ethics Review Board.

There were 18825 persons aged 20 years or older in NHANES III that met inclusion criteria for this study. The exclusion criteria included: (1) Ungradable or missing ultrasound results for hepatic steatosis, (2) excessive alcohol consumption, (3) hepatitis B or hepatitis C infection (4) fasting period outside of 8-24 h (5) incomplete or missing data on physical examination and laboratory testing. Participants were divided into two groups: NAFLD participants (study group) and non-NAFLD participants (control group).

As presented in Table 1, we included 28 factors associated with NAFLD as independent variables in this study: Demographic (*i.e.*, age, gender, race/ethnicity), body measurement (*i.e.*, BMI and waist circumference), general biochemistry tests [*i.e.*, iron, total iron-binding capacity (TIBC), transferrin saturation, ferritin, cholesterol, triglyceride, HDL cholesterol, C-reactive protein, and uric acid], liver chemistry [aspartate aminotransferase (AST), Alanine aminotransferase (ALT), gamma glutamyl transferase (GGT), alkaline phosphatase (ALP), total bilirubin, total protein, and albumin], and diabetes testing profile [*i.e.*, HbA1c, fasting plasma glucose (FPG), fasting C-peptide, and fasting insulin]. Besides demographic variables, the above variables were selected as the risk factors based on the usage in clinical practice and the supporting evidence that demonstrated the association with NAFLD or its commonly accepted risk factors (*i.e.*, obesity, insulin resistance, and liver fibrosis).

Table 1 Baseline characteristics of participants with non-alcoholic fatty liver disease and controls *n* (%)

	NAFLD, <i>n</i> = 817	Controls, <i>n</i> = 2418
Demographic		
Age (yr)	48.47 ± 15.75	44.36 ± 16.20
Gender (male)	368 (45.04)	1004 (41.52)
Race/ethnicity		
White (non-Hispanic)	308 (37.70)	1046 (43.13)
Black (non-Hispanic)	193 (23.62)	705 (29.16)
Mexican American	280 (34.27)	550 (22.75)
Others	36 (4.41)	120 (4.96)
Body measurement		
Body mass index (kg/m ²)	30.38 ± 6.95	26.56 ± 5.42
Waist circumference (cm)	101.73 ± 16.32	90.84 ± 13.45
Biochemistry tests		
Iron (μg/dL)	75.35 ± 29.71	77.71 ± 32.75
TIBC (μg/dL)	364.79 ± 58.05	359.86 ± 56.59
Transferrin saturation (%)	21.13 ± 8.73	22.09 ± 9.65
Ferritin (ng/mL)	161.75 ± 152.55	110.16 ± 114.71
Cholesterol (mg/dL)	212.23 ± 44.95	202.73 ± 42.19
Triglyceride (mg/dL)	202.91 ± 137.97	136.58 ± 95.79
HDL cholesterol (mg/dL)	46.72 ± 16.80	52.10 ± 15.61
C-reactive protein (mg/dL) ¹	0.56 ± 0.80	0.45 ± 0.65
Uric acid (mg/dL)	5.62 ± 1.52	5.04 ± 1.42
Liver chemistry		
AST (U/L) ¹	24.76 ± 19.62	20.72 ± 14.71
ALT(U/L) ¹	22.78 ± 17.86	15.96 ± 12.14
GGT (U/L) ¹	42.87 ± 66.68	28.14 ± 41.69
ALP (U/L) ¹	93.72 ± 33.61	86.04 ± 36.29
Total bilirubin (mg/dL)	0.55 ± 0.30	0.54 ± 0.28
Total protein (g/dL)	7.49 ± 0.46	7.37 ± 0.45
Albumin (g/dL)	4.11 ± 0.35	4.12 ± 0.36
Diabetes testing profile		
HbA1c (%) ¹	6.02 ± 1.62	5.50 ± 1.09
FPG (mg/dL) ¹	114.50 ± 65.84	97.85 ± 35.68
Fasting C-peptide (pmol/mL)	1.11 ± 0.68	0.69 ± 0.53
Fasting insulin (μU/mL) ¹	21.85 ± 27.82	12.76 ± 19.24

¹The distribution is positively skewed with skewness > 3. ALP: Alkaline phosphatase; ALT: Alanine aminotransferase; AST: Aspartate aminotransferase; FPG: Fasting plasma glucose; GGT: Gamma glutamyl transferase; HbA1c: Glycated hemoglobin; TIBC: Total iron-binding capacity.

Definitions

In this study, NAFLD is defined as: (1) Diagnosed with moderate to severe hepatic steatosis on ultrasound; (2) no history of excessive alcohol intake in the past 12 mo; (3) not infected with hepatitis B or hepatitis C.

To evaluate the presence and extent of hepatic steatosis, readers used five main criteria: (1) Parenchymal brightness, (2) liver to kidney contrast, (3) deep beam attenuation, (4) bright vessel walls, and (5) gall-bladder wall definition. Based on the presence or absence of these five criteria, a main finding was categorized as normal, mild, moderate or severe^[20]. It is worth nothing that participants aged above 74 were not eligible for ultrasound study in NHANES III. For this reason, patients aged above 74 were excluded from this study.

Excessive alcohol intake is defined as more than 2 drinks per day for men or 1 drink per day for women in the past 12 mo, in which one drink of alcoholic beverage is equivalent to a 12 oz beer, a 5 oz glass of wine, or 1.5 oz of liquor. The average number of drinks per day is calculated from number of drinking days × number of drinks on drinking day/365 d. To qualify as hepatitis viral infection, participants must have tested positive for serum hepatitis B surface antigen or serum hepatitis C

antibody HCP (anti-HCV).

Statistical analysis

Statistical analyses were performed using STATA Release 14 (StataCorp LP, TX, United States). Numbers are presented in mean ± SD or number (%). All continuous factors were first tested for skewness; if the distributions were extremely skewed to the right (herein defined as skewness > 3), the factors were log transformed before using them as predictors in regression models. Since the response variable is dichotomous variable (NAFLD or non-NAFLD), logistic model is an appropriate model for determining if predictors are significantly associated with the response variable. As a result, logistic regression was used to determine if NAFLD is associated with any predictor in univariate level. Then, the significant factors from univariate analysis were included as predictors in step-wise logistic regression to determine the significant predictors in multivariate level. The significance level is 0.05.

RESULTS

Out of 18825 participants aged ≥ 20, there were 3235




Figure 1 Study design and study population.

participants ($n = 3235$) that passed the exclusion criteria as shown in Figure 1. Based on ultrasound findings, 817 (25.26%) participants were classified as NAFLD. Baseline characteristics of participants in study group and control group are summarized in Table 1.

For continuous variables, there were 8 factors having skewness greater than 3. Subsequently, the log transformation was applied to these factors, including C-reactive protein, AST, ALT, GGT, alkaline phosphatase, glycated hemoglobin, plasma glucose, and insulin. As shown in Table 2, there are 24 variables significantly associated with NAFLD in univariate level; the P -value of these significant factors mostly below 0.001.

As presented in Table 3, the number of significant factors reduced from 24 to 9 in multivariate analysis. The top three factors ranked by the highest change in pseudo R^2 (ΔR^2) are waist circumference (OR = 1.03, ΔR^2 = 2.13%, $P < 0.001$), C-peptide level (OR = 1.82, ΔR^2 = 1.33%, $P < 0.001$), and log_e of ALT (OR = 1.76, ΔR^2 = 1.16%, $P < 0.001$). The pseudo R^2 of the multivariate model is 16.68%.

DISCUSSION

The most significant NAFLD risk factor in both univariate and multivariate levels is waist circumference. Since waist circumference and BMI are highly interrelated surrogate markers of obesity^[21], it is not surprising to see one factor eliminated in multivariate level. Waist circumference—a measure of excess abdominal adiposity—has been identified as an independent risk factor for many obesity-related conditions, such as cardiovascular disease, type 2 diabetes, dyslipidemia, and hypertension, metabolic syndrome, polycystic ovary syndrome^[22–27]. The results from this study support that waist circumference is also an independent and probably the most important risk

factor for NAFLD in the United States population.

Insulin resistance is another well-known condition commonly found in NAFLD patients^[28,29]. Indeed, all diabetes test profiles were positively correlated with NAFLD by univariate analysis. In fact, type 2 diabetes can be diagnosed directly from HbA1c level ($\geq 6.5\%$) or FPG (≥ 126 mg/dL)^[30], while C-peptide and insulin are not routinely used in clinical practice to diagnose type 2 diabetes. While there are situations where C-peptide or insulin levels are useful—the diagnosis of insulinoma^[31], surreptitious use of insulin^[32], the diagnosis of type 2 diabetes in the young^[33], and the diagnosis of latent autoimmune diabetes in adults^[34], direct measurement of insulin levels and not of C-peptide has been used to assess insulin resistance. Previous studies often found that C-peptide levels are raised in patients with NASH^[35–37]. However, the application of C-peptide as a biomarker for interventions designed to improve insulin sensitivity remains to be determined. In case of NAFLD, there is only limited evidence; C-peptide was found to be associated with NAFLD in specific groups of population (i.e., obese adolescents and adults, latent autoimmune diabetes, and diabetes patients)^[13,14,38]. Our results are the first to show that C-peptide has a role not only as an independent risk factor for NAFLD but can also be useful for screening or monitoring the degree of insulin resistance in NAFLD in the general population. Based on ΔR^2 , we conclude that insulin resistance, as indicated by fasting C-peptide, is the second most important condition leading to NAFLD, second only to obesity as diagnosed by waist circumference, and is superior to measurement of fasting insulin levels.

Liver chemistries are used as an indicator of liver inflammation or liver cell damage. Commonly used liver chemistries include AST, ALT, ALP, GGT, total bilirubin, total protein, and albumin. For example, predominance of AST and ALT indicates hepatocellular injury; predominance of ALP and total bilirubin indicates cholestatic injury; an elevated ALP of hepatic origin may be confirmed GGT^[39–41]. As shown in Table 2, total protein and the natural log of AST, ALT, ALP, and GGT were positively associated with NAFLD in univariate analysis. However, only total protein and natural log of ALT were positively correlated with NAFLD in multivariate level. AST and ALT are the most widely used liver chemistries. The fact that ALT is included in multivariate model is not unexpected since ALT is generally higher than AST level in NAFLD^[40]. On the other hand, total protein is a non-specific marker of health, nutrition and liver synthetic capacity. Due to the fact that total protein consists of albumin and multiple subtypes of globulin, further investigation into the association between NAFLD and each subtype of globulin may provide a clearer explanation of our findings.

Ferritin is a protein that mainly stores iron in the body and serum ferritin level is the most accurate blood test to diagnose iron deficiency anemia^[42]. Recently, the role of ferritin as a biomarker in inflammatory diseases has been increasingly recognized^[43–45]. As an acute phase

Table 2 Univariate analysis for predictors of non-alcoholic fatty liver disease

	Beta	Standard error	Odds ratio	P value
Demographic				
Age (yr)	0.0157	0.0025	1.02	< 0.001 ^a
Gender (male)	0.1435	0.0815	1.15	0.078
Race/ethnicity				
White (non-Hispanic)	-0.226	0.0831	0.80	0.007 ^a
Black (non-Hispanic)	-0.2857	0.0937	0.75	0.002 ^a
Mexican American	0.5715	0.0882	1.77	< 0.001 ^a
Others	-0.1248	0.1945	0.88	0.521
Body measurement				
Body mass index (kg/m ²)	0.1004	0.0069	1.11	< 0.001 ^a
Waist circumference (cm)	0.0506	0.003	1.05	< 0.001 ^a
Biochemistry tests				
Iron (μg/dL)	-0.0024	0.0013	1.00	0.069
TIBC (μg/dL)	0.0015	0.0007	1.00	0.032 ^a
Transferrin saturation (%)	-0.0111	0.0044	0.99	0.012 ^a
Ferritin (ng/mL)	0.0029	0.0003	1.00	< 0.001 ^a
Cholesterol (mg/dL)	0.005	0.0009	1.01	< 0.001 ^a
Triglyceride (mg/dL)	0.0049	0.0004	1.00	< 0.001 ^a
HDL cholesterol (mg/dL)	-0.0261	0.0031	0.97	< 0.001 ^a
C-reactive protein (mg/dL) ¹	0.3398	0.0534	1.40	< 0.001 ^a
Uric acid (mg/dL)	0.2649	0.0277	1.30	< 0.001 ^a
Liver chemistry				
AST (U/L) ¹	1.004	0.1153	1.02	< 0.001 ^a
ALT (U/L) ¹	1.0274	0.0777	1.04	< 0.001 ^a
GGT (U/L) ¹	0.7441	0.0612	1.01	< 0.001 ^a
ALP (U/L) ¹	0.9011	0.1303	1.01	< 0.001 ^a
Total bilirubin (mg/dL)	0.0914	0.1395	1.09	0.512
Total protein (g/dL)	0.581	0.0896	1.79	< 0.001 ^a
Albumin (g/dL)	-0.0871	0.1141	0.92	0.445
Diabetes testing profile				
HbA1c (%) ¹	2.2201	0.2184	1.34	< 0.001 ^a
FPG (mg/dL) ¹	1.3125	0.1428	1.01	< 0.001 ^a
Fasting C-peptide (pmol/mL)	1.1976	0.0753	3.31	< 0.001 ^a
Fasting insulin (μU/mL) ¹	0.9646	0.059	1.02	< 0.001 ^a

¹Log transformation was applied to this factor before including in regression model. ^a*P* < 0.05. ALP: Alkaline phosphatase; ALT: Alanine aminotransferase; AST: Aspartate aminotransferase; FPG: Fasting plasma glucose; GGT: Gamma glutamyl transferase; HbA1c: Glycated hemoglobin; TIBC: Total iron-binding capacity. ^a*P* < 0.05.

Table 3 Risk factors of non-alcoholic fatty liver disease from step-wise logistic regression

	Beta	Standard error	Odds ratio	Change in pseudo R ²	P value
Demographic					
Race/ethnicity					
White (non-Hispanic)	0.2969	0.1162	1.35	0.18%	0.011
Mexican American	0.5495	0.1207	1.73	0.57%	0.020
Body measurement					
Waist circumference (cm)	0.0308	0.0035	1.03	2.13%	< 0.001
Biochemistry tests					
Ferritin (ng/mL)	0.0013	0.0004	1.00	0.15%	< 0.001
Triglyceride (mg/dL)	0.0008	0.0004	1.00	0.32%	< 0.001
Liver chemistry					
ALT (U/L) ¹	0.5658	0.0875	1.76	1.16%	< 0.001
Total protein (g/dL)	0.5319	0.1045	1.70	0.72%	< 0.001
Diabetes testing profile					
HbA1c (%) ¹	0.9266	0.2492	2.53	0.38%	< 0.001
Fasting C-peptide (pmol/mL)	0.6009	0.0877	1.82	1.33%	< 0.001

¹Log transformation was applied to this factor before including in regression model. Note: Pseudo R² = 16.68%; constant (Y-intercept) = 0.0000153; change in Pseudo R² is an incremental increase in Pseudo R² resulting from adding variable to model last. ALT: Alanine aminotransferase; HbA1c: Glycated hemoglobin.

response protein, ferritin concentrations increase during inflammation and may not reflect the size of total body

iron stores^[44]. Moreover, ferritin was found be associated with histologic severity and advanced fibrosis in patients

with NAFLD^[46-48].

Other factors significantly associated with NAFLD include race/ethnicity (non-Hispanic white and Mexican American), and triglyceride level. Race/ethnicity were often found associated with obesity-related diseases in United States based population^[49,50]. Triglyceride is an important biomarker of cardiovascular disease risk^[51], another condition highly interrelated with NAFLD.

There are several limitations in this study. First, the diagnosis of NAFLD in this study is based on the hepatic ultrasound results although liver biopsy remains the gold standard for the diagnosis of NAFLD. Second, the statistical analysis used is logistic regression. Since the relationship among these factors are complex, inter-related, and non-linear, linearity assumptions embedded in logistic regression may not be able to address all aspects of NAFLD. Furthermore, given a pseudo R^2 of 16.68%, only 16.68% of variation can be explained by multivariate model in Table 3.

In conclusion, NAFLD is associated with many conditions and factors. Three most important factors from multivariate model in this study are waist circumference, fasting C-peptide, and ALT. Further study is needed to validate the clinical utility of C-peptide in diagnosis or monitoring insulin resistance in NAFLD patients.

ARTICLE HIGHLIGHTS

Research background

Non-alcoholic fatty liver disease (NAFLD) is the most common cause of chronic liver disease in the United States. Additionally, NAFLD can progress to more severe liver diseases, such as non-alcoholic steatohepatitis, cirrhosis, and hepatocellular carcinoma. Many factors were found to be independently associated with NAFLD and/or severity of liver fibrosis in NAFLD. Nevertheless, there is limited evidence of the association between NAFLD and C-peptide.

Research motivation

Among many risk factors that are associated with NAFLD, obesity and insulin resistance are probably the most well-known ones. C-peptide levels can be used to measure insulin secretion and a surrogate marker of insulin resistance. However, C-peptide is not routinely used in clinical practice to diagnose type 2 diabetes or monitor insulin resistance status in NAFLD.

Research objectives

The objective of this study was to determine if fasting C-peptide is independently associated with NAFLD using multivariate analysis in the United States general population.

Research methods

Using the National Health and Nutrition Examination Survey 1988-1994, NAFLD participants aged 20 or greater without any other liver diseases were included in this study. The participants with excessive alcohol intake (> 2 drinks per day for males and > 1 drink per day for female) were excluded from the study. C-peptide and 27 other factors known to be associated with NAFLD (e.g., age, gender, body mass index, waist circumference, race/ethnicity, liver chemistries, and other diabetes tests) were selected as predictors in regression model. Univariate logistic regression and multivariate step-wise logistic regression were used to determine if the significant predictors of NAFLD, respectively.

Research results

There were 3235 participants ($n = 3235$) that passed the exclusion criteria. Based on ultrasound findings, 817 (25.26%) participants were classified as

NAFLD. Twenty-four variables were significantly associated with NAFLD in univariate level; the P -value of these significant factors mostly below 0.001. Using multivariate analysis, we found 9 out of 24 factors to be significantly associated with NAFLD. Ranked by ΔR^2 , the top three factors ranked are waist circumference (OR = 1.03, $\Delta R^2 = 2.13\%$, $P < 0.001$), C-peptide level (OR = 1.82, $\Delta R^2 = 1.33\%$, $P < 0.001$), and log_e of ALT (OR = 1.76, $\Delta R^2 = 1.16\%$, $P < 0.001$). The pseudo R^2 of the multivariate model is 16.68%.

Research conclusions

C-peptide is the second most important predictor of NAFLD in United States population after waist circumference.

Research perspectives

Further prospective research is needed to validate the clinical utility of fasting C-peptide in diagnosis or monitoring insulin resistance in NAFLD patients. Moreover, C-peptide should be considered as a potential factor for calculative liver scores to evaluate the fibrosis level.

REFERENCES

- 1 **Fraser A**, Longnecker MP, Lawlor DA. Prevalence of elevated alanine aminotransferase among US adolescents and associated factors: NHANES 1999-2004. *Gastroenterology* 2007; **133**: 1814-1820 [PMID: 18054554 DOI: 10.1053/j.gastro.2007.08.077]
- 2 **Vernon G**, Baranova A, Younossi ZM. Systematic review: the epidemiology and natural history of non-alcoholic fatty liver disease and non-alcoholic steatohepatitis in adults. *Aliment Pharmacol Ther* 2011; **34**: 274-285 [PMID: 21623852 DOI: 10.1111/j.1365-2036.2011.04724.x]
- 3 **Williams CD**, Stengel J, Asike MI, Torres DM, Shaw J, Contreras M, Landt CL, Harrison SA. Prevalence of nonalcoholic fatty liver disease and nonalcoholic steatohepatitis among a largely middle-aged population utilizing ultrasound and liver biopsy: a prospective study. *Gastroenterology* 2011; **140**: 124-131 [PMID: 20858492 DOI: 10.1053/j.gastro.2010.09.038]
- 4 **Beste LA**, Leipeztz SL, Green PK, Dominitz JA, Ross D, Ioannou GN. Trends in burden of cirrhosis and hepatocellular carcinoma by underlying liver disease in US veterans, 2001-2013. *Gastroenterology* 2015; **149**: 1471-1482.e5; quiz e17-8 [PMID: 26255044 DOI: 10.1053/j.gastro.2015.07.056]
- 5 **Rinella ME**. Nonalcoholic fatty liver disease: a systematic review. *JAMA* 2015; **313**: 2263-2273 [PMID: 26057287 DOI: 10.1001/jama.2015.5370]
- 6 **Buzzetti E**, Pinzani M, Tsochatzis EA. The multiple-hit pathogenesis of non-alcoholic fatty liver disease (NAFLD). *Metabolism* 2016; **65**: 1038-1048 [PMID: 26823198 DOI: 10.1016/j.metabol.2015.12.012]
- 7 **Da Silva HE**, Teterina A, Comelli EM, Taibi A, Arendt BM, Fischer SE, Lou W, Allard JP. Nonalcoholic fatty liver disease is associated with dysbiosis independent of body mass index and insulin resistance. *Sci Rep* 2018; **8**: 1466 [PMID: 29362454 DOI: 10.1038/s41598-018-19753-9]
- 8 **Ruhl CE**, Everhart JE. Fatty liver indices in the multiethnic United States National Health and Nutrition Examination Survey. *Aliment Pharmacol Ther* 2015; **41**: 65-76 [PMID: 25376360 DOI: 10.1111/apt.13012]
- 9 **Angulo P**, Hui JM, Marchesini G, Bugianesi E, George J, Farrell GC, Enders F, Saksena S, Burt AD, Bida JP, Lindor K, Sanderson SO, Lenzi M, Adams LA, Kench J, Thorneau TM, Day CP. The NAFLD fibrosis score: a noninvasive system that identifies liver fibrosis in patients with NAFLD. *Hepatology* 2007; **45**: 846-854 [PMID: 17393509 DOI: 10.1002/hep.21496]
- 10 **Harrison SA**, Oliver D, Arnold HL, Gogia S, Neuschwander-Tetri BA. Development and validation of a simple NAFLD clinical scoring system for identifying patients without advanced disease. *Gut* 2008; **57**: 1441-1447 [PMID: 18390575 DOI: 10.1136/gut.2007.146019]
- 11 **Ma H**, Xu C, Xu L, Yu C, Miao M, Li Y. Independent association of HbA1c and nonalcoholic fatty liver disease in an elderly Chinese

- population. *BMC Gastroenterol* 2013; **13**: 3 [PMID: 23294935 DOI: 10.1186/1471-230X-13-3]
- 12 **Van Cauter E**, Mestrez F, Sturis J, Polonsky KS. Estimation of insulin secretion rates from C-peptide levels. Comparison of individual and standard kinetic parameters for C-peptide clearance. *Diabetes* 1992; **41**: 368-377 [PMID: 1551497 DOI: 10.2337/diabetes.41.3.368]
- 13 **Chen M**, Li T, Zhang R, Li L, Lu J, Bao Y, Jia W. [Clinical features of non-alcoholic fatty liver disease and its relationship with serum C-peptide levels in patients with latent autoimmune diabetes in adults]. *Zhonghua Yi Xue Za Zhi* 2015; **95**: 3575-3578 [PMID: 26813367]
- 14 **Tricò D**, Caprio S, Umano GR, Pierpont B, Nouws J, Galderisi A, Kim G, Mata MM, Santoro N. Metabolic Features of Nonalcoholic Fatty Liver (NAFL) in Obese Adolescents: Findings from a Multi-ethnic Cohort. *Hepatology* 2018; [PMID: 29665034 DOI: 10.1002/hep.30035]
- 15 **Leighton E**, Sainsbury CA, Jones GC. A Practical Review of C-Peptide Testing in Diabetes. *Diabetes Ther* 2017; **8**: 475-487 [PMID: 28484968 DOI: 10.1007/s13300-017-0265-4]
- 16 **Wang L**, Lin P, Ma A, Zheng H, Wang K, Li W, Wang C, Zhao R, Liang K, Liu F, Hou X, Song J, Lu Y, Zhu P, Sun Y, Chen L. C-Peptide Is Independently Associated with an Increased Risk of Coronary Artery Disease in T2DM Subjects: A Cross-Sectional Study. *PLoS One* 2015; **10**: e0127112 [PMID: 26098780 DOI: 10.1371/journal.pone.0127112]
- 17 **Li Y**, Li Y, Meng L, Zheng L. Association between serum C-peptide as a risk factor for cardiovascular disease and high-density lipoprotein cholesterol levels in nondiabetic individuals. *PLoS One* 2015; **10**: e112281 [PMID: 25559358 DOI: 10.1371/journal.pone.0112281]
- 18 **Gonzalez-Mejia ME**, Porchia LM, Torres-Rasgado E, Ruiz-Vivanco G, Pulido-Pérez P, Báez-Duarte BG, Pérez-Fuentes R. C-Peptide Is a Sensitive Indicator for the Diagnosis of Metabolic Syndrome in Subjects from Central Mexico. *Metab Syndr Relat Disord* 2016; **14**: 210-216 [PMID: 26863426 DOI: 10.1089/met.2015.0067]
- 19 **National Center for Health Statistics**. Third National Health and Nutrition Examination Survey Data (NHANES III). Accessed May 29, 2018 Available from: URL: <https://www.cdc.gov/nchs/nhanes/nhanes3.htm>
- 20 **National Center for Health Statistics**. Third National Health and Nutrition Examination Survey: Hepatic Steatosis Ultrasound Images Assessment Procedures Manual. Accessed May 29, 2018 Available from: URL: https://www.cdc.gov/nchs/data/nhanes/nhanes3/hepatic_steatosis_ultrasound_procedures_manual.pdf
- 21 **Jensen MD**, Ryan DH, Apovian CM, Ard JD, Comuzzie AG, Donato KA, Hu FB, Hubbard VS, Jakicic JM, Kushner RF, Loria CM, Millen BE, Nonas CA, Pi-Sunyer FX, Stevens J, Stevens VJ, Wadden TA, Wolfe BM, Yanovski SZ; American College of Cardiology/American Heart Association Task Force on Practice Guidelines; Obesity Society. 2013 AHA/ACC/TOS guideline for the management of overweight and obesity in adults: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines and The Obesity Society. *J Am Coll Cardiol* 2014; **63**: 2985-3023 [PMID: 24239920 DOI: 10.1016/j.jacc.2013.11.004]
- 22 **Dimitriadis K**, Tsioufis C, Mazaraki A, Liataxis I, Koutra E, Kordalis A, Kasiakogias A, Flessas D, Tentolouris N, Tousoulis D. Waist circumference compared with other obesity parameters as determinants of coronary artery disease in essential hypertension: a 6-year follow-up study. *Hypertens Res* 2016; **39**: 475-479 [PMID: 26865004 DOI: 10.1038/hr.2016.8]
- 23 **Grundy SM**, Cleeman JI, Daniels SR, Donato KA, Eckel RH, Franklin BA, Gordon DJ, Krauss RM, Savage PJ, Smith SC Jr, Spertus JA, Costa F; American Heart Association; National Heart, Lung, and Blood Institute. Diagnosis and management of the metabolic syndrome: an American Heart Association/National Heart, Lung, and Blood Institute Scientific Statement. *Circulation* 2005; **112**: 2735-2752 [PMID: 16157765 DOI: 10.1161/CIRCULATIONAHA.105.169404]
- 24 **Janssen I**, Katzmarzyk PT, Ross R. Waist circumference and not body mass index explains obesity-related health risk. *Am J Clin Nutr* 2004; **79**: 379-384 [PMID: 14985210 DOI: 10.1093/ajcn/79.3.379]
- 25 **Lindström J**, Tuomilehto J. The diabetes risk score: a practical tool to predict type 2 diabetes risk. *Diabetes Care* 2003; **26**: 725-731 [PMID: 12610029 DOI: 10.2337/diacare.26.3.725]
- 26 **Pazderska A**, Kyaw Tun T, Phelan N, McGowan A, Sherlock M, Behan L, Boran G, Gibney J. In women with PCOS, waist circumference is a better surrogate of glucose and lipid metabolism than disease status per se. *Clin Endocrinol (Oxf)* 2018; **88**: 565-574 [PMID: 29285778 DOI: 10.1111/cen.13542]
- 27 **Pouliot MC**, Després JP, Lemieux S, Moorjani S, Bouchard C, Tremblay A, Nadeau A, Lupien PJ. Waist circumference and abdominal sagittal diameter: best simple anthropometric indexes of abdominal visceral adipose tissue accumulation and related cardiovascular risk in men and women. *Am J Cardiol* 1994; **73**: 460-468 [PMID: 8141087 DOI: 10.1016/0002-9149(94)90676-9]
- 28 **Gaggini M**, Morelli M, Buzzigoli E, DeFronzo RA, Bugianesi E, Gastaldello A. Non-alcoholic fatty liver disease (NAFLD) and its connection with insulin resistance, dyslipidemia, atherosclerosis and coronary heart disease. *Nutrients* 2013; **5**: 1544-1560 [PMID: 23666091 DOI: 10.3390/nu5051544]
- 29 **Asrih M**, Jornayvaz FR. Inflammation as a potential link between nonalcoholic fatty liver disease and insulin resistance. *J Endocrinol* 2013; **218**: R25-R36 [PMID: 23833274 DOI: 10.1530/JOE-13-0201]
- 30 **American Diabetes Association**. Diagnosis and classification of diabetes mellitus. *Diabetes Care* 2014; **37** Suppl 1: S81-S90 [PMID: 24357215 DOI: 10.2337/dc14-S081]
- 31 **Okabayashi T**, Shima Y, Sumiyoshi T, Kozuki A, Ito S, Ogawa Y, Kobayashi M, Hanazaki K. Diagnosis and management of insulinoma. *World J Gastroenterol* 2013; **19**: 829-837 [PMID: 23430217 DOI: 10.3748/wjg.v19.i6.829]
- 32 **Waickus CM**, de Bustros A, Shakil A. Recognizing factitious hypoglycemia in the family practice setting. *J Am Board Fam Pract* 1999; **12**: 133-136 [PMID: 10220236 DOI: 10.3122/jabfm.12.2.133]
- 33 **Hattersley A**, Bruining J, Shield J, Njolstad P, Donaghue K; International Society for Pediatric and Adolescent Diabetes. ISPAD Clinical Practice Consensus Guidelines 2006-2007. The diagnosis and management of monogenic diabetes in children. *Pediatr Diabetes* 2006; **7**: 352-360 [PMID: 17212604 DOI: 10.1111/j.1399-5448.2006.00217.x]
- 34 **Stenström G**, Gottsäter A, Bakhtadze E, Berger B, Sundkvist G. Latent autoimmune diabetes in adults: definition, prevalence, beta-cell function, and treatment. *Diabetes* 2005; **54** Suppl 2: S68-S72 [PMID: 16306343 DOI: 10.2337/diabetes.54.suppl_2.S68]
- 35 **Franque SM**, Verrijken A, Mertens I, Hubens G, Van Marck E, Pelckmans P, Michielsens P, Van Gaal L. Noninvasive assessment of nonalcoholic fatty liver disease in obese or overweight patients. *Clin Gastroenterol Hepatol* 2012; **10**: 1162-1168; quiz e87 [PMID: 22796457 DOI: 10.1016/j.cgh.2012.06.019]
- 36 **Hui JM**, Hodge A, Farrell GC, Kench JG, Kriketos A, George J. Beyond insulin resistance in NASH: TNF-alpha or adiponectin? *Hepatology* 2004; **40**: 46-54 [PMID: 15239085 DOI: 10.1002/hep.20280]
- 37 **Chitturi S**, Abeygunasekera S, Farrell GC, Holmes-Walker J, Hui JM, Fung C, Karim R, Lin R, Samarasinghe D, Liddle C, Weltman M, George J. NASH and insulin resistance: Insulin hypersecretion and specific association with the insulin resistance syndrome. *Hepatology* 2002; **35**: 373-379 [PMID: 11826411 DOI: 10.1053/jhep.2002.30692]
- 38 **Hua X**, Li M, Pan F, Xiao Y, Cui W, Hu Y. Non-alcoholic fatty liver disease is an influencing factor for the association of SHBG with metabolic syndrome in diabetes patients. *Sci Rep* 2017; **7**: 14532 [PMID: 29109457 DOI: 10.1038/s41598-017-15232-9]
- 39 **Kawachi I**, Robinson GM, Stace NH. A combination of raised serum AST:ALT ratio and erythrocyte mean cell volume level detects excessive alcohol consumption. *N Z Med J* 1990; **103**: 145-148 [PMID: 2342668]
- 40 **Kwo PY**, Cohen SM, Lim JK. ACG Clinical Guideline: Evaluation of Abnormal Liver Chemistries. *Am J Gastroenterol* 2017; **112**:

- 18-35 [PMID: 27995906 DOI: 10.1038/ajg.2016.517]
- 41 **Nyblom H**, Berggren U, Balldin J, Olsson R. High AST/ALT ratio may indicate advanced alcoholic liver disease rather than heavy drinking. *Alcohol Alcohol* 2004; **39**: 336-339 [PMID: 15208167 DOI: 10.1093/alcac/agh074]
- 42 **Short MW**, Domagalski JE. Iron deficiency anemia: evaluation and management. *Am Fam Physician* 2013; **87**: 98-104 [PMID: 23317073]
- 43 **Kell DB**, Pretorius E. Serum ferritin is an important inflammatory disease marker, as it is mainly a leakage product from damaged cells. *Metallomics* 2014; **6**: 748-773 [PMID: 24549403 DOI: 10.1039/c3mt00347g]
- 44 **Abitbol V**, Borderie D, Polin V, Maksimovic F, Sarfati G, Esch A, Tabouret T, Dhooge M, Dreanic J, Perkins G, Coriat R, Chaussade S. Diagnosis of Iron Deficiency in Inflammatory Bowel Disease by Transferrin Receptor-Ferritin Index. *Medicine (Baltimore)* 2015; **94**: e1011 [PMID: 26131803 DOI: 10.1097/MD.0000000000001011]
- 45 **Mecklenburg I**, Reznik D, Fasler-Kan E, Drewe J, Beglinger C, Hruz P; Swiss IBD Cohort Study Group. Serum hepcidin concentrations correlate with ferritin in patients with inflammatory bowel disease. *J Crohns Colitis* 2014; **8**: 1392-1397 [PMID: 24825446 DOI: 10.1016/j.crohns.2014.04.008]
- 46 **Sabrina N**, Bai CH, Chang CC, Chien YW, Chen JR, Chang JS. Serum Iron:Ferritin Ratio Predicts Healthy Body Composition and Reduced Risk of Severe Fatty Liver in Young Adult Women. *Nutrients* 2017; **9** [PMID: 28777296 DOI: 10.3390/nu9080833]
- 47 **Parikh P**, Patel J, Ingle M, Sawant P. Serum ferritin levels predict histological severity in patients with nonalcoholic fatty liver disease in India. *Indian J Gastroenterol* 2015; **34**: 200-208 [PMID: 26108652 DOI: 10.1007/s12664-015-0572-5]
- 48 **Kowdley KV**, Belt P, Wilson LA, Yeh MM, Neuschwander-Tetri BA, Chalasani N, Sanyal AJ, Nelson JE; NASH Clinical Research Network. Serum ferritin is an independent predictor of histologic severity and advanced fibrosis in patients with nonalcoholic fatty liver disease. *Hepatology* 2012; **55**: 77-85 [PMID: 21953442 DOI: 10.1002/hep.24706]
- 49 **Schmiegelow MD**, Hedlin H, Mackey RH, Martin LW, Vitolins MZ, Stefanick ML, Perez MV, Allison M, Hlatky MA. Race and ethnicity, obesity, metabolic health, and risk of cardiovascular disease in postmenopausal women. *J Am Heart Assoc* 2015; **4** [PMID: 25994446 DOI: 10.1161/JAHA.114.001695]
- 50 **Davidson MB**, Schriger DL. Effect of age and race/ethnicity on HbA1c levels in people without known diabetes mellitus: implications for the diagnosis of diabetes. *Diabetes Res Clin Pract* 2010; **87**: 415-421 [PMID: 20061043 DOI: 10.1016/j.diabres.2009.12.013]
- 51 **Nordestgaard BG**, Varbo A. Triglycerides and cardiovascular disease. *Lancet* 2014; **384**: 626-635 [PMID: 25131982 DOI: 10.1016/S0140-6736(14)61177-6]

P- Reviewer: Abenavoli L, Gheita TA, Hegardt FG, Tziomalos K
S- Editor: Wang XJ **L- Editor:** A **E- Editor:** Yin SY





Published by **Baishideng Publishing Group Inc**
7901 Stoneridge Drive, Suite 501, Pleasanton, CA 94588, USA
Telephone: +1-925-223-8242
Fax: +1-925-223-8243
E-mail: bpgoffice@wjgnet.com
Help Desk: <http://www.f6publishing.com/helpdesk>
<http://www.wjgnet.com>



ISSN 1007-9327

