

World Journal of *Gastroenterology*

World J Gastroenterol 2018 March 7; 24(9): 957-1062



REVIEW

- 957 Quantitative and noninvasive assessment of chronic liver diseases using two-dimensional shear wave elastography
Xie LT, Yan CH, Zhao QY, He MN, Jiang TA

ORIGINAL ARTICLE

Basic Study

- 971 Can bacterial virulence factors predict antibiotic resistant *Helicobacter pylori* infection?
Brennan DE, Dowd C, O'Morain C, McNamara D, Smith SM
- 982 Dysregulation of PARP1 is involved in development of Barrett's esophagus
Zhang C, Ma T, Luo T, Li A, Gao X, Wang ZG, Li F
- 992 Autophagy activation by Jiang Zhi Granule protects against metabolic stress-induced hepatocyte injury
Zheng YY, Wang M, Shu XB, Zheng PY, Ji G
- 1004 Long noncoding RNA RP4 functions as a competing endogenous RNA through miR-7-5p sponge activity in colorectal cancer
Liu ML, Zhang Q, Yuan X, Jin L, Wang LL, Fang TT, Wang WB

Retrospective Study

- 1013 Quality indicators in pediatric colonoscopy in a low-volume center: Implications for training
Lee WS, Tee CW, Koay ZL, Wong TS, Zahraq F, Foo HW, Ong SY, Wong SY, Ng RT
- 1022 Prognostic value of lymph nodes count on survival of patients with distal cholangiocarcinomas
Lin HP, Li SW, Liu Y, Zhou SJ

Clinical Practice Study

- 1035 Phenotypic and genotypic characterization of inflammatory bowel disease in children under six years of age in China
Fang YH, Luo YY, Yu JD, Lou JG, Chen J

Randomized Controlled Trial

- 1046 Effect of polyglycolic acid sheet plus esophageal stent placement in preventing esophageal stricture after endoscopic submucosal dissection in patients with early-stage esophageal cancer: A randomized, controlled trial
Chai NL, Feng J, Li LS, Liu SZ, Du C, Zhang Q, Linghu EQ

CASE REPORT

- 1056** Four cancer cases after esophageal atresia repair: Time to start screening the upper gastrointestinal tract
Vergouwe FW, Gottrand M, Wijnhoven BP, IJsselstijn H, Piessen G, Bruno MJ, Wijnen RM, Spaander MC

ABOUT COVER

Editorial board member of *World Journal of Gastroenterology*, Sang Chul Lee, MD, PhD, Chief Doctor, Full Professor, Professor, Surgeon, Surgical Oncologist, Division of Colorectal Surgery, Department of General Surgery, Catholic University of Korea, Daejeon St. Mary's Hospital, Daejeon 34943, South Korea

AIMS AND SCOPE

World Journal of Gastroenterology (*World J Gastroenterol*, *WJG*, print ISSN 1007-9327, online ISSN 2219-2840, DOI: 10.3748) is a peer-reviewed open access journal. *WJG* was established on October 1, 1995. It is published weekly on the 7th, 14th, 21st, and 28th each month. The *WJG* Editorial Board consists of 642 experts in gastroenterology and hepatology from 59 countries.

The primary task of *WJG* is to rapidly publish high-quality original articles, reviews, and commentaries in the fields of gastroenterology, hepatology, gastrointestinal endoscopy, gastrointestinal surgery, hepatobiliary surgery, gastrointestinal oncology, gastrointestinal radiation oncology, gastrointestinal imaging, gastrointestinal interventional therapy, gastrointestinal infectious diseases, gastrointestinal pharmacology, gastrointestinal pathophysiology, gastrointestinal pathology, evidence-based medicine in gastroenterology, pancreatology, gastrointestinal laboratory medicine, gastrointestinal molecular biology, gastrointestinal immunology, gastrointestinal microbiology, gastrointestinal genetics, gastrointestinal translational medicine, gastrointestinal diagnostics, and gastrointestinal therapeutics. *WJG* is dedicated to become an influential and prestigious journal in gastroenterology and hepatology, to promote the development of above disciplines, and to improve the diagnostic and therapeutic skill and expertise of clinicians.

INDEXING/ABSTRACTING

World Journal of Gastroenterology (*WJG*) is now indexed in Current Contents®/Clinical Medicine, Science Citation Index Expanded (also known as SciSearch®), Journal Citation Reports®, Index Medicus, MEDLINE, PubMed, PubMed Central and Directory of Open Access Journals. The 2017 edition of Journal Citation Reports® cites the 2016 impact factor for *WJG* as 3.365 (5-year impact factor: 3.176), ranking *WJG* as 29th among 79 journals in gastroenterology and hepatology (quartile in category Q2).

EDITORS FOR THIS ISSUE

Responsible Assistant Editor: *Xiang Li*
Responsible Electronic Editor: *Yan Huang*
Proofing Editor-in-Chief: *Lian-Sheng Ma*

Responsible Science Editor: *Ze-Mao Gong*
Proofing Editorial Office Director: *Jin-Lei Wang*

NAME OF JOURNAL
World Journal of Gastroenterology

ISSN
 ISSN 1007-9327 (print)
 ISSN 2219-2840 (online)

LAUNCH DATE
 October 1, 1995

FREQUENCY
 Weekly

EDITORS-IN-CHIEF
Damian Garcia-Olmo, MD, PhD, Doctor, Professor, Surgeon, Department of Surgery, Universidad Autonoma de Madrid; Department of General Surgery, Fundacion Jimenez Diaz University Hospital, Madrid 28040, Spain

Stephen C Strom, PhD, Professor, Department of Laboratory Medicine, Division of Pathology, Karolinska Institutet, Stockholm 141-86, Sweden

Andrzej S Tarnawski, MD, PhD, DSc (Med), Professor of Medicine, Chief Gastroenterology, VA Long Beach Health Care System, University of California, Irvine, CA, 5901 E. Seventh Str., Long Beach,

CA 90822, United States

EDITORIAL BOARD MEMBERS
 All editorial board members resources online at <http://www.wjgnet.com/1007-9327/editorialboard.htm>

EDITORIAL OFFICE
 Ze-Mao Gong, Director
World Journal of Gastroenterology
 Baishideng Publishing Group Inc
 7901 Stoneridge Drive, Suite 501,
 Pleasanton, CA 94588, USA
 Telephone: +1-925-2238242
 Fax: +1-925-2238243
 E-mail: editorialoffice@wjgnet.com
 Help Desk: <http://www.f0publishing.com/helpdesk>
<http://www.wjgnet.com>

PUBLISHER
 Baishideng Publishing Group Inc
 7901 Stoneridge Drive, Suite 501,
 Pleasanton, CA 94588, USA
 Telephone: +1-925-2238242
 Fax: +1-925-2238243
 E-mail: bpgoffice@wjgnet.com
 Help Desk: <http://www.f0publishing.com/helpdesk>
<http://www.wjgnet.com>

PUBLICATION DATE
 March 7, 2018

COPYRIGHT
 © 2018 Baishideng Publishing Group Inc. Articles published by this Open-Access journal are distributed under the terms of the Creative Commons Attribution Non-commercial License, which permits use, distribution, and reproduction in any medium, provided the original work is properly cited, the use is non commercial and is otherwise in compliance with the license.

SPECIAL STATEMENT
 All articles published in journals owned by the Baishideng Publishing Group (BPG) represent the views and opinions of their authors, and not the views, opinions or policies of the BPG, except where otherwise explicitly indicated.

INSTRUCTIONS TO AUTHORS
 Full instructions are available online at <http://www.wjgnet.com/bpg/gerinfo/204>

ONLINE SUBMISSION
<http://www.f0publishing.com>

Retrospective Study

Quality indicators in pediatric colonoscopy in a low-volume center: Implications for training

Way-Seah Lee, Chun-Wei Tee, Zhong-Lin Koay, Tat-Seng Wong, Fatimah Zahraq, Hee-Wei Foo, Sik-Yong Ong, Shin-Yee Wong, Ruey-Terng Ng

Way-Seah Lee, Chun-Wei Tee, Zhong-Lin Koay, Tat-Seng Wong, Fatimah Zahraq, Hee-Wei Foo, Sik-Yong Ong, Shin-Yee Wong, Ruey-Terng Ng, Department of Paediatrics, University Malaya Medical Centre, Kuala Lumpur 59100, Malaysia

Way-Seah Lee, Paediatric and Child Health Research Group, University Malaya, Kuala Lumpur 50603, Malaysia

ORCID number: Way-Seah Lee (0000-0001-9163-2828); Chun-Wei Tee (0000-0002-1119-2971); Zhong-Lin Koay (0000-0002-8724-9903); Tat-Seng Wong (0000-0002-3076-228X); Fatimah Zahraq (0000-0002-2658-1945); Hee-Wei Foo (0000-0001-7761-0550); Sik-Yong Ong (0000-0001-9647-5788); Shin-Yee Wong (0000-0002-7486-0594); Ruey-Terng Ng (0000-0003-1656-5797).

Author contributions: Lee WS conceived the idea of the research; Lee WS, Ng RT, Ong SY and Foo HW provided the clinical data; Tee CW, Koay ZL, Wong TS and Zahraq F collected the data; Wong SY performed the statistical analysis; Lee WS and Tee CW analyzed the data; Lee WS wrote the first draft; All authors contributed equally to writing of the final draft; All authors authorized the final version of the manuscript.

Supported by Research grant from Ministry of Higher Education, Malaysia, No. UM.C/625/HIR/MOHE/CHAN/13/1.

Institutional review board statement: This study was reviewed and approved by the Ethics Committee of the University Malaya Medical Centre (MEC reference: 902.15).

Informed consent statement: Patients were not required to give informed consent to the study because the analysis performed used anonymous clinical data that were obtained after each patient agreed to treatment by written consent.

Conflict-of-interest statement: All authors declare no conflicts-of-interest related to this article.

Data sharing statement: No additional data are available.

Open-Access: This article is an open-access article which was

selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

Manuscript source: Unsolicited manuscript

Correspondence to: Way-Seah Lee, MBBS, FRCPC, MD, Professor, Department of Paediatrics, Level 9, Women's and Children's Block, University Malaya Medical Center, Kuala Lumpur 59100, Malaysia. leews@ummc.edu.my
Telephone: +60-3-79492065
Fax: +60-3-79494704

Received: November 24, 2017

Peer-review started: November 25, 2017

First decision: December 27, 2017

Revised: January 15, 2018

Accepted: January 19, 2018

Article in press: January 19, 2018

Published online: March 7, 2018

Abstract

AIM

To study implications of measuring quality indicators on training and trainees' performance in pediatric colonoscopy in a low-volume training center.

METHODS

We reviewed retrospectively the performance of pediatric colonoscopies in a training center in Malaysia over 5 years (January 2010-December 2015), benchmarked against five quality indicators: appropriateness of indications, bowel preparations, cecum and ileal

examination rates, and complications. The European Society of Gastrointestinal Endoscopy guideline for pediatric endoscopy and North American Society for Pediatric Gastroenterology, Hepatology and Nutrition training guidelines were used as benchmarks.

RESULTS

Median (\pm SD) age of 121 children [males = 74 (61.2%)] who had 177 colonoscopies was 7.0 (\pm 4.6) years. On average, 30 colonoscopies were performed each year (range: 19-58). Except for investigations of abdominal pain (21/177, 17%), indications for colonoscopies were appropriate in the remaining 83%. Bowel preparation was good in 87%. One patient (0.6%) with severe Crohn's disease had bowel perforation. Cecum examination and ileal intubation rate was 95% and 68.1%. Ileal intubation rate was significantly higher in diagnosing or assessing inflammatory bowel disease (IBD) than non-IBD (72.9% *vs* 50.0% $P = 0.016$). Performance of four trainees was consistent throughout the study period. Average cecum and ileal examination rate among trainees were 97% and 77%.

CONCLUSION

Benchmarking against established guidelines helps units with a low-volume of colonoscopies to identify area for further improvement.

Key words: Pediatric colonoscopies; Quality indicators; Performance

© **The Author(s) 2018.** Published by Baishideng Publishing Group Inc. All rights reserved.

Core tip: Competency in colonoscopy is an essential component in the training for pediatric gastroenterology worldwide. We measured the performance of pediatric colonoscopy from a low-volume training center on quality indicators against established guidelines. The unit, which performed an average of 30 colonoscopies each year, performed well in clear indication for colonoscopy, good bowel preparation, safety and high rate of cecal examination (95%) but needs improvement for ileal intubation (at 68%). Benchmarking against established guidelines helps units with a low volume of colonoscopies to identify area for improvement.

Lee WS, Tee CW, Koay ZL, Wong TS, Zahraq F, Foo HW, Ong SY, Wong SY, Ng RT. Quality indicators in pediatric colonoscopy in a low-volume center: Implications for training. *World J Gastroenterol* 2018; 24(9): 1013-1021 Available from: URL: <http://www.wjgnet.com/1007-9327/full/v24/i9/1013.htm> DOI: <http://dx.doi.org/10.3748/wjg.v24.i9.1013>

INTRODUCTION

Colonoscopy is an essential diagnostic procedure for the evaluation and treatment of lower gastrointestinal

pathologies in children^[1-4]. Major indications for colonoscopy in children include rectal bleeding, investigation of diarrhea, failure to thrive and perianal lesions, and as initial diagnostic evaluation for inflammatory bowel disease (IBD)^[5-10].

Approximately 20%-30% of IBD patients have onset of disease in childhood^[11]. A study from Australia showed that IBD was the diagnosis in 58% of children who had initial diagnostic colonoscopy^[5]. The percentage is much lower in Asia, ranging from 10.9% in Taiwanese children^[8] to 19.6% in Korean children^[7]. The incidence of IBD is increasing worldwide^[12,13]. The incidence in Asia, including Malaysia, is increasing as well, although it is still relatively uncommon as compared to North America and Western Europe^[13,14].

Competency in colonoscopy has become an essential component in the training syllabus for both adult and pediatric gastroenterology worldwide^[15,16]. In adult colonoscopy, cecal intubation and detection for adenoma are considered as standard quality measures^[15]. In children, however, routine screening for adenomas is generally not recommended^[15]. On the other hand, ileal intubation is essential for accurate diagnosis of IBD, particularly Crohn's disease (CD)^[17]. Thus, appropriate indication for colonoscopy, complete examination including inspection of cecum and terminal ileum, adequate bowel preparation and free of complications are all important quality indicators in pediatric colonoscopy^[16].

In areas where the prevalence of IBD is low, such as Malaysia, the volume of pediatric colonoscopies performed may be limited^[7-10,18]. The reported cecum examination and ileal intubation rates vary. In Hong Kong, the cecal and ileal intubation rates were 97.6% and 75.6%^[10]. In Taiwan, the ileal intubation rate was 77.5%^[8]. In Australia, where the incidence of IBD is high, Singh *et al*^[5] reported that the cecal and ileal intubation rates were 96.3% and 92.4%, respectively.

A study on quality indicators published previously showed that performance benchmarked against quality indicators varies in different centers^[16]. To the best of our knowledge, however, no study on performance benchmarked against quality indicators from low-volume centers has been published previously.

We aimed to ascertain the performance of our unit when benchmarked against established quality indicators in pediatric colonoscopy covering the following areas: indications, quality of bowel preparation, safety and complications, cecal examination and terminal ileum intubation rates. We also assessed the implications of our performance to ascertain opportunities for improvements to the training program in this training center.

MATERIALS AND METHODS

This was a retrospective review on all pediatric colonoscopies performed between January 2010 and

December 2015 at the Paediatric Gastroenterology Unit, University Malaya Medical Centre (UMMC), Malaysia. The present study was approved by the institutional ethics review committee (MEC reference: 902.15). During the study period, the unit was staffed by one full-time consultant and a part-time visiting consultant. They were assisted by fellows-in-training, who spent the first 18 mo of a 3-year fellowship training program in the unit.

Data collection

Cases were retrieved from hospital and the unit database. The following data were collected: demographics and clinical features; indications for colonoscopy; laboratory data; degree of bowel preparation; extent of colonoscopic examination; and complications. Colonoscopic and pathological diagnoses were ascertained. Cases were excluded if the data were incomplete.

Quality indicators

The following areas were used as quality indicators: (1) appropriateness of indications; (2) quality of bowel preparation and extent of colonoscopic examination, including (3) cecum examination and (4) ileal intubation; and (5) safety (including anesthesia and sedation) and complications. Factors affecting extent of examination and the performances of trainees were also analyzed.

Indications and preparations for colonoscopy

In our unit, each referral for colonoscopy was screened and decided by a consultant. Generally, the indications for colonoscopy followed the established guidelines^[3,4], and has been reported previously^[18]. For the purpose of the present study, the European Society of Gastrointestinal Endoscopy (ESGE) guideline for pediatric endoscopy was used as a benchmark^[4].

Sedation and anesthesia

In our unit, colonoscopies were usually performed under general anesthesia. In adolescents, sedation (generally a combination of midazolam and pethidine) was used occasionally at the discretion of the anesthetist.

Bowel preparation

Bowel preparation has been standardized throughout the study period. Two days prior to colonoscopy, each patient was allowed a low residue diet. On the night before the procedure, each patient had bowel cleansing with polyethylene glycol solution and glycerin rectal enema. The degree of bowel preparation observed during colonoscopy was not standardized. It was judged by the endoscopist as poor, fair, good or excellent^[6].

Extent of colonoscopy

The extent of the colonoscopy was confirmed by visual identification of the colonic wall appearance, and the anatomy the cecum and terminal ileum. The biopsy of the terminal ileum was also used as an additional

confirmation. The recommendation by North American Society for Pediatric Gastroenterology, Hepatology and Nutrition (NASPGHAN) guidelines for training in pediatric gastroenterology was used as a benchmark^[19]. The NASPGHAN guidelines recommends a cecum and ileal examination rate of between 90%-95%^[19].

Performance

For the purpose of the present study, analysis on performance was confined to colonoscopies where an inspection of ileum was intended. This included cases where intubation of terminal ileum was indicated (*i.e.*, in diagnosing or assessing IBD), feasible (acceptable quality of bowel preparation where full examination was feasible) or safe (benefit of ileal intubation outweighs the risk of full examination, such as bowel perforation).

Performance by trainees

Analysis on the performance by trainees was confined to trainees who had completed a minimum of 12 mo training in the unit during the study period. Number of colonoscopies performed, cecum examination and ileal intubation rates were noted. In colonoscopies where trainees encountered technical difficulties during the procedure and were subsequently taken over by the consultant, the procedures were logged as performed by the consultant.

Statistical analysis

Data were collected and managed by using a statistical software program (SPSS version 20.0; SPSS Inc., IBM Corp., Armonk, NY, United States). Descriptive data were described in percentage, mean and median. Categorical data were analyzed using a two-tailed χ^2 test.

RESULTS

During the 6-year study period, 194 colonoscopies were performed in the unit. Data on 17 procedures were incomplete and were excluded from analysis. Of the remaining 177 colonoscopies, 56 were repeated procedures. Thus, 121 patients who had 177 colonoscopies were analyzed. The results are presented in two parts: (1) indications, colonoscopic findings and diagnosis of 121 patients who had first colonoscopy; and (2) quality indicators of 177 colonoscopies performed.

Volume of procedures

There was a steady increase in the number of colonoscopies performed each year during the study period (Figure 1). On average, 30 colonoscopies were performed each year during the study period, ranging from 19 procedures per year in the first 2 years to 58 procedures in 2015. Among the procedures, 15% (27/177) were logged as performed by consultants, while the remaining 85% (150/177) were performed

Number of colonoscopies performed during the study period




Figure 1 Number of colonoscopies performed each year during the study period.

by trainees supervised by a consultant.

Epidemiology, demographic features in 121 patients

There was a male preponderance (males = 74, 61%) in the 121 patients who had first colonoscopy (Table 1). The median (\pm SD) age was 7.5 (\pm 4.5) years. Eighty-one (67%) patients also had concomitant esophago-gastro-duodenoscopy (EGDS).

Indications for first colonoscopy in 121 patients

The most common indication was confirming the diagnosis of IBD (36/121, 30%; Table 1). Others were investigation of anemia or rectal bleeding (25/121, 21%). Investigation of abdominal pain was the indication in 17% (21/121). Most of the repeat colonoscopies were for disease assessment in IBD.

Colonoscopic findings in 121 patients

A positive finding was noted in 68 (56.2%) colonoscopies, while the remaining 53 (43.8%) had a normal colonoscopic finding. Indications for colonoscopy for the 53 patients with a negative finding were: excluding IBD ($n = 18$); disease assessment of pre-existing IBD ($n = 4$); assessment of abdominal pain ($n = 8$); ascertainment of lower gastrointestinal bleeding ($n = 10$); and, miscellaneous ($n = 13$).

Diagnosis in 121 patients

In addition to 16 patients who had colonoscopic assessment of preexisting IBD, a new clinical diagnosis or institution of new therapeutic measures was made in another 87 patients following colonoscopy. Thus, a total of 103 patients had a positive diagnosis. The diagnostic yield was 85%.

The colonoscopic diagnoses are shown in Table 1. Overall, 50 (41%) patients had a diagnosis of IBD (newly diagnosed, $n = 34$; diagnosis confirmed elsewhere, $n = 16$; Figure 2). Of these, 30 patients had CD and 20 had ulcerative colitis, respectively.

Another 22 (18%) patients had focal inflammation of the rectum (IBD-unclassified) or solitary rectal ulcer syndrome.

Trend for diagnosis of IBD in 121 patients

The number of new cases of IBD seen in the unit during the study period is shown in Figure 2. There was an increasing trend in the number of new IBD cases, especially for CD.

Sedation and anesthesia in 177 colonoscopies

Vast majority (165/177, 93.2%) of the procedures were performed under general anesthesia. The remaining 12 procedures (6.8%) were performed under sedation, being administered by anesthetist. No major events related to anesthesia or sedation were observed during the study period.

Bowel preparation in 177 colonoscopies

Bowel preparation was judged to be good by the endoscopist in 87% (155/177) of the patients, moderate in 0.6% (1/177) and bad in 12% (21/177) patients.

Cecum and terminal ileum intubation in 177 colonoscopies

Information on the extent of colonoscopy was available in all 177 procedures (Table 1). The overall ileal intubation rate was 54.2% (96/177). Cecum was examined in an additional 22.0% (39/177). Thus, the cecum was reached in 76.3% (135/177) of patients. The extents of colonoscopy of the remaining 42 procedures are shown in Table 1.

In 36 patients, full colonoscopic examination and ileal intubation were not intended. They were not indicated in 18 cases, including those for confirmation or surveillance of graft-versus-host disease ($n = 10$), confirmation of rectal metastasis ($n = 2$), tissue biopsy for malabsorption (1 each for food protein-induced

Table 1 Indications, diagnoses and quality indicators in 121 children who had 177 colonoscopies

	<i>n</i>	%
Sex, males	74	61
Age in yr, median ± SD	7.5 ± 4.5	
Concomitant esophagogastroduodenoscopy	81	67
Indications, <i>n</i> = 121 ¹		
Suspected of inflammatory bowel disease, new patients	36	30
Per rectal bleeding/investigations of anemia	25	21
Investigation of gastrointestinal symptoms	21	17
Assessment of inflammatory bowel disease, diagnosed elsewhere	16	13
Suspected of colonic polyps	7	6
Exclusion of graft-versus-host disease or colonic malignancies	7	6
Assessment of failure to thrive/malabsorption	4	3.3
Others	10	8
Colonoscopic diagnosis, <i>n</i> = 121		
Crohn's disease	30	25
Ulcerative colitis	20	17
Non-specific colitis or solitary rectal ulcer syndrome	22	18
Infective colitis	9	7
Colonic polyps	8	7
Graft-versus-host disease	5	4
Malabsorption	2	2
Allergic colitis	2	2
Miscellaneous diagnosis	5	4
No diagnosis	18	15
Extent of colonoscopic examination, 177 colonoscopies		
Terminal ileum	96	54
Cecum	38	21
Ascending colon	9	5
Transverse colon	13	7
Descending colon	12	7
Sigmoid colon	7	4
Rectum	1	0.6
Reached cecum but no terminal ileum intubation	134	76
Ileal intubation not intended, 177 colonoscopies ²	36	
Not indicated	18	
Distorted anatomy due to previous surgery	1	
External stricture	2	
Previous colostomy in Crohn's disease	4	
Large polyp at rectum	2	
Risk of perforation outweighs benefit of full examination due to severe colitis	5	
Poor bowel preparation	4	
Full colonoscopic examination intended, 177 colonoscopies ²	141	
Ileal intubation	96	68
Cecum examination	134	95

¹Some patients had more than one indication; ²Not intended include not indicated, not feasible or the risks of full examination outweigh the benefit of full examination.

enterocolitis syndrome and autoimmune enteropathy), trichuriasis (*n* = 2), and assessment of previously confirmed solitary rectal ulcer syndrome (*n* = 2). Risk of perforation was judged to outweigh benefit of full examination in 5 patients. All had IBD with severe colitis and friable mucosal wall (ulcerative colitis, *n* = 3; CD, *n* = 2). Poor bowel preparation prevented a complete examination in 4 patients. A large rectal polyp obstructing the lumen (*n* = 1) and excessive bleeding in a patient with a large rectal polyp (*n* = 1) prevented a complete colonoscopy in 2 patients.

Of the remaining 141 patients in whom inspection of the terminal ileum was intended, the cecum examination and ileal intubation rates were 95.0%

(134/141) and 68.1% (96/141), respectively. Overall, 45.8% of colonoscopies did not include an inspection of terminal ileum, and 31.9% did not reach terminal ileum when it was intended. Similarly, 24.3% of the procedures did not reach the cecum, and 5.0% failed to reach cecum when it was intended.

Factors affecting complete examination

Rate of complete examination was not significantly affected by the age of patients [ileal intubation rate; < 5 years (35/141) vs ≥ 5 years (106/141) = 62.9% vs 69.8% *P* = 0.44]. However, the ileal intubation rate was significantly higher when the indication for the colonoscopy was for the diagnosis or assessment of

Table 2 Colonoscopic performance by trainees

	Trainee A	Trainee B	Trainee C	Trainee D	Average
Duration of training in mo	18	18	24	18	19.5
Number of colonoscopies performed during training period	19	17	44	36	29
Number of colonoscopies where intubation of terminal ileum was intended	16	13	39	28	24
Intubation of terminal ileum	12 (75)	10 (77)	34 (87)	18 (64)	77%
Examination of cecum and terminal ileum	16 (100)	13 (100)	38 (97)	26 (93)	97%

Data are presented as *n* or *n* (%), unless otherwise indicated. Note: In colonoscopies where trainees encountered technical difficulties during the procedure and were subsequently taken over by the consultant, the procedures were logged as performed by the consultant.

Number of new cases of IBD seen in the unit; 2010-2015




Figure 2 Number of new cases of pediatric inflammatory bowel disease seen in the unit during the study period. CD: Crohn's disease; IBD: Inflammatory bowel disease; UC: Ulcerative colitis.

IBD (73.0% vs 50%; $P = 0.016$).

Performance by trainees

During the study period, four trainees completed a minimum of 12-mo training in pediatric colonoscopy in the unit (Table 2). Part of the training period of another three trainees fell outside the study period and were not included in the analysis. The average number of colonoscopies performed was 29. The overall cecum examination rate was 97%, while the overall ileal intubation rate was 77%. There was a consistent performance by the trainees (Table 2).

Complications in 177 colonoscopies

A 7-year-old boy with CD who had gross delay in referral, severe malnutrition and severe mucosal ulcerations had iatrogenic perforation of the colon during the procedure. The patient had an uneventful recovery after colostomy and repair. Another patient with rectal polyp developed bleeding while it was removed during colonoscopy. The bleeding stopped after cauterization. No blood transfusion was required. No other complications were noted in the remaining 175 procedures, either associated with the procedure,

anesthesia or sedation.

DISCUSSION

To the best of our knowledge, the present study is the first of its kind on performance benchmarked against established quality indicators in pediatric colonoscopy from a center where the volume of colonoscopy is low. During the study period, the number of colonoscopies performed in the center each year ranged from 19 to 58, with an average of 30. It was conducted in a setting with a low but rapidly increasing incidence of IBD.

Our results reveal that of the five quality indicators which were benchmarked against, the performance was high in four indicators. Most of the indications for colonoscopy performed in the unit were appropriate, following the recommendations of ESGE guidelines. Performances were good in bowel preparation, patient safety and complications. The cecum examination rate reached the expected rate at 95% by NASPGHAN, but was somewhat limited in ileal intubation rate at 68.1%. The performance of all trainees was consistently good in cecum examination but variable in ileal intubation.

There are several important implications on the results of the present study. Firstly, the indications for colonoscopies in the present study were mostly appropriate according to established guidelines^[4]. The overall diagnostic yield was 85%. Previously, we observed that 99.7% of all EGDS and colonoscopies performed in the unit were appropriate^[18]. The most common indications for colonoscopy were diagnosing or assessing IBD and ascertaining the cause of anemia. There was an increasing trend in the number of newly-diagnosed IBD cases during the study period. Investigation of abdominal symptoms, mainly abdominal pain, was the indication in only 17% of the patients^[4].

In adult colonoscopy, screening for colon cancer is the main quality indicator^[19]. Surrogate measures, such as colonic adenoma detection and complete examination of the colon, have been found to correlate with high-quality colonoscopy^[20]. Pediatric colonoscopy, on the other hand, is fundamentally a different procedure with different indications, technical requirement, and different quality indicators^[21]. The most important difference between adult and pediatric colonoscopies are the size of the patients and indications^[21].

Screening for IBD is an important indication in pediatric colonoscopy. Differentiating between ulcerative colitis and CD is important in the diagnosis of IBD. Thus, examination of the cecum and terminal ileum are important quality indicators in pediatric colonoscopy^[22]. In particular, intubation of the terminal ileum and sampling biopsy is essential for confirming the diagnosis of CD^[23,24]. Pediatric endoscopy training guidelines suggest the cecal intubation rate to be at least 90% to 95%, with a comparable terminal ileum intubation rate^[16,19].

Based on these quality indicators, the performances of centers vary. In Australia where screening for IBD was the major indication for colonoscopy, the reported cecum and ileal examination rates were 96.3% and 92.4%, respectively^[5]. In Taiwan, the reported ileal intubation rate was 77.5%, while in China it was 81.7%^[8,9]. The ileal intubation rates in the multicenter consortium review from the United States ranged from 30% to 90%^[16].

After excluding procedures where either complete colonoscopy was not indicated, unsafe or not feasible, the cecum examination and ileal intubation rates in the present study were 95% and 68.1%, respectively. The cecum examination was comparable to the NASPGHAN pediatric gastroenterology training guidelines^[19], but the ileal intubation rate was somewhat lower. Nevertheless, these figures were comparable with those reported in the region and the United States multicenter consortium study^[8,16].

Bowel preparation was noted to be good in the majority of the cases (87%). Colonic perforation was observed in 1 patient who had severe long-standing CD prior to referral. The perforation rate was 0.5%.

We reported two perforations in 66 colonoscopies previously^[18].

The present study showed that all the trainees have a satisfactory cecum intubation rate, although the ileal intubation rate needs to be improved. To achieve cecal intubation examination and ileal intubation rate of 90%-95%, NASPGHAN guidelines for training in pediatric gastroenterology recommended 120 colonoscopies for pediatric trainees^[19]. In this center, each trainee performed about 12-20 colonoscopies in children each year during their training. Before starting pediatric colonoscopy under supervision, the trainees were required to undergo training in adult colonoscopy, performing at least 50 colonoscopies in adult patients.

The implication of this is that the trainees would not have adequate training opportunity of reaching the recommended 120 colonoscopies in pediatric colonoscopy if the entire training was spent in this unit alone. Trainees spent between 18 mo to 24 mo of training in this center. The trainees usually start the training in endoscopic procedures with the adult gastroenterologists before being allowed to perform pediatric colonoscopy. The final year of the training is usually spent in a center of excellence for training in Australia or United Kingdom, where there is abundant opportunity for pediatric colonoscopy. Thus, even though the ileal intubation rate of the trainees noted in the present study did not reach the intended goal suggested by the NASPGHAN training guidelines^[19], it is expected that their performance will be enhanced further once the overseas training is completed.

Another potential way to enhance colonoscopic skills is simulated colonoscopy training^[25]. Studies involving trainees in adult colonoscopy showed improved performances during patient-based colonoscopy^[25]. However, to date, no such study has been published involving training in pediatric colonoscopy.

Other important quality indicators which were not included in the present study were documentation of American Society of Anesthesiologist Physical Status Classification System (ASA) risk assessment and procedure duration. ASA risk assessment was routinely checked by the anesthetist. It was documented separately but was not captured in the present study. Additional potential measures of high-quality pediatric colonoscopy which have been mentioned but not included in this study are minimization of air insufflation, ease of patient sedation, duration of procedure, performance of mucosal biopsy sampling, patient recovery time and ongoing procedural competency assessment^[22].

An obvious limitation to this study is its retrospective nature. The data was only collected from one center. However, our center is the only training center for pediatric gastroenterology in the whole country; we are unable to benchmark our performance against other centers. Nevertheless, with the exception of a somewhat lower ileal intubation rate, the performance in other quality indicators is excellent.

In conclusion, there is an increasing trend in the number of colonoscopies performed each year in this center. This is due to the increasing number of newly-diagnosed IBD cases during the study period. The performance of pediatric colonoscopy in this center was excellent in four of the five quality indicators benchmarked. The ileal intubation rate needs to be improved. A period of training in a center with a large volume of IBD will enhance the skills and performance of colonoscopy among the trainees of pediatric gastroenterology in Malaysia.

ARTICLE HIGHLIGHTS

Research background

The incidence of inflammatory bowel disease is on the rise worldwide, including in Asia. The gold standard for the diagnosis of inflammatory bowel disease is histologic confirmation from tissue biopsies obtained during esophago-gastro-duodenoscopy and colonoscopy. In particular, differentiating Crohn's disease from ulcerative colitis is dependent upon inspection and biopsy of the terminal ileum. Thus, intubation of the terminal ileum is considered as an important quality indicator in pediatric colonoscopy. Current guidelines by the North American Society of Pediatric Gastroenterology, Hepatology and Nutrition recommend the cecum examination and terminal ileum of 95%. This target should be used by pediatric gastroenterology centers worldwide to benchmark their performance.

Research motivation

Current literature showed that reported ileal intubation rate in pediatric colonoscopy from centers around Asia ranged from 75.6% to 77.5%. The performance of our unit, from an area where the incidence of inflammatory bowel disease is currently low but is on the rise, is unknown. Thus, we were motivated to benchmark our performance against current recommendation to identify areas for improvement to enhance the quality of our training program.

Research objectives

The main objective was to benchmark the performance of our unit, in particular the completeness of colonoscopic examination, *i.e.* cecal examination and ileal intubation, against current recommended guidelines. We also evaluated other indicators such as appropriateness of indications, level of bowel preparation, as well as safety and complications of the procedures encountered.

Research methods

We conducted a retrospective analysis on all the pediatric colonoscopies performed in a pediatric gastroenterology training center in Malaysia over a period of 6 years. We included the following indicators: appropriateness of indications; quality of bowel preparation; safety and complications; as well as cecal examination and terminal ileum intubation rates. The performances of trainees in the cecal and ileal examination rates were ascertained separately.

Research results

We found that of the 177 colonoscopies performed, the diagnostic yield was 85%, quality of bowel preparation was good in 87%, while one of 177 procedures was complicated by perforation during the procedure. The overall cecum examination rate was 76.3% and ileal intubation rate was 54.2%. After excluding colonoscopy where full colonoscopic examination was either not indicated, not feasible because of poor bowel preparation or unsafe (severe colitis), the cecum examination rate was 95.0% and ileal intubation rate was 68.1%. Among four trainees who completed a minimum of 12 mo training, the overall cecum rate was 97% while the overall ileal intubation rate was 77%. The performance of all trainees was consistent. Thus, the cecum examination rate of our unit was satisfactory but the rate of terminal ileum intubation needs further improvement. To improve the rate of ileal intubation, the trainees would spend part of their training program in a center of excellence with adequate

volume of pediatric colonoscopy.

Research conclusions

The present study was the first attempt by a pediatric gastroenterology unit in Asia to benchmark its performance in pediatric colonoscopy against established international guidelines. Our study suggests that such a benchmark is both applicable and desirable. The study allows our unit to identify areas for further improvement. Trainees from our unit now routinely spend part of their training in a center of excellence to enhance their skills in colonoscopy.

Research perspectives

Benchmarking against established guidelines has been adopted as part of quality assurance of our unit. We plan to conduct a prospective study to include other indicators of good practice not included in this retrospective review. These include anesthetic risk assessment, duration of procedure, ease of sedation, quality of mucosal biopsy sampling and patient recovery time.

REFERENCES

- 1 **Teague RH**, Salmon PR, Read AE. Fiberoptic examination of the colon: a review of 255 cases. *Gut* 1973; **14**: 139-142 [PMID: 4540492]
- 2 **Hassall E**, Barclay GN, Ament ME. Colonoscopy in childhood. *Pediatrics* 1984; **73**: 594-599 [PMID: 6718114]
- 3 **ASGE Standards of Practice Committee**, Lightdale JR, Acosta R, Shergill AK, Chandrasekhara V, Chathadi K, Early D, Evans JA, Fanelli RD, Fisher DA, Fonkalsrud L, Hwang JH, Kashab M, Muthusamy VR, Pasha S, Saltzman JR, Cash BD; American Society for Gastrointestinal Endoscopy. Modifications in endoscopic practice for pediatric patients. *Gastrointest Endosc* 2014; **79**: 699-710 [PMID: 24593951 DOI: 10.1016/j.gie.2013.08.014]
- 4 **Thomson M**, Tringali A, Dumonceau JM, Tavares M, Tabbers MM, Furlano R, Spaander M, Hassan C, Tzvinikos C, Ijsselstijn H, Viala J, Dall'Oglio L, Benninga M, Orel R, Vandenplas Y, Keil R, Romano C, Brownstone E, Hlava Š, Gerner P, Dolak W, Landi R, Huber WD, Everett S, Vecsei A, Aabakken L, Amil-Dias J, Zambelli A. Paediatric Gastrointestinal Endoscopy: European Society for Paediatric Gastroenterology Hepatology and Nutrition and European Society of Gastrointestinal Endoscopy Guidelines. *J Pediatr Gastroenterol Nutr* 2017; **64**: 133-153 [PMID: 27622898 DOI: 10.1097/MPG.0000000000001408]
- 5 **Singh HK**, Withers GD, Ee LC. Quality indicators in pediatric colonoscopy: an Australian tertiary center experience. *Scand J Gastroenterol* 2017; **52**: 1453-1456 [PMID: 28936881 DOI: 10.1080/00365521.2017.1380224]
- 6 **Wahid AM**, Devarajan K, Ross A, Zilbauer M, Heuschkel R. Paediatric gastrointestinal endoscopy: a qualitative study. *Eur J Gastroenterol Hepatol* 2016; **28**: 25-29 [PMID: 26473298 DOI: 10.1097/MEG.0000000000000488]
- 7 **Lee YW**, Chung WC, Sung HJ, Kang YG, Hong SL, Cho KW, Kang D, Lee IH, Jeon EJ. Current status and clinical impact of pediatric endoscopy in Korea. *Korean J Gastroenterol* 2014; **64**: 333-339 [PMID: 25530584]
- 8 **Wu CT**, Chen CA, Yang YJ. Characteristics and Diagnostic Yield of Pediatric Colonoscopy in Taiwan. *Pediatr Neonatol* 2015; **56**: 334-338 [PMID: 25850637 DOI: 10.1016/j.pedneo.2015.01.005]
- 9 **Lei P**, Gu F, Hong L, Sun Y, Li M, Wang H, Zhong B, Chen M, Cui Y, Zhang S. Pediatric colonoscopy in South China: a 12-year experience in a tertiary center. *PLoS One* 2014; **9**: e95933 [PMID: 24759776 DOI: 10.1371/journal.pone.0095933]
- 10 **Tam YH**, Lee KH, Chan KW, Sihoe JD, Cheung ST, Mou JW. Colonoscopy in Hong Kong Chinese children. *World J Gastroenterol* 2010; **16**: 1119-1122 [PMID: 20205284 DOI: 10.3748/wjg.v16.i9.1119]
- 11 **Ruel J**, Ruane D, Mehandru S, Gower-Rousseau C, Colombel JF. IBD across the age spectrum: is it the same disease? *Nat Rev*

- Gastroenterol Hepatol* 2014; **11**: 88-98 [PMID: 24345891 DOI: 10.1038/nrgastro.2013.240]
- 12 **Ng SC**, Tang W, Ching JY, Wong M, Chow CM, Hui AJ, Wong TC, Leung VK, Tsang SW, Yu HH, Li MF, Ng KK, Kamm MA, Studd C, Bell S, Leong R, de Silva HJ, Kasturiratne A, Mufeeena MN, Ling KL, Ooi CJ, Tan PS, Ong D, Goh KL, Hilmi I, Pisespongsa P, Manatsathit S, Rerknimitr R, Aniwan S, Wang YF, Ouyang Q, Zeng Z, Zhu Z, Chen MH, Hu PJ, Wu K, Wang X, Simadibrata M, Abdullah M, Wu JC, Sung JJ, Chan FK; Asia-Pacific Crohn's and Colitis Epidemiologic Study (ACCESS) Study Group. Incidence and phenotype of inflammatory bowel disease based on results from the Asia-pacific Crohn's and colitis epidemiology study. *Gastroenterology* 2013; **145**: 158-165.e2 [PMID: 23583432 DOI: 10.1053/j.gastro.2013.04.007]
 - 13 **Malmborg P**, Hildebrand H. The emerging global epidemic of paediatric inflammatory bowel disease--causes and consequences. *J Intern Med* 2016; **279**: 241-258 [PMID: 26355194 DOI: 10.1111/joim.12413]
 - 14 **Hilmi I**, Jaya F, Chua A, Heng WC, Singh H, Goh KL. A first study on the incidence and prevalence of IBD in Malaysia--results from the Kinta Valley IBD Epidemiology Study. *J Crohns Colitis* 2015; **9**: 404-409 [PMID: 25744112 DOI: 10.1093/ecco-jcc/jjv039]
 - 15 **Walsh CM**, Ling SC, Mamula P, Lightdale JR, Walters TD, Yu JJ, Carnahan H. The gastrointestinal endoscopy competency assessment tool for pediatric colonoscopy. *J Pediatr Gastroenterol Nutr* 2015; **60**: 474-480 [PMID: 25564819 DOI: 10.1097/MPG.0000000000000686]
 - 16 **Thakkar K**, Holub JL, Gilger MA, Shub MD, McOmber M, Tsou M, Fishman DS. Quality indicators for pediatric colonoscopy: results from a multicenter consortium. *Gastrointest Endosc* 2016; **83**: 533-541 [PMID: 26253014 DOI: 10.1016/j.gie.2015.06.028]
 - 17 **Levine A**, Koletzko S, Turner D, Escher JC, Cucchiara S, de Ridder L, Kolho KL, Veres G, Russell RK, Paerregaard A, Buderus S, Greer ML, Dias JA, Veereman-Wauters G, Lionetti P, Sladek M, Martin de Carpi J, Staiano A, Ruemmele FM, Wilson DC; European Society of Pediatric Gastroenterology, Hepatology, and Nutrition. ESPGHAN revised porto criteria for the diagnosis of inflammatory bowel disease in children and adolescents. *J Pediatr Gastroenterol Nutr* 2014; **58**: 795-806 [PMID: 24231644 DOI: 10.1097/MPG.0000000000000239]
 - 18 **Lee WS**, Zainuddin H, Boey CC, Chai PF. Appropriateness, endoscopic findings and contributive yield of pediatric gastrointestinal endoscopy. *World J Gastroenterol* 2013; **19**: 9077-9083 [PMID: 24379634 DOI: 10.3748/wjg.v19.i47.9077]
 - 19 **Leichtner AM**, Gillis LA, Gupta S, Heubi J, Kay M, Narkewicz MR, Rider EA, Rufo PA, Sferra TJ, Teitelbaum J; NASPGHAN Training Committee; North American Society for Pediatric Gastroenterology. NASPGHAN guidelines for training in pediatric gastroenterology. *J Pediatr Gastroenterol Nutr* 2013; **56** Suppl 1: S1-S8 [PMID: 23263531 DOI: 10.1097/MPG.0b013e31827a78d6]
 - 20 **Reh DK**. Quality in colonoscopy: cecal intubation first, then what? *Am J Gastroenterol* 2006; **101**: 732-734 [PMID: 16635220 DOI: 10.1111/j.1572-0241.2006.00483.x]
 - 21 **Reh DK**. Colonoscopy. *Gastrointest Endosc* 2013; **78**: 444-449 [PMID: 23948194 DOI: 10.1016/j.gie.2013.06.025]
 - 22 **Kay M**, Wylie R. It's all about the loop: quality indicators in pediatric colonoscopy. *Gastrointest Endosc* 2016; **83**: 542-544 [PMID: 26897046 DOI: 10.1016/j.gie.2015.08.058]
 - 23 **Sawczenko A**, Sandhu BK. Presenting features of inflammatory bowel disease in Great Britain and Ireland. *Arch Dis Child* 2003; **88**: 995-1000 [PMID: 14612366]
 - 24 **Batres LA**, Maller ES, Ruchelli E, Mahboubi S, Baldassano RN. Terminal ileum intubation in pediatric colonoscopy and diagnostic value of conventional small bowel contrast radiography in pediatric inflammatory bowel disease. *J Pediatr Gastroenterol Nutr* 2002; **35**: 320-323 [PMID: 12352520]
 - 25 **Grover SC**, Scaffidi MA, Khan R, Garg A, Al-Mazroui A, Alomani T, Yu JJ, Plener IS, Al-Awamy M, Yong EL, Cino M, Ravindran NC, Zasowski M, Grantcharov TP, Walsh CM. Progressive learning in endoscopy simulation training improves clinical performance: a blinded randomized trial. *Gastrointest Endosc* 2017; **86**: 881-889 [PMID: 28366440 DOI: 10.1016/j.gie.2017.03.1529]

P- Reviewer: Choi YS **S- Editor:** Gong ZM **L- Editor:** Filipodia
E- Editor: Huang Y





Published by **Baishideng Publishing Group Inc**
7901 Stoneridge Drive, Suite 501, Pleasanton, CA 94588, USA
Telephone: +1-925-223-8242
Fax: +1-925-223-8243
E-mail: bpgoffice@wjgnet.com
Help Desk: <http://www.f6publishing.com/helpdesk>
<http://www.wjgnet.com>



ISSN 1007-9327

