



35878-Review.doc

Quotes Excluded
Bibliography Excluded9%
SIMILARName of Journal: *World Journal of Obstetrics and Gynecology*

Manuscript NO: 35878

Manuscript Type: Original Article

Retrospective Cohort Study

Observed prevalence and risk factors of birth defects in Shanghai, China

Min Yang, Xiao-Xia An, Hui-Juan Wang, Ji-Mei Wang

Abstract

AIM—To investigate the prevalence and related risk factors of birth defects in Shanghai.

MATERIALS AND METHODS—This report describes a population-based study of all births at the Obstetrics and Gynecology Hospital of Fudan University in Shanghai, China from January 2008 to December 2014. A logistic

Match Overview

- | Rank | Source | Words | Match % |
|------|----------|--|---------|
| 1 | Internet | 95 words
crawled on 29-Dec-2016
bmcpediatr.biomedcentral.com | 2% |
| 2 | Crossref | 53 words
Liu, Jufen, Le Zhang, Zhiwen Li, Lei Jin, Yali Zhang, Rongwei Ye, Jianmeng Liu, and Aiguo Ren. "Prevalence and tren | 1% |
| 3 | Crossref | 51 words
Ge Sun. "Twelve-year prevalence of common neonatal c...
genital malformations in Zhejiang Province, China", World | 1% |
| 4 | Crossref | 50 words
Yang, Mei, Shaoqiang Zhang, and Yukai Du. "Epidemiolog
y characteristics of birth defects in Shenzhen city during ... | 1% |
| 5 | Crossref | 38 words
Cambra, K., B. Ibanez, D. Urzelai, I. Portillo, I. Montoya, S. Esnaola, and F. B. Cirarda. "Trends in the prevalences of c | 1% |
| 6 | Internet | 27 words
crawled on 12-Aug-2017
www.scilit.net | 1% |



[全部](#)[图片](#)[新闻](#)[视频](#)[地图](#)[更多](#)[设置](#)[工具](#)

找到约 80,700 条结果 (用时 0.52 秒)

Google 学术 : Observed prevalence and risk factors of Birth defects in Shanghai ,China

... or without cleft palate in Shanghai, China, from 1980 to ... - Cooper - 被引用次数 : 155

... -control study of paternal smoking and birth defects - Zhang - 被引用次数 : 88

... vary with season in Chinese women of childbearing ... - Ronnenberg - 被引用次数 : 107

Prevalence and risk factors of congenital heart defects among live ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5237335/> - 翻译此页

作者 : L Pei - 2017 - 被引用次数 : 3 - 相关文章

2017年1月13日 - Prevalence and risk factors of congenital heart defects among live births: a ... In China, CHD is the most prevalent birth defect, accounting for over 10% of corresponding to an overall observed prevalence of 187.2 (95% CI: 14.8, ... (82.8 per 10000), but lower than that in Shanghai (187.2 per 1000 births).

Observed Prevalence of Congenital Heart Defects From a ... - NCBI - NIH

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - 翻译此页

作者 : Y Zhang - 2011 - 被引用次数 : 25 - 相关文章

The prevalence of all congenital heart defects in the 4 areas of China studied was ... Congenital heart defects are the leading cause of infant mortality due to birth defects, of Participants and Nonparticipants in Study Areas Near Shanghai, China, Sonographic examinations were performed regardless of risk factors or ...

Prevalence of Congenital Heart Disease in Xinjiang Multi-Ethnic ...

journals.plos.org/plosone/article?id=10.1371/journal.pone.0133961 ▾ 翻译此页

作者 : F Liu - 2015 - 被引用次数 : 6 - 相关文章

全部 图片 视频 新闻 地图 图书

找到约 105,000 条结果

Google 学术 : Observed prevalence and risk factors of birth defects in Shanghai, China

... or without cleft palate in Shanghai, China, from 1980 to ... - Cooper - 被引用次数 : 158

... -control study of paternal smoking and birth defects - Zhang - 被引用次数 : 91

... vary with season in Chinese women of childbearing ... - Ronnenberg - 被引用次数 : 107

Prevalence and risk factors of congenital heart defects ... - NCBI - NIH

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5237335/>

13 Jan 2017 ... **Prevalence and risk factors** of congenital heart defects among live births: a ... In **China**, CHD is the most prevalent **birth defect**, accounting for over 10% of ... corresponding to an overall **observed prevalence** of 187.2 (95% CI: 14.8, ... (82.8 per 10000), but lower than that in **Shanghai** (187.2 per 1000 births).

Prevalence and risk factors of congenital heart defects among live ...

https://www.researchgate.net/.../312362866_Prevalence_and_risk_factors_of_congenital_heart_defects_among_live_births_A_population-based_cross-...

17 Jan 2017 ... **Prevalence and risk factors** of congenital heart defects among live births: A population-based cross-sectional survey in Shaanxi province In **China**, CHD is the most prevalent **birth defect**, **Shanghai**, a previous surveillance program identified 75 ... an overall **observed prevalence** of 187.2 (95% CI: 14.8, ..

Observed Prevalence of Congenital Heart Defects From a ...

pubmedcentralcanada.ca/pmc/articles/PMC4469985/

Congenital heart defects are the leading cause of infant mortality due to **birth defects**. of Participants and Nonparticipants in Study Areas Near **Shanghai, China**, Sonographic examinations were performed regardless of **risk factors** or ...

[全部](#)[图片](#)[新闻](#)[视频](#)[地图](#)[更多](#)[设置](#)[工具](#)

找到约 64,000 条结果（用时 0.55 秒）

Google 学术: Observed prevalence and risk factors of birth defects in Shanghai, China

... or without cleft palate in Shanghai, China, from 1980 to ... - Cooper - 被引用次数: 159

... -control study of paternal smoking and birth defects - Zhang - 被引用次数: 91

... vary with season in Chinese women of childbearing ... - Ronnenberg - 被引用次数: 107

Prevalence and risk factors of congenital heart defects ... - NCBI - NIH

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5237335/> - 翻译此页

作者: L Pei - 2017 - 被引用次数: 3 - 相关文章

2017年1月13日 - Prevalence and risk factors of congenital heart defects among live ... In China, CHD is the most prevalent birth defect, accounting for over In four areas surrounding Shanghai, a previous surveillance ... corresponding to an overall observed prevalence of 187.2 (95% CI: 14.8, 23.5) per 10000 births [7].

Prevalence and risk factors of congenital heart defects ... - ResearchGate

https://www.researchgate.net/.../312362866_Prevalence_and_risk_factors_of... - 翻译此页

2017年1月17日 - Prevalence and risk factors of congenital heart defects among live births: A population-based cross-sectional survey in Shaanxi province In China, CHD is the most prevalent birth defect, Shanghai, a previous surveillance program identified 75 ... an overall observed prevalence of 187.2 (95% CI: 14.8, ...

Prevalence of Congenital Heart Disease in Xinjiang Multi-Ethnic ...

journals.plos.org/plosone/article?id=10.1371/journal.pone.0133961 ▼ 翻译此页

作者: F Liu - 2015 - 被引用次数: 9 - 相关文章

2015年8月28日 - Background The prevalence and risk factors of congenital heart disease ... heart disease (CHD) in Xinjiang, northwestern part of China. ... of fallot (TOF, 4.2%) were common cyanotic and cyanotic