Primary pancreatic paraganglioma harboring lymph node metasta:

■ ♀ Q



ALL

VIDEOS

13.000 Results

IMAGES

Any time ▼

Pancreatic gangliocytic paraganglioma harboring lymph node ...

https://tohoku.pure.elsevier.com/en/publications/... ▼

Background: Gangliocytic paraganglioma (GP) is a rare neuroendocrine neoplasm, which occurs mostly in the periampullary portion of the duodenum; the majority of the reported cases of duodenal GP has been of benign nature with a low incidence of regional lymph node metastasis. GP arising from the pancreas is extremely rare. To date, only three cases have been reported and its clinical characteristics are largely unknown. Case presentation: A nodule located in the pancreatic ...

Cited by: 5 Author: Keisuke Nonaka, Yoko Matsuda, Akira Oka...

Publish Year: 2017

CT and MR Imaging Findings of Pancreatic Paragangliomas: A ...


https://europepmc.org/article/MED/26945413 ▼

Mar 01, 2016 · Currently, lymph node metastasis and metastases in other organs are still definite $evidence\ for\ the\ confirmation\ of\ malignant\ paraganglioma.\ 26\ Among\ the\ previously\ reported\ 3\ \textbf{cases}\ of$ malignant pancreatic paragangliomas, 2 cases showed only 1 lymph node metastasis, whereas the remaining case presented with multiple intrahepatic metastases. 16,17,19 In the case report by Al ...

Author: Wenjie Liang, Shunliang Xu Cited by: 5 Publish Year: 2016 Estimated Reading Time: 8 mins

Malignant extra-adrenal pancreatic paraganglioma: case ...

https://europepmc.org/abstract/MED/24138700 •





Primary pancreatic paraganglioma harboring lymph node metastas







ALL

IMAGES

VIDEOS

13,000 Results

Any time -

Pancreatic gangliocytic paraganglioma harboring lymph ...

https://tohoku.pure.elsevier.com/en/publications/... •

Background: Gangliocytic paraganglioma (GP) is a rare neuroendocrine neoplasm, which occurs mostly in the periampullary portion of the duodenum; the majority of the reported cases of duodenal GP has been of benign nature with a low incidence of regional lymph node metastasis. GP arising from the pancreas is extremely rare. To date, only three cases have been reported and its clinical characteristics are largely unknown. Case presentation: A nodule located in the pancreatic ...

Cited by: 5 Author: Keisuke Nonaka, Yoko Matsuda, Akira O...

Publish Year: 2017

CT and MR Imaging Findings of Pancreatic Paragangliomas: ...

https://europepmc.org/article/MED/26945413 -

Mar 01, 2016 · Currently, **lymph node** metastasis and metastases in other organs are still definite evidence for the confirmation of malignant paraganglioma. 26 Among the previously reported 3 cases of **malignant pancreatic paragangliomas**, 2 cases showed only 1 lymph node metastasis, whereas the remaining case presented with multiple intrahepatic metastases. 16,17,19 In the case report by Al ...

Cited by: 5 Author: Wenjie Liang, Shunliang Xu Publish Year: 2016 Estimated Reading Time: 8 mins

(PDF) CT and MR Imaging Findings of Pancreatic ...

https://www.researchgate.net/publication/297609754...

Pancreatic gangliocytic paraganglioma harboring lymph node metastasis: A case report and literature review ... One of these cases demonstrates metastasis to lymph node, while another case ...

Estimated Reading Time: 7 mins

Malignant extra-adrenal pancreatic paraganglioma: case ...

https://europepmc.org/abstract/MED/24138700 -

Oct 20, 2013 - In the previously reported 16 primary pancreatic paragangliomas, there was only one case that was considered malignant with lymph node metastasis and one case that showed hormonal activity [1,5]. In this report, we present a malignant pancreatic paraganglioma in a 19-year-old woman.