

Prognostic value of Onodera's nutritional index for ...

https://wiso.biomedcentral.com/articles/10.1186/s12957-021-02345-9 -

Aug 03, 2021 - Onodera's prognostic nutritional index (OPNI) was initially used to evaluate the immune-nutritional state of patients who are given gastrointestinal surgery. Several studies have shown that OPNI is a crucial prognostic factor in some specific human cancers, such as gastric cancer [19], pancreatic cancer [20], colorectal cancer [21], and esophageal cancer [22].

[PDF] Prognostic value of Onodera's nutritional index for ...

https://wjso.biomedcentral.com/track/pdf/10.1186/s12957-021-02345-9

ratio (NLR), platelet-to-lymphocyte ratio (PLR) and Onodera's prognostic nutritional index (OPNI), have gained consid- erable attention and have been preliminarily revealed as prognostic markers of gastrointestinal stromal tumors (GISTs).

Onodera's prognostic nutritional index correlates with ...

https://www.sciencedirect.com/science/article/pii/S1936523320303429

Dec 01, 2020 - Dec 01, 2020 - The Onodera prognostic nutritional index (OPNI) is an indicator calculated from serum albumin and total lymphocyte counts in peripheral blood, which can be utilized to evaluate patients' nutritional and immune status.

Cited by: 2 Author: Ryoji Yoshida, Shunsuke Gohara, Junki ...

Publish Year: 2020

Pretreatment prognostic nutritional index is a prognostic ...

https://jtd.amegroups.com/article/view/44712/html -

Onodera et al. proposed the concept of prognostic nutritional index (PNI) which is the mostly used marker of nutritional status. PNI is based on serum albumin concentrations and total lymphocyte counts in the peripheral blood and evaluates the potential impact of hypoalbuminemia and lymphocytopenia.

Cited by: 1 Author: Jingjing Liu, Shuang Li, Shuang Zhang, C...

Publish Year: 2020

[PDF] tumors treated with or without Tyrosine Kinase ...

https://www.researchgate.net/publication/350208696...

Prognostic value of Onodera's Nutritional Index for intermediate and high risk gastrointestinal stromal tumors treated with or without Tyrosine Kinase Inhibitors Feng Wang



Onodera's Prognostic Nutritional Index is a novel and useful progn







ALL

IMAGES

VIDEOS

3,810 Results

Any time -



wjso.biomedcentral.com - Aug 03, 2021

Prognostic value of Onodera's nutritional index for ...

Onodera's prognostic nutritional index (OPNI) was initially used to evaluate the immune-nutritional state of patients who are given gastrointestinal surgery . Several studies have shown that OPNI is a crucial prognostic factor in some specific human cancers, such as gastric cancer [19], pancreatic cancer [20], colorectal cancer [21], and esophageal cancer [22].

https://wjso.biomedcentral.com/articles/10.1186/s12957-021-02345-9 -

[PDF] Prognostic value of Onodera's nutritional index for ...

https://wjso.biomedcentral.com/track/pdf/10.1186/s12957-021-02345-9

ratio (NLR), platelet-to-lymphocyte ratio (PLR) and Onodera's prognostic nutritional index (OPNI), have gained consid- erable attention and have been preliminarily revealed as prognostic markers of gastrointestinal stromal tumors (GISTs).

Onodera's prognostic nutritional index correlates with ...

https://www.sciencedirect.com/science/article/pii/S1936523320303429

Dec 01, 2020 · Pretreatment nutritional and immunological status is useful for predicting survival outcomes for various types of malignant tumors. Our objective was to determine the impact of the pretreatment Onodera's prognostic nutritional index (OPNI) on outcomes of patients who underwent definitive chemoradiotherapy for advanced oral squamous cell carcinoma (OSCC).

Cited by: 2 Author: Ryoji Yoshida, Shunsuke Gohara, Junki ...

Publish Year: 2020

Pretreatment prognostic nutritional index is a prognostic ...

https://jtd.amegroups.com/article/view/44712/html -

Onodera et al. proposed the concept of prognostic nutritional index (PNI) which is the mostly used marker of nutritional status. PNI is based on serum albumin concentrations and total lymphocyte counts in the peripheral blood and evaluates the potential impact of bypoalbuminemia and lymphocytopenia