

[全部](#)[新闻](#)[图片](#)[视频](#)[购物](#)[更多](#)[设置](#)[工具](#)

找到约 11,500 条结果 (用时 0.70 秒)

[PDF] Critical Limb Ischemia Medtech Commercial Model - LimFlow

www.limflow.com/wp-content/uploads/pdf/Ischemia%20LimFlow.pdf - 翻译此页

2017年11月27日 - OUTSIDE OPINION. **Rethinking** the ... The minimally invasive **treatment** of peripheral **vascular** disease began much of it must be **treated** below the knee (**BTK**), which lim- its device ... which was founded in 2013, has raised \$3.6 million for the **de-** ... Flow is in business to give them a third and **better option**..

Luca Farchioni's research works | Azienda Ospedaliera Santa Maria ...

https://www.researchgate.net/scientific.../71601426_Luca_Farchioni - 翻译此页

Its prompt diagnosis and **treatment** are mandatory to restore renal function. ... Djurre **de** Waard In the contemporary **endovascular** era the decision to deny repair arbitrarily to older patients with rAAAs must be **revisited**. From available data, the **best** results of **endovascular** first **approach** for **BTK** lesions can be achieved ...

Drug-coated balloons and their place in treating peripheral arterial ...

www.tandfonline.com/doi/full/10.1586/erd.12.67?scroll=top...true - 翻译此页

2014年1月9日 - While it revolutionized the **treatment** of **vascular** stenosis and occlusion, the ... soon attracted great interest as an **alternative treatment** to plain balloon angioplasty and stent implantation. ... The authors will **revisit** this topic below. In short below the knee (**BTK**) lesions, **DES** have proven to yield statistically ...

[PDF] cirse 2017 - cSlide

<https://cslide.ctimeetingtech.com/cirse2017/confcal?external>

作者: C RSE - 相关文章

embolization (**DE**) requiring treatment 7.3%, stent thrombosis 4.9% and planned SRFA may even be a **good alternative** to surgical resection of focal liver lesions in mistic report, the results of drug-eluting balloons **BTK** have been disappointing. Hemodialysis arteriovenous fistula patency

Match Overview



There are no matching sources for this report.

1

1 **Revisiting endovascular treatment in below-the-knee disease. Are drug-eluting stents**
2 **the best option?**

3
4 Spiliopoulos S *et al.* Infrapopliteal DES

5
6 Stavros Spiliopoulos, Panagiotis Kitrou, Elias Brountzos

7
8 **Abstract**

9 Patients with below the knee arterial disease are mainly individuals suffering from
10 critical limb ischemia (CLI), while a large percentage of these patients are also suffering
11 from diabetes or chronic renal failure or both. Available data from randomized
12 controlled trials and their metaanalysis demonstrated that the use of infrapopliteal
13 drug-eluting stents (DES), in short- to medium- length lesions, obtains significantly

全部

新闻

视频

图片

更多

设置

工具

找到约 53,300 条结果 (用时 0.53 秒)

Drug eluting stents in the treatment of below the knee arterial ... - NCBI

<https://www.ncbi.nlm.nih.gov/pubmed/23640354> ▼ 翻译此页

作者: RL Varcoe - 2013 - 被引用次数: 6 - 相关文章

Drug eluting stents in the treatment of below the knee arterial occlusive disease. ... The use of endovascular techniques for the treatment of occlusive arterial ... Arterial Occlusive Diseases/physiopathology; Arterial Occlusive Diseases/surgery* ...

缺少字词: revisiting best option

Drug-eluting stents below the knee. - NCBI

<https://www.ncbi.nlm.nih.gov/pubmed/21460773> ▼ 翻译此页

作者: M Bosiers - 2011 - 被引用次数: 17 - 相关文章

(1)Department of Vascular Surgery, A.Z. Sint-Blasius, Dendermonde, Belgium. ... Drug-eluting stent (DES) technology was developed to prevent early thrombosis and ... Arterial Disease/physiopathology; Peripheral Arterial Disease/therapy* ...

缺少字词: revisiting best option

[PDF] Below-the-Knee Drug-Eluting Stents - Endovascular Today

https://evtoday.com/pdfs/EVT0811_feature_feiring.pdf ▼ 翻译此页

2011年8月1日 - presence of severe below-the-knee (BTK) arteri- ... When evaluating the worth of therapeutic options, we ... angiogram 18 months after treating in-stent restenosis with a 3.5- X 23-mm Cypher lesion length, patient gender, chronic kidney disease, dia- ... better at 50% for Rutherford class 4 and 5 patients. A.

缺少字词: revisiting

Endovascular Today - Below-the-Knee Drug-Eluting Stents (August ...

<https://evtoday.com/2011/08/below-the-knee-drug-eluting-stents/> ▼ 翻译此页

When evaluating the worth of therapeutic options, we have come to rely on the ... an abundance of evidence supporting its efficacy and safety: drug-eluting stents (DES). ... Because a comprehensive review of data on DES for treating BTK CLI is Infragenicular stent implantation for belowthe- knee atherosclerotic disease: ...

缺少字词: revisiting

找到约 89,500 条结果 (用时 0.53 秒)

(PDF) Endovascular treatment of infrapopliteal... - ResearchGate

https://www.researchgate.net/.../259985375_Endovascular_treatment_of_inf... - 翻译此页

Endovascular treatment of infrapopliteal arteries: Angioplasty vs stent in the ... On the basis of the results available, the drug-eluting devices produced **better** results. DEB can be considered the leading approach in below-the-knee disease. ... Drug-coated balloons and drug-eluting stents: Clinical effectiveness revisited.

Drug-eluting stents below the knee. - NCBI

<https://www.ncbi.nlm.nih.gov/pubmed/21460773> ▼ 翻译此页

作者: M Bosiers - 2011 - 被引用次数: 17 - 相关文章

(1)Department of Vascular Surgery, A.Z. Sint-Blasius, Dendermonde, Belgium. ... Drug-eluting stent (DES) technology was developed to prevent early thrombosis and ... Arterial Disease/physiopathology; Peripheral Arterial Disease/therapy* ...

缺少字词: revisiting best option

Drug eluting stents in the treatment of below the knee arterial ... - NCBI

<https://www.ncbi.nlm.nih.gov/pubmed/23640354> ▼ 翻译此页

作者: RL Varcoe - 2013 - 被引用次数: 6 - 相关文章

Drug eluting stents in the treatment of below the knee arterial occlusive disease. ... The use of endovascular techniques for the treatment of occlusive arterial ... Arterial Occlusive Diseases/physiopathology; Arterial Occlusive Diseases/surgery* ...

缺少字词: revisiting best option

Evidence for the use of drug eluting stents in below-the-knee lesions.

<https://www.ncbi.nlm.nih.gov/pubmed/25428562> ▼ 翻译此页

作者: D Trombert - 2015 - 被引用次数: 7 - 相关文章

2014年11月27日 - Evidence for the use of drug eluting stents in below-the-knee lesions. ... Peripheral arterial disease has become more and more present in daily practice, ... Nowadays, endovascular treatment is usually the first choice, given the high ... Angiographic results are better, compared to BMS, in terms of primary ...

缺少字词: revisiting

Drug-Eluting Balloons are Better for Below-Knee Lesions Than ...

<https://www.cathlabdigest.com/.../Drug-Eluting-Balloons-are-Better-Below-K...> ▼ 翻译此页

Drug-Eluting Balloons are Better for Below-Knee Lesions Than Sirolimus-Eluting Stents ... sirolimus-eluting stents as a primary treatment for below-the-knee lesions? ... in below-the-knee lesions (with the Xience stent from Abbott Vascular), the ... and regardless of the baseline vessel condition, whether it is