

World Journal of *Gastroenterology*

World J Gastroenterol 2020 October 7; 26(37): 5534-5744



OPINION REVIEW

- 5534 Review of inflammatory bowel disease and COVID-19
Sultan K, Mone A, Durbin L, Khuwaja S, Swaminath A

REVIEW

- 5543 Hepatitis E virus: Epidemiology, diagnosis, clinical manifestations, and treatment
Aslan AT, Balaban HY
- 5561 Transjugular intrahepatic portosystemic shunt in cirrhosis: An exhaustive critical update
Rajesh S, George T, Philips CA, Ahamed R, Kumbar S, Mohan N, Mohanan M, Augustine P

MINIREVIEWS

- 5597 Calcifying fibrous tumor of the gastrointestinal tract: A clinicopathologic review and update
Turbiville D, Zhang X
- 5606 Artificial intelligence technologies for the detection of colorectal lesions: The future is now
Attardo S, Chandrasekar VT, Spadaccini M, Maselli R, Patel HK, Desai M, Capogreco A, Badalamenti M, Galtieri PA, Pellegatta G, Fugazza A, Carrara S, Anderloni A, Occhipinti P, Hassan C, Sharma P, Repici A
- 5617 Application of artificial intelligence in the diagnosis and treatment of hepatocellular carcinoma: A review
Jiménez Pérez M, Grande RG

ORIGINAL ARTICLE**Basic Study**

- 5629 Antioxidant activity and hepatoprotective effect of 10 medicinal herbs on CCl₄-induced liver injury in mice
Meng X, Tang GY, Liu PH, Zhao CJ, Liu Q, Li HB

Case Control Study

- 5646 Short- and long-term outcomes associated with enhanced recovery after surgery protocol vs conventional management in patients undergoing laparoscopic gastrectomy
Tian YL, Cao SG, Liu XD, Li ZQ, Liu G, Zhang XQ, Sun YQ, Zhou X, Wang DS, Zhou YB

Retrospective Cohort Study

- 5661 Periodontitis combined with smoking increases risk of the ulcerative colitis: A national cohort study
Kang EA, Chun J, Kim JH, Han K, Soh H, Park S, Hong SW, Moon JM, Lee J, Lee HJ, Park JB, Im JP, Kim JS

Retrospective Study

- 5673 Preliminary experience of hybrid endoscopic submucosal dissection by duodenoscope for recurrent laterally spreading papillary lesions

Wang ZK, Liu F, Wang Y, Wang XD, Tang P, Li W

- 5682 *Helicobacter pylori* infection with atrophic gastritis: An independent risk factor for colorectal adenomas

Chen QF, Zhou XD, Fang DH, Zhang EG, Lin CJ, Feng XZ, Wang N, Wu JS, Wang D, Lin WH

Clinical Trials Study

- 5693 Endoscopic ultrasound-fine needle biopsies of pancreatic lesions: Prospective study of histology quality using Franseen needle

Stathopoulos P, Pehl A, Breitling LP, Bauer C, Grote T, Gress TM, Denkert C, Denzer UW

Prospective Study

- 5705 Risk prediction rule for advanced neoplasia on screening colonoscopy for average-risk individuals

Sharara AI, El Mokahal A, Harb AH, Khalaf N, Sarkis FS, M El-Halabi M, Mansour NM, Malli A, Habib R

EVIDENCE-BASED MEDICINE

- 5718 Endoscopic retrograde cholangiopancreatography in the treatment of pancreaticopleural fistula in children

Zhang J, Gao LC, Guo S, Mei TL, Zhou J, Wang GL, Yu FH, Fang YL, Xu BP

CASE REPORT

- 5731 Abernethy syndrome in Slovenian children: Five case reports and review of literature

Peček J, Fister P, Homan M

ABOUT COVER

Editorial Board of *World Journal of Gastroenterology*, Dr. Angelo Zambam de Mattos is a Professor of Medicine – Gastroenterology at the Federal University of Health Sciences of Porto Alegre (UFCSPA), where he is also a permanent faculty member of the Graduate Program in Medicine: Hepatology (the only Brazilian graduate program specialized specifically in Hepatology). His research focuses on cirrhosis and its complications, culminating in > 50 academic papers. He also carries out clinical work at Irmandade Santa Casa de Misericórdia of Porto Alegre, one of the largest hospital complexes in southern Brazil. Prof. Mattos received his Medical degree in 2005, Master's degree in 2012 and PhD in 2015, all from UFCSPA. He is a member of the Brazilian Federation of Gastroenterology, Brazilian Association of Hepatology, and Brazilian Association of Digestive Endoscopy, and he is past president of the Gastroenterology Association of Rio Grande do Sul, Brazil (2017-2018). (L-Editor: Filipodia)

AIMS AND SCOPE

The primary aim of *World Journal of Gastroenterology* (*WJG, World J Gastroenterol*) is to provide scholars and readers from various fields of gastroenterology and hepatology with a platform to publish high-quality basic and clinical research articles and communicate their research findings online. *WJG* mainly publishes articles reporting research results and findings obtained in the field of gastroenterology and hepatology and covering a wide range of topics including gastroenterology, hepatology, gastrointestinal endoscopy, gastrointestinal surgery, gastrointestinal oncology, and pediatric gastroenterology.

INDEXING/ABSTRACTING

The *WJG* is now indexed in Current Contents®/Clinical Medicine, Science Citation Index Expanded (also known as SciSearch®), Journal Citation Reports®, Index Medicus, MEDLINE, PubMed, PubMed Central, and Scopus. The 2020 edition of Journal Citation Report® cites the 2019 impact factor (IF) for *WJG* as 3.665; IF without journal self cites: 3.534; 5-year IF: 4.048; Ranking: 35 among 88 journals in gastroenterology and hepatology; and Quartile category: Q2.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: *Yu-Jie Ma*; Production Department Director: *Xiang Li*; Editorial Office Director: *Ze-Mao Gong*.

NAME OF JOURNAL

World Journal of Gastroenterology

ISSN

ISSN 1007-9327 (print) ISSN 2219-2840 (online)

LAUNCH DATE

October 1, 1995

FREQUENCY

Weekly

EDITORS-IN-CHIEF

Andrzej S Tarnawski, Subrata Ghosh

EDITORIAL BOARD MEMBERS

<http://www.wjgnet.com/1007-9327/editorialboard.htm>

PUBLICATION DATE

October 7, 2020

COPYRIGHT

© 2020 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjgnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>

Application of artificial intelligence in the diagnosis and treatment of hepatocellular carcinoma: A review

Miguel Jiménez Pérez, Rocío González Grande

ORCID number: Miguel Jiménez Pérez [0000-0001-7550-1019](https://orcid.org/0000-0001-7550-1019); Rocío González Grande [0000-0002-7691-5755](https://orcid.org/0000-0002-7691-5755).

Author contributions: Jiménez Pérez M and Grande RG contributed equally to this work.

Conflict-of-interest statement: Authors have no conflict of interest.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

Manuscript source: Invited manuscript

Received: July 8, 2020

Peer-review started: July 8, 2020

First decision: August 8, 2020

Revised: September 1, 2020

Accepted: September 18, 2020

Article in press: September 18, 2020

Miguel Jiménez Pérez, Rocío González Grande, UGC de Aparato Digestivo, Unidad de Hepatología-Trasplante Hepático, Hospital Regional Universitario de Málaga, Málaga 29010, Spain

Corresponding author: Miguel Jiménez Pérez, MD, PhD, Chief Doctor, UGC de Aparato Digestivo, Unidad de Hepatología-Trasplante Hepático, Hospital Regional Universitario de Málaga, Avenida Carlos Haya, Málaga 29010, Spain. mjimenezp@commalaga.com

Abstract

Although artificial intelligence (AI) was initially developed many years ago, it has experienced spectacular advances over the last 10 years for application in the field of medicine, and is now used for diagnostic, therapeutic and prognostic purposes in almost all fields. Its application in the area of hepatology is especially relevant for the study of hepatocellular carcinoma (HCC), as this is a very common tumor, with particular radiological characteristics that allow its diagnosis without the need for a histological study. However, the interpretation and analysis of the resulting images is not always easy, in addition to which the images vary during the course of the disease, and prognosis and treatment response can be conditioned by multiple factors. The vast amount of data available lend themselves to study and analysis by AI in its various branches, such as deep-learning (DL) and machine learning (ML), which play a fundamental role in decision-making as well as overcoming the constraints involved in human evaluation. ML is a form of AI based on automated learning from a set of previously provided data and training in algorithms to organize and recognize patterns. DL is a more extensive form of learning that attempts to simulate the working of the human brain, using a lot more data and more complex algorithms. This review specifies the type of AI used by the various authors. However, well-designed prospective studies are needed in order to avoid as far as possible any bias that may later affect the interpretability of the images and thereby limit the acceptance and application of these models in clinical practice. In addition, professionals now need to understand the true usefulness of these techniques, as well as their associated strengths and limitations.

Key Words: Artificial intelligence; Machine learning; Hepatocellular carcinoma; Diagnosis; Treatment; Prognosis

©The Author(s) 2020. Published by Baishideng Publishing Group Inc. All rights reserved.

Published online: October 7, 2020**P-Reviewer:** Azer S, Du Z, Gao YT**S-Editor:** Yan JP**L-Editor:** A**P-Editor:** Ma YJ

Core Tip: The biological variability in the behavior of hepatocellular carcinoma (HCC), conditioned by multiple factors, makes it difficult to establish general standards of action applicable equally to all patients. Analysis of the vast amount of data now available and their relation with tumor behavior is fundamental to be able to establish an efficient approach to HCC. It is here that the computational power of artificial intelligence can play a determining role, though it is necessary to understand the strengths and limitations of this technology before it can be applied in clinical practice.

Citation: Jiménez Pérez M, Grande RG. Application of artificial intelligence in the diagnosis and treatment of hepatocellular carcinoma: A review. *World J Gastroenterol* 2020; 26(37): 5617-5628

URL: <https://www.wjgnet.com/1007-9327/full/v26/i37/5617.htm>

DOI: <https://dx.doi.org/10.3748/wjg.v26.i37.5617>

INTRODUCTION

In the era of Big Data, the need for the efficient management of the great amount of information available has led to the development and application of artificial intelligence (AI) and its various techniques in the general field of medicine. Although the concept of AI arose in the 1950s^[1], it was not until a few years ago that it really came to experience its breakthrough.

The term AI refers to those computer programs that try to reproduce human cognitive functions, like learning or problem solving. Initially, machine learning (ML), developed as a branch of AI, analyzed data in order to create algorithms that can detect patterns of behavior from which predictive models can be established. Different ML techniques, such as support vector machines (SVM), artificial neural networks (ANNs) or classification and regression trees, have all been used in multiple studies in the field of medicine^[2]. Technological advances over the last ten years have resulted in the appearance of deep learning (DL) as a new model of ML to develop multi-layered neural network algorithms, using such techniques as convolutional neural network (CNN), a multilayer of ANN, that has proven of great use in the analysis of radiological images^[3,4].

Although the application of AI in various fields of medicine has shown promising results, we should nevertheless be aware of its limitations. The retrospective manner of many of these studies and the use of not particularly suitable databases, with their inherent bias, can affect the accuracy of AI. Thus, it is necessary to draw up prospective, well-designed, multicenter studies that are free from bias which could affect their interpretability and thereby limit their acceptance and application in clinical practice. And of course, we must not forget such other aspects as cost-effectiveness, regulations by the health authorities and ethical considerations.

AI has been used in the field of hepatology for the diagnosis, treatment and prognostic prediction of various different disorders, though with special relevance in the study of hepatocellular carcinoma (HCC) as this is a very common tumor. Estimates by the American Cancer Society put the number of new cases of liver and intrahepatic bile duct cancer during 2020 to be 42810, with 30160 deaths^[5]. HCC has particular radiological features that enable its diagnosis without the need for any histological study. Accordingly, the analysis of imaging tests takes on special relevance as their interpretation is not always easy, in addition to which they vary over the course of the disease, as do the prognosis and response to treatment, which are all affected by multiple factors. This all results in a vast amount of data whose integration and efficient analysis lend themselves to study by AI. Indeed, several studies have recently been undertaken to aid in the decision-making process and overcome the limitations of human evaluation^[4,6].

Figure 1 gives a schematic idea of how AI, with its variants, could be used to study a patient with HCC, whether diagnosed or suspected. AI is able to perform a combined analysis of radiological, clinical and histological data, producing information that can aid in the diagnostic accuracy, tumor staging, treatment planning using methods of segmentation and evaluation of the presence of microvascular invasion, in addition to giving a prognostic estimate.




Figure 1 Graphic presentation of the applications of artificial intelligence in the approach to hepatocellular carcinoma.

AI IN THE DIAGNOSIS OF HCC

In the field of liver cancer, the use of AI techniques to aid traditional diagnostic techniques is promising. CNN is a multilayer ANN interconnected in such a way that all input data traverse all the various layers during which the information is processed to produce output data. It can be considered an advanced form of DL with its own learning capacity. CNN can increase the diagnostic yield of ultrasound studies, abdominal computerized tomography or abdominal magnetic resonance imaging (MRI), positron emission tomography (PET) and histology.

Abdominal ultrasound

HCC usually, though not always, develops in a cirrhotic liver. Accordingly, clinical practice guidelines recommend regular abdominal ultrasound in patients with hepatic cirrhosis; indeed, it is considered the method of choice for screening of space-occupying lesions. Ultrasound is therefore the main tool to evaluate liver disease and detect new lesions. However, image interpretation is not always easy and may present interobserver variability.

To assess the underlying disease, Bharti *et al*^[7] proposed an ANN model to differentiate four stages of liver disease using data obtained from ultrasound images: normal liver, chronic liver disease, cirrhosis, and HCC. The classification accuracy of the model was 96.6%^[7]. Liu *et al*^[8] designed an algorithm to classify ultrasound images. They selected the liver capsule to determine the presence of cirrhosis, even in early stages when the usual findings reported by the radiologist, such as a nodular liver outline, enlarged porta or splenomegaly, are still not obvious. Using their analysis of the morphology of the liver capsule, they were able to determine the presence or absence of cirrhosis, with an area under the curve of 0.968^[8].

The human yield when characterizing a liver lesion from ultrasound images is limited. Schmauch *et al*^[9] designed a DL system able to detect and classify space-occupying lesions in the liver as benign or malignant. After a supervised training using a database of 367 images together with the radiological reports, the resulting algorithm detected and characterized the lesions with a mean receiver operating characteristic of 0.93 and 0.916, respectively. Although the system requires validation, it could increase the diagnostic yield of ultrasound and warn of possibly malignant lesions^[9].

AI has also been used in contrast-enhanced ultrasound (C-US), improving its ability to identify characteristics suggestive of cancer. Guo *et al*^[10] demonstrated that DL applied to the behavior of liver lesions seen on C-US in three phases (arterial, portal and late) increased the accuracy, sensitivity and specificity of the study^[10].

Abdominal computed tomography with IV contrast

When a follow-up ultrasound shows a new liver lesion, other imaging studies are undertaken, mainly dynamic contrast-enhanced computed tomography (CT) or MRI, to obtain a more precise evaluation. The radiological behavior of liver lesions in

dynamic CT or MRI studies is useful for characterization of the lesion. If a liver lesion > 1 cm fulfils certain radiological criteria, almost pathognomonic for HCC such as the presence of hyperenhancement at arterial phase and washout at portal or late phases in a cirrhotic patient, no further studies are needed for its diagnosis or histological confirmation. However, liver nodules often present an indeterminate behavior on CT and a biopsy of the lesion is required, as recommended in the European Association for the Study of the Liver guidelines^[11]. This, though, requires assuming the risks involved in the procedure, or close follow-up, as indicated in the American Association for the Study of Liver Diseases guidelines^[12], with a high number of studies and the possibility of not diagnosing a malignant lesion in time. Mokrane *et al.*^[13] undertook a retrospective analysis of 178 patients with cirrhosis and liver nodules in whom the Liver Reporting and Data System criteria were unable to distinguish the neoplastic from the non-neoplastic lesions, thereby necessitating a biopsy; 77% proved to be malignant on biopsy. Using DL techniques to classify nodules as HCC or non-HCC achieved an area under the curve (AUC) of 0.70. Another retrospective study, by Yasaka *et al.*^[14], analyzed the yield of an ANN, composed of three layers, classifying liver masses using contrast-enhanced CT into five categories: A, classic HCC; B, malignant tumors apart from HCC (cholangiocarcinoma, hepatocholangiocarcinoma or metastasis); C, indeterminate masses, dysplastic nodules or early HCC and benign masses other than cysts or hemangiomas; D, hemangiomas; E, cysts. After supervised training using over 55000 image sets the authors obtained a high accuracy for the classification of liver lesions, especially for the differentiation between categories A-B and C-D.

Quantifying the tumor load may be useful, particularly for the detection of tumor recurrence in follow-up CT studies. As tumor relapses can be small and go unnoticed, Vivanti *et al.*^[15] described an automated detection method of recurrence, based on the initial appearance of the tumor, its CT behavior, and the quantification of the tumor load at baseline and during the follow-up. The technique had a high rate of true positives in the identification of tumor recurrence, with an accuracy of 86%.

Liver segmentation is of great importance to assess liver lesions and for planning the ideal treatment. However, manual segmentation is made more difficult by the heterogeneity of the lesions or their diffuse borders. Li *et al.*^[16] proposed a CNN that can segment liver tumors based on CT images, with an accuracy of 82.67% ± 1.43%, better than that of traditional techniques, thereby favoring suitable treatment planning.

MRI of the abdomen

CNN applied to MRI has also been analyzed. Hamm *et al.*^[17] developed and validated a DL system based on a CNN that classifies MRI liver lesions, with an accuracy of 92%, a sensitivity of 92% and a specificity of 98%; and an average computation time of 5.6 ms.

Other studies have associated additional MRI sequences and risk factors plus the patient's clinical data to apply an automated classification system cataloguing liver lesions as adenoma, cyst, hemangioma, HCC and metastasis, with a sensitivity and specificity of 0.8/0.78, 0.93/0.93, 0.84/0.82, 0.73/0.56 and 0.62/0.77, respectively^[18]. Zhang *et al.*^[19] described a training model using MRI in 20 patients to classify liver tissues. Their results were promising, improving the yield of the reference models used.

PET

Preis *et al.*^[20] evaluated the yield of fluorine 18 fluorodeoxyglucose positron emission tomography/computed tomography (18F-FDG PET/CT) using a neural network to analyze liver uptake of 18F together with patient and laboratory data. They achieved a high sensitivity and specificity to detect liver malignancy unidentified visually, showing that this technique could complement the radiologist in the interpretation of PET, though their main aim was to evaluate metastatic liver disease, where 18F-FDG PET/CT has greater applicability.

Histology

The histopathological classification of a liver lesion and the differentiation of the tumor strain is crucial for treatment planning and a prognostic estimate of the disease, though this can sometimes prove challenging even for expert pathologists. Kiani *et al.*^[21] used AI as a support for the pathologists, focusing on the histological differentiation between HCC and cholangiocarcinoma. They analyzed prospectively the impact of this aid in the diagnostic yield of 11 pathologists, finding that it made no change to their mean accuracy.

Others have described how a deep CNN using previous histopathological images of HCC can make an automated diagnosis of HCC and distinguish healthy tissue from tumor tissue, in addition to identifying certain biological predictors^[22]. **Table 1** shows the main studies using AI techniques in the diagnosis of HCC.

AI IN THE TREATMENT OF HCC

The individual biological variability between patients in the behavior of HCC hinders evidence-based clinical assessment that is applicable to all patients. Accordingly, powerful, standardized risk stratification systems are needed in order to optimize treatment strategies and assess their effects. It is here that AI can play an important role in the therapeutic approach to HCC. Most studies on the use of AI in the treatment of HCC are aimed at the analysis of certain particular tumor characteristics, such as radiological, histological or genetic features, or a combination of the clinical data in order to predict the response to a particular treatment. This, in turn, will allow for the suitable selection of patients for particular treatment options.

AI strategies based on radiomics

In usual clinical practice the diagnosis and treatment assessment of HCC are commonly done with such imaging techniques as C-US, CT and MRI, after analyzing certain tumor features like vascularization or behavior after the administration of contrast material^[23]. However, these characteristics are liable to subjectivity in their interpretation by the radiologist, in addition to the lack of high resolution dimensional images. But a new technology has recently appeared in the area of radiology and cancer, radiomics^[24]. Although it is not yet in extensive use in clinical practice, it has nevertheless awoken great interest. This technology allows for the extraction of a great amount of quantifiable objective data contained in the radiological images and their later association with the underlying biological processes. Analysis of all these data with AI software can provide useful diagnostic and prognostic information with predictive accuracy^[24,25].

Evaluation of surgical resection

Early tumor recurrence after surgical resection is associated with a poor prognosis. The preoperative identification of patients at high risk of recurrence is fundamental to avoid unnecessary treatment. Computer models have been developed that analyze certain tumor characteristics and aid in the preoperative prediction of the risk of recurrence or the evaluation of survival after resection.

Vascular microinvasion (VMI) has been established as an independent predictive factor of recurrence, associated with poor results after tumor resection^[26]. Although the preoperative availability of information about VMI would be of much benefit, radiological techniques currently in clinical use do not provide for an adequate direct diagnosis.

Several studies have managed to elaborate radiomic signatures that enable prediction of the preoperative status of VMI, based on contrast-enhanced CT^[27,28] or MRI^[29]. However, these techniques involve radiological exposure, and are laborious to perform and costly. Recently, Dong *et al*^[30] published a study using radiomic algorithms based on grayscale ultrasound images to elaborate radiomic signatures with the potential to aid in the prediction of VMI, with promising results. Ji *et al*^[31] created predictive models for recurrence after surgical resection using radiomic techniques to analyze contrast-enhanced CT images, with a C-index of 0.633-0.699. In conjunction with the inclusion of clinical data, the model can be used to establish a personalized risk stratification facilitating the individual management of HCC.

Survival after surgical resection has also been assessed in several studies using ML techniques^[32-34], and more recently with more advanced DL models based on digitalized histological images of the tumor. Saillard *et al*^[35] drew up a predictive model of survival after resection, attaining a C-index for survival prediction of 0.78. A recent prospective study by Schoenberg *et al*^[36] involving 180 patients also led to a predictive model based on the analysis of 26 preoperative routine clinical variables, obtaining a predictive value of 0.78.

Evaluation of transcatheter arterial chemoembolization

Transcatheter arterial chemoembolization (TACE) is the treatment of choice for intermediate stage B HCC, in the Barcelona Clinical Liver Cancer (BCLC) classification^[37]. Adequate selection of patients who might benefit from this treatment

Table 1 Studies applying artificial intelligence in the diagnosis of hepatocellular carcinoma

Ref.	Title	Aim of the study of the use of AI in imaging techniques	Diagnostic technique studied	AI tool used	University/department
Bharti <i>et al</i> ^[7] 2018	Preliminary study of chronic liver classification on ultrasound images using an ensemble model	Classification of liver disease in four stages; normal liver, chronic liver disease, cirrhosis and HCC	Ultrasound	CNN	Thapar Institute of Engineering & Technology, Patiala, India
Liu <i>et al</i> ^[8] 2017	Accurate prediction of responses to transarterial chemoembolization for patients with hepatocellular carcinoma by using artificial intelligence in contrast-enhanced ultrasound	Early identification of the presence of cirrhosis	Ultrasound	ML	Sun Yat-sen University, Guangzhou, China
Schmauch <i>et al</i> ^[9] 2019	Diagnosis of focal liver lesions from ultrasound using deep learning	Classify liver lesions as benign or malignant	Ultrasound	DL	Owkin Inc, Research and Development Laboratory, Paris, France
Guo <i>et al</i> ^[10] 2018	A two-stage multi-view learning framework based computer-aided diagnosis of liver tumors with contrast enhanced ultrasound images	Characterize liver lesions and identify data of malignancy	C-US	ML	University School of Medicine, Shanghai, China
Mokrane <i>et al</i> ^[13] 2020	Radiomics machine-learning signature for diagnosis of hepatocellular carcinoma in cirrhotic patients with indeterminate liver nodules	Identify malignancy in hepatic space-occupying lesions catalogued as indeterminate	CT	Radiomics	Department of Radiology, New York Presbyterian Hospital, Columbia University Vagelos College of Physicians and Surgeons, New York City, NY, United States
Yasaka <i>et al</i> ^[14] 2018	Deep learning with convolutional neural network for differentiation of liver masses at dynamic contrast-enhanced CT: A preliminary study	Classification of liver lesions in five categories	CT	CNN	Department of Radiology, The University of Tokyo Hospital, Tokyo, Japan
Vivanti <i>et al</i> ^[15] 2017	Automatic detection of new tumors and tumor burden evaluation in longitudinal liver CT scan studies	Detection of tumor recurrence analyzing volume/tumor load	CT	CNN	The Rachel and Selim Benin School of Computer Science and Engineering, The Hebrew University of Jerusalem, Jerusalem, Israel
Li <i>et al</i> ^[16] 2015	Automatic segmentation of liver tumor in CT Images with deep convolutional neural networks	Liver tumor segmentation	CT	CNN	Research Lab for Medical Imaging and Digital Surgery, Shenzhen Institutes of Advanced Technology, Chinese Academy of Sciences, Shenzhen, China
Hamm <i>et al</i> ^[17] 2019	Deep learning for liver tumor diagnosis part I: development of a convolutional neural network classifier for multi-phasic MRI	Classification of liver lesions	MRI	DL	Department of Radiology and Biomedical Imaging, Yale School of Medicine, United States
Jansen <i>et al</i> ^[18] 2019	Automatic classification of focal liver lesions based on MRI and risk factors	Classification of liver lesions in: Adenomas, cysts, hemangiomas, HCC and metastasis	MRI	ML	Image Sciences Institute, University Medical Center Utrecht & Utrecht University, Utrecht, the Netherlands
Zhang <i>et al</i> ^[19] 2018	Liver tissue classification using an auto-context-based deep neural network with a multi-phase training framework	Classification of liver tissue	MRI	CNN	Department of Biomedical Engineering, Yale University, New Haven, CT, United States
Preis <i>et al</i> ^[20] 2011	Neural network evaluation of pet scans of the liver: A potentially useful adjunct in clinical interpretation	Identify metastatic liver disease	PET	CNN	Department of Radiology, Massachusetts General Hospital and Harvard Medical School, Boston
Kiani <i>et al</i> ^[21] 2020	Impact of a deep learning assistant on the histopathologic classification of liver cancer	Differentiate HCC from cholangiocarcinoma	Histology	DL	Department of Computer Science, Stanford University, Stanford, CA, United States
Liao <i>et al</i> ^[22] 2020	Deep learning-based classification and mutation prediction from histopathological images of hepatocellular carcinoma	Automated identification of liver tumor tissue, differentiating it from healthy	Histology	DL	Department of Liver Surgery & Liver Transplantation, State Key Laboratory of Biotherapy and Cancer Center, West China Hospital, Sichuan University and

Note: AI: Artificial intelligence; HCC: Hepatocellular carcinoma; CNN: Convolutional neural network; ML: Machine learning; DL: Deep learning; C-US: Contrast-enhanced ultrasound; CT: Computed tomography; MRI: Magnetic resonance imaging; PET: Positron emission tomography.

is vital in order to avoid unnecessary examinations that can sometimes have undesirable secondary effects for the patient and waste costs for the health system. Studies have been developed based on AI techniques to attempt to predict the response to treatment with TACE and aid adequate patient selection. Most of these studies are based on imaging analysis, though some have used genomic signatures. Morshid *et al*^[38] elaborated a fully automated ML algorithm using the combination of quantitative characteristics of CT images plus pretreatment patient clinical data to predict the response to TACE. They achieved a prediction accuracy rate of 74.2% using a combination of the BCLC stage plus quantitative image features *vs* using just the BCLC stage alone. Peng *et al*^[39] validated a DL model to predict the response to TACE using CT images from a total of 789 patients in three different hospitals. They obtained an accuracy of 84% and an AUC of 0.97 to predict complete response. Liu *et al*^[40] constructed and validated a DL model (DL radiomics-based C-US model) but based on the quantitative analysis of C-US cine recordings. It was highly reproducible and had an AUC of 0.93 (95%CI: 0.80-0.98) to predict the response to TACE.

Other studies have used ML techniques combining MRI with clinical data to predict the response to TACE. Abajian *et al*^[41] studied 36 patients who underwent MRI before TACE. They developed a predictive model of response with an accuracy of 78%, sensitivity of 62.5% and specificity of 82%.

The efficacy of TACE has also been examined by survival analysis of the patients after its application. Mähringer-Kunz *et al*^[42] developed a prediction model of survival after TACE by constructing an ANN, using all the parameters of the main conventional prediction scores (ART^[43], ABCR^[44] and SNACOR^[45]). They predicted a one-year survival with an AUC of 0.77, sensitivity of 78% and specificity of 81%, better results compared to those of the conventional scores mentioned.

Although most studies evaluating the use of AI to assess TACE have used radiomics, some have also evaluated the prediction of response to TACE using genetic analysis. Ziv *et al*^[46] studied genetic mutations using SVM techniques to predict the tumor response after TACE, though it was a retrospective study with a low number of cases.

Evaluation of radiofrequency ablation

Radiofrequency ablation (RFA) as a therapy aimed at curing HCC in early stages^[37] has also been evaluated. Liang *et al*^[47] drew up a predictive model of HCC recurrence based on SVM. They studied 83 patients with HCC who underwent RFA, obtaining an AUC of 0.69, sensitivity of 67% and specificity of 86%, with which they were able to identify patients at a high risk of recurrence. Table 2 summarizes the studies using AI

Table 2 Studies applying artificial intelligence for the treatment of hepatocellular carcinoma

Ref.	Title	n	Study type	Study aim	AI tool used	Outcome	University/department
Dong <i>et al</i> ^[30] 2020	Preoperative prediction of microvascular invasion in hepatocellular carcinoma: initial application of a radiomic algorithm based on grayscale ultrasound images	322	Retrospective	Prediction of VMI using C-US	Radiomics	AUC: 0.73; Sen: 0.919; Spe: 0.359	Department of Ultrasound, Zhongshan Hospital, Fudan University, Shanghai, China
Xu <i>et al</i> ^[28] 2019	Radiomic analysis of contrast-enhanced CT predicts microvascular invasion and outcome in hepatocellular carcinoma	495	Retrospective	Prediction of VMI using C-CT	Radiomics	AUC: 0.90	Department of Radiology, The First Affiliated Hospital with Nanjing Medical University, Nanjing, Jiangsu Province, China
Ma <i>et al</i> ^[27] 2019	Preoperative radiomics nomogram for microvascular invasion prediction in hepatocellular carcinoma using contrast-enhanced CT	157	Retrospective	Prediction of VMI using C-CT	Radiomics	AUC: 0.73	Department of Diagnostic Radiology, National Cancer Center/Cancer Hospital, Chinese Academy of Medical Sciences and Peking Union Medical College, China
Zhou <i>et al</i> ^[29] 2017	Malignancy characterization of hepatocellular carcinomas based on texture analysis of contrast-enhanced MR images	46	Retrospective	Prediction of VMI using C-MRI		AUC: 0.918; Sen: 92%; Spe: 66%	Laboratory for Health Informatics, Shenzhen Institutes of Advanced Technology, Chinese Academy of Sciences, Shenzhen, China
Ziv <i>et al</i> ^[46] 2017	Gene signature associated with upregulation of the Wnt/ β -Catenin signaling pathway predicts tumor response to transarterial embolization	17	Retrospective	Prediction of response to TACE using signature gene		Prediction accuracy: 70%	Interventional Radiology Service, Department of Radiology, Memorial Sloan Kettering Cancer Center, New York, NY, United States
Morshid <i>et al</i> ^[38] 2019	A machine learning model to predict hepatocellular carcinoma response to transcatheter arterial chemoembolization	105	Retrospective	Prediction of response to TACE using CT	ML	Acc: 74%	Departments of Imaging Physics, Diagnostic Radiology, Gastrointestinal Oncology and Interventional Radiology, The University of Texas, MD Anderson Cancer Center, Houston
Liu <i>et al</i> ^[40] 2020	Accurate prediction of responses to transarterial chemoembolization for patients with hepatocellular carcinoma by using artificial intelligence in contrast-enhanced ultrasound	130	Retrospective	Prediction of response to TACE using C-US	DL	AUC: 0.93	Department of Medical Ultrasonics, Institute of Diagnostic and Interventional Ultrasound, The First Affiliated Hospital of Sun Yat-sen University, China
Peng <i>et al</i> ^[39] 2020	Residual convolutional neural network for predicting response of transarterial chemoembolization in hepatocellular carcinoma from CT imaging	789	Retrospective	Prediction of response to TACE using CT	CNN	AUC: 0.97; Acc: 84.3%	Hepatology Unit and Department of Infectious Diseases, Nanfang Hospital, Southern Medical University, China
Abajian <i>et al</i> ^[41] 2018	Predicting treatment response to intra-arterial therapies for hepatocellular carcinoma with the use of supervised machine learning-an artificial intelligence concept	36	Retrospective	Prediction of response to TACE using MRI	ML	Acc: 78%; Sen: 62%; Spe: 82%	Yale School of Medicine, Department of Radiology and Biomedical Imaging, United States
Mähringer-Kunz <i>et al</i> ^[42] 2020	Predicting survival after transarterial chemoembolization for hepatocellular carcinoma using a neural network: A pilot study	282	Retrospective	Prediction of survival after TACE	CNN	Acc: 0.77; Sen: 78%; Spe: 81%	Department of Diagnostic and Interventional Radiology, University Medical Center of the Johannes Gutenberg-University Mainz, Mainz, Germany
Saillard <i>et al</i> ^[35] 2020	Predicting survival after hepatocellular carcinoma resection using deep-learning on histological slides	194	Retrospective	Prediction of survival after surgical resection	DL	C-index: 0.78	Owkin Lab, Owkin
Liang <i>et al</i> ^[47] 2014	Recurrence predictive models for patients with hepatocellular carcinoma after radiofrequency ablation using support vector machines with feature selection methods	83	Prospective	Prediction of recurrence after RFA	ML	AUC: 67%; Sen: 86%; Spe: 82%	Department of Internal Medicine, National Taiwan University Hospital and National Taiwan University College of Medicine, Taipei, Taiwan
Ji	Machine-learning analysis of contrast-enhanced CT radiomics	470	Retrospective	Prediction of	ML	C-index: 0.633-	Hepatobiliary Center, The First Affiliated Hospital of Nanjing

<i>et al</i> ^[31] 2019	predicts recurrence of hepatocellular carcinoma after resection: A multi-institutional study	recurrence after resection	0.699	Medical University, Nanjing, PR China
-----------------------------------	--	----------------------------	-------	---------------------------------------

Note: All studies were retrospective studies in their design, except the study by Liang *et al*^[47] was prospective. AI: Artificial intelligence; CNN: Convolutional neural network; ML: Machine learning; DL: Deep learning; VMI: Vascular microinvasion; C-US: Contrast-enhanced ultrasound; AUC: Area under the curve; Acc: Accuracy; Sen: Sensitivity; Spe: Specificity; CT: Computed tomography; C-CT: Contrast-enhanced CT; MRI: Magnetic resonance imaging; C-MRI: Contrast-enhanced MRI; TACE: Transcatheter arterial chemoembolization; RFA: Radiofrequency ablation.

techniques in the treatment of HCC.

PREDICTING OVERALL SURVIVAL OF HCC

The prediction of overall survival of HCC, apart from the application of any therapy, has also been assessed using AI techniques. Current evidence on the relation of abnormalities in DNA methylation and HCC^[48-50] was the basis for the study by Dong *et al*^[51]. These authors used ML techniques (SVM) to analyze DNA methylation data from 377 HCC samples, and constructed three risk categories to predict overall survival, with a mean 10-fold cross-validation score of 0.95.

FUTURE DIRECTIONS

Larger studies are needed comparing the yield of medical professionals with the support of AI *vs* other professionals without such support in order to demonstrate its benefit as an aid in medicine. In particular, to assess liver masses and study HCC these trials should focus on aspects related to treatment and prognosis, such as the characterization of hepatic lesions catalogued as indeterminate, the presence of vascular invasion and the response to percutaneous therapy. Another important aspect is the use of AI in the analysis of the behavior of HCC in cirrhotic and non-cirrhotic patients, as well as the differentiation of primary and metastatic liver lesions^[52] and especially the differential diagnosis with cholangiocarcinoma that can be complicated with currently available techniques and whose management and prognosis are completely different from those of HCC. At the same time, it is also necessary to start training health-care professionals to be prepared for the future incorporation of AI in daily practice in the field of liver cancer.

CONCLUSION

The incorporation of AI technologies in medicine has represented one of the most

relevant advances in recent years. It will doubtless experience a progressively increasing rise, due to its usefulness in the processing and analysis of the enormous amount of data currently available. Nevertheless, we should be aware that certain constraints still exist that can limit its acceptance and applicability in clinical practice. Health care professionals must learn the true usefulness of AI and accept the need for its coexistence with the indispensable need for human evaluation, accepting that AI is here to support human intelligence, never to replace it. Despite the great progress represented by AI, it is nevertheless vital to guarantee that medical protocols remain rigorously transparent.

REFERENCES

- 1 **Turing AM.** I.—Computing Machinery and Intelligence. *Mind* 1950; **LIX**: 433-460 [DOI: [10.1093/mind/LIX.236.433](https://doi.org/10.1093/mind/LIX.236.433)]
- 2 **Kaul V,** Enslin S, Gross SA. History of artificial intelligence in medicine. *Gastrointest Endosc* 2020 [PMID: [32565184](https://pubmed.ncbi.nlm.nih.gov/32565184/) DOI: [10.1016/j.gie.2020.06.040](https://doi.org/10.1016/j.gie.2020.06.040)]
- 3 **Yang YJ,** Bang CS. Application of artificial intelligence in gastroenterology. *World J Gastroenterol* 2019; **25**: 1666-1683 [PMID: [31011253](https://pubmed.ncbi.nlm.nih.gov/31011253/) DOI: [10.3748/wjg.v25.i14.1666](https://doi.org/10.3748/wjg.v25.i14.1666)]
- 4 **Le Berre C,** Sandborn WJ, Aridhi S, Devignes MD, Fournier L, Smaïl-Tabbone M, Danese S, Peyrin-Biroulet L. Application of Artificial Intelligence to Gastroenterology and Hepatology. *Gastroenterology* 2020; **158**: 76-94.e2 [PMID: [31593701](https://pubmed.ncbi.nlm.nih.gov/31593701/) DOI: [10.1053/j.gastro.2019.08.058](https://doi.org/10.1053/j.gastro.2019.08.058)]
- 5 **Siegel RL,** Miller KD, Jemal A. Cancer statistics, 2020. *CA Cancer J Clin* 2020; **70**: 7-30 [PMID: [31912902](https://pubmed.ncbi.nlm.nih.gov/31912902/) DOI: [10.3322/caac.21590](https://doi.org/10.3322/caac.21590)]
- 6 **Briceño J.** Artificial intelligence and organ transplantation: challenges and expectations. *Curr Opin Organ Transplant* 2020; **25**: 393-398 [PMID: [32487888](https://pubmed.ncbi.nlm.nih.gov/32487888/) DOI: [10.1097/MOT.0000000000000775](https://doi.org/10.1097/MOT.0000000000000775)]
- 7 **Bharti P,** Mittal D, Ananthasivan R. Preliminary Study of Chronic Liver Classification on Ultrasound Images Using an Ensemble Model. *Ultrason Imaging* 2018; **40**: 357-379 [PMID: [30015593](https://pubmed.ncbi.nlm.nih.gov/30015593/) DOI: [10.1177/0161734618787447](https://doi.org/10.1177/0161734618787447)]
- 8 **Liu X,** Song JL, Wang SH, Zhao JW, Chen YQ. Learning to Diagnose Cirrhosis with Liver Capsule Guided Ultrasound Image Classification. *Sensors (Basel)* 2017; **17**: 149 [PMID: [28098774](https://pubmed.ncbi.nlm.nih.gov/28098774/) DOI: [10.3390/s17010149](https://doi.org/10.3390/s17010149)]
- 9 **Schmauch B,** Herent P, Jehanno P, Dehaene O, Saillard C, Aubé C, Luciani A, Lassau N, Jégou S. Diagnosis of focal liver lesions from ultrasound using deep learning. *Diagn Interv Imaging* 2019; **100**: 227-233 [PMID: [30926443](https://pubmed.ncbi.nlm.nih.gov/30926443/) DOI: [10.1016/j.diii.2019.02.009](https://doi.org/10.1016/j.diii.2019.02.009)]
- 10 **Guo LH,** Wang D, Qian YY, Zheng X, Zhao CK, Li XL, Bo XW, Yue WW, Zhang Q, Shi J, Xu HX. A two-stage multi-view learning framework based computer-aided diagnosis of liver tumors with contrast enhanced ultrasound images. *Clin Hemorheol Microcirc* 2018; **69**: 343-354 [PMID: [29630528](https://pubmed.ncbi.nlm.nih.gov/29630528/) DOI: [10.3233/CH-170275](https://doi.org/10.3233/CH-170275)]
- 11 **European Association for the Study of the Liver.** EASL Clinical Practice Guidelines: Management of hepatocellular carcinoma. *J Hepatol* 2018; **69**: 182-236 [PMID: [29628281](https://pubmed.ncbi.nlm.nih.gov/29628281/) DOI: [10.1016/j.jhep.2018.03.019](https://doi.org/10.1016/j.jhep.2018.03.019)]
- 12 **Heimbach JK,** Kulik LM, Finn RS, Sirlin CB, Abecassis MM, Roberts LR, Zhu AX, Murad MH, Marrero JA. AASLD guidelines for the treatment of hepatocellular carcinoma. *Hepatology* 2018; **67**: 358-380 [PMID: [28130846](https://pubmed.ncbi.nlm.nih.gov/28130846/) DOI: [10.1002/hep.29086](https://doi.org/10.1002/hep.29086)]
- 13 **Mokrane FZ,** Lu L, Vavasseur A, Otal P, Peron JM, Luk L, Yang H, Ammari S, Saenger Y, Rousseau H, Zhao B, Schwartz LH, Dercle L. Radiomics machine-learning signature for diagnosis of hepatocellular carcinoma in cirrhotic patients with indeterminate liver nodules. *Eur Radiol* 2020; **30**: 558-570 [PMID: [31444598](https://pubmed.ncbi.nlm.nih.gov/31444598/) DOI: [10.1007/s00330-019-06347-w](https://doi.org/10.1007/s00330-019-06347-w)]
- 14 **Yasaka K,** Akai H, Abe O, Kiryu S. Deep Learning with Convolutional Neural Network for Differentiation of Liver Masses at Dynamic Contrast-enhanced CT: A Preliminary Study. *Radiology* 2018; **286**: 887-896 [PMID: [29059036](https://pubmed.ncbi.nlm.nih.gov/29059036/) DOI: [10.1148/radiol.2017170706](https://doi.org/10.1148/radiol.2017170706)]
- 15 **Vivanti R,** Szeskin A, Lev-Cohain N, Sosna J, Joskowicz L. Automatic detection of new tumors and tumor burden evaluation in longitudinal liver CT scan studies. *Int J Comput Assist Radiol Surg* 2017; **12**: 1945-1957 [PMID: [28856515](https://pubmed.ncbi.nlm.nih.gov/28856515/) DOI: [10.1007/s11548-017-1660-z](https://doi.org/10.1007/s11548-017-1660-z)]
- 16 **Li W,** Jia F, Hu Q. Automatic Segmentation of Liver Tumor in CT Images with Deep Convolutional Neural Networks. *J Comput Commun* 2015; **3**: 146-151 [DOI: [10.4236/jcc.2015.311023](https://doi.org/10.4236/jcc.2015.311023)]
- 17 **Hamm CA,** Wang CJ, Savic LJ, Ferrante M, Schobert I, Schlachter T, Lin M, Duncan JS, Weinreb JC, Chapiro J, Letzen B. Deep learning for liver tumor diagnosis part I: development of a convolutional neural network classifier for multi-phasic MRI. *Eur Radiol* 2019; **29**: 3338-3347 [PMID: [31016442](https://pubmed.ncbi.nlm.nih.gov/31016442/) DOI: [10.1007/s00330-019-06205-9](https://doi.org/10.1007/s00330-019-06205-9)]
- 18 **Jansen MJA,** Kuijff HJ, Veldhuis WB, Wessels FJ, Viergever MA, Pluim JPW. Automatic classification of focal liver lesions based on MRI and risk factors. *PLoS One* 2019; **14**: e0217053 [PMID: [31095624](https://pubmed.ncbi.nlm.nih.gov/31095624/) DOI: [10.1371/journal.pone.0217053](https://doi.org/10.1371/journal.pone.0217053)]
- 19 **Zhang F,** Yang J, Nezami N, Laage-Gaupp F, Chapiro J, De Lin M, Duncan J. Liver Tissue Classification Using an Auto-context-based Deep Neural Network with a Multi-phase Training Framework. *Patch Based Tech Med Imaging (2018)* 2018; **11075**: 59-66 [PMID: [32432233](https://pubmed.ncbi.nlm.nih.gov/32432233/) DOI: [10.1007/978-3-030-00500-9_7](https://doi.org/10.1007/978-3-030-00500-9_7)]
- 20 **Preis O,** Blake MA, Scott JA. Neural network evaluation of PET scans of the liver: a potentially useful adjunct in clinical interpretation. *Radiology* 2011; **258**: 714-721 [PMID: [21339347](https://pubmed.ncbi.nlm.nih.gov/21339347/) DOI: [10.1148/radiol.10100547](https://doi.org/10.1148/radiol.10100547)]
- 21 **Kiani A,** Uyumazturk B, Rajpurkar P, Wang A, Gao R, Jones E, Yu Y, Langlotz CP, Ball RL, Montine TJ, Martin BA, Berry GJ, Ozawa MG, Hazard FK, Brown RA, Chen SB, Wood M, Allard LS, Ylagan L, Ng AY, Shen J. Impact of a deep learning assistant on the histopathologic classification of liver cancer. *NPJ Digit Med* 2020; **3**: 23 [PMID: [32140566](https://pubmed.ncbi.nlm.nih.gov/32140566/) DOI: [10.1038/s41746-020-0232-8](https://doi.org/10.1038/s41746-020-0232-8)]

- 22 **Liao H**, Xiong T, Peng J, Xu L, Liao M, Zhang Z, Wu Z, Yuan K, Zeng Y. Classification and Prognosis Prediction from Histopathological Images of Hepatocellular Carcinoma by a Fully Automated Pipeline Based on Machine Learning. *Ann Surg Oncol* 2020; **27**: 2359-2369 [PMID: [31916093](#) DOI: [10.1245/s10434-019-08190-1](#)]
- 23 **Villanueva A**. Hepatocellular Carcinoma. *N Engl J Med* 2019; **380**: 1450-1462 [PMID: [30970190](#) DOI: [10.1056/NEJMra1713263](#)]
- 24 **Gillies RJ**, Kinahan PE, Hricak H. Radiomics: Images Are More than Pictures, They Are Data. *Radiology* 2016; **278**: 563-577 [PMID: [26579733](#) DOI: [10.1148/radiol.2015151169](#)]
- 25 **Lambin P**, Leijenaar RTH, Deist TM, Peerlings J, de Jong EEC, van Timmeren J, Sanduleanu S, Larue RTHM, Even AJG, Jochems A, van Wijk Y, Woodruff H, van Soest J, Lustberg T, Roelofs E, van Elmpt W, Dekker A, Mottaghy FM, Wildberger JE, Walsh S. Radiomics: the bridge between medical imaging and personalized medicine. *Nat Rev Clin Oncol* 2017; **14**: 749-762 [PMID: [28975929](#) DOI: [10.1038/nrclinonc.2017.141](#)]
- 26 **Erstad DJ**, Tanabe KK. Prognostic and Therapeutic Implications of Microvascular Invasion in Hepatocellular Carcinoma. *Ann Surg Oncol* 2019; **26**: 1474-1493 [PMID: [30788629](#) DOI: [10.1245/s10434-019-07227-9](#)]
- 27 **Ma X**, Wei J, Gu D, Zhu Y, Feng B, Liang M, Wang S, Zhao X, Tian J. Preoperative radiomics nomogram for microvascular invasion prediction in hepatocellular carcinoma using contrast-enhanced CT. *Eur Radiol* 2019; **29**: 3595-3605 [PMID: [30770969](#) DOI: [10.1007/s00330-018-5985-y](#)]
- 28 **Xu X**, Zhang HL, Liu QP, Sun SW, Zhang J, Zhu FP, Yang G, Yan X, Zhang YD, Liu XS. Radiomic analysis of contrast-enhanced CT predicts microvascular invasion and outcome in hepatocellular carcinoma. *J Hepatol* 2019; **70**: 1133-1144 [PMID: [30876945](#) DOI: [10.1016/j.jhep.2019.02.023](#)]
- 29 **Zhou W**, Zhang L, Wang K, Chen S, Wang G, Liu Z, Liang C. Malignancy characterization of hepatocellular carcinomas based on texture analysis of contrast-enhanced MR images. *J Magn Reson Imaging* 2017; **45**: 1476-1484 [PMID: [27626270](#) DOI: [10.1002/jmri.25454](#)]
- 30 **Dong Y**, Zhou L, Xia W, Zhao XY, Zhang Q, Jian JM, Gao X, Wang WP. Preoperative Prediction of Microvascular Invasion in Hepatocellular Carcinoma: Initial Application of a Radiomic Algorithm Based on Grayscale Ultrasound Images. *Front Oncol* 2020; **10**: 353 [PMID: [32266138](#) DOI: [10.3389/fonc.2020.00353](#)]
- 31 **Ji GW**, Zhu FP, Xu Q, Wang K, Wu MY, Tang WW, Li XC, Wang XH. Machine-learning analysis of contrast-enhanced CT radiomics predicts recurrence of hepatocellular carcinoma after resection: A multi-institutional study. *EBioMedicine* 2019; **50**: 156-165 [PMID: [31735556](#) DOI: [10.1016/j.ebiom.2019.10.057](#)]
- 32 **Ho WH**, Lee KT, Chen HY, Ho TW, Chiu HC. Disease-free survival after hepatic resection in hepatocellular carcinoma patients: a prediction approach using artificial neural network. *PLoS One* 2012; **7**: e29179 [PMID: [22235270](#) DOI: [10.1371/journal.pone.0029179](#)]
- 33 **Chiu HC**, Ho TW, Lee KT, Chen HY, Ho WH. Mortality predicted accuracy for hepatocellular carcinoma patients with hepatic resection using artificial neural network. *ScientificWorldJournal* 2013; **2013**: 201976 [PMID: [23737707](#) DOI: [10.1155/2013/201976](#)]
- 34 **Shi HY**, Lee KT, Lee HH, Ho WH, Sun DP, Wang JJ, Chiu CC. Comparison of artificial neural network and logistic regression models for predicting in-hospital mortality after primary liver cancer surgery. *PLoS One* 2012; **7**: e35781 [PMID: [22563399](#) DOI: [10.1371/journal.pone.0035781](#)]
- 35 **Saillard C**, Schmauch B, Laifa O, Moarii M, Toldo S, Zaslavskiy M, Pronier E, Laurent A, Amaddeo G, Regnault H, Sommacale D, Ziolo M, Pawlotsky JM, Mulé S, Luciani A, Wainrib G, Clozet T, Courtiol P, Calderaro J. Predicting survival after hepatocellular carcinoma resection using deep-learning on histological slides. *Hepatology* 2020 [PMID: [32108950](#) DOI: [10.1002/hep.31207](#)]
- 36 **Schoenberg MB**, Bucher JN, Koch D, Börner N, Hesse S, De Toni EN, Seidensticker M, Angele MK, Klein C, Bazhin AV, Werner J, Guba MO. A novel machine learning algorithm to predict disease free survival after resection of hepatocellular carcinoma. *Ann Transl Med* 2020; **8**: 434 [PMID: [32395478](#) DOI: [10.21037/atm.2020.04.16](#)]
- 37 **Forner A**, Reig M, Bruix J. Hepatocellular carcinoma. *Lancet* 2018; **391**: 1301-1314 [PMID: [29307467](#) DOI: [10.1016/S0140-6736\(18\)30010-2](#)]
- 38 **Morshid A**, Elsayer KM, Khalaf AM, Elmohr MM, Yu J, Kaseb AO, Hassan M, Mahvash A, Wang Z, Hazle JD, Fuentes D. A machine learning model to predict hepatocellular carcinoma response to transcatheter arterial chemoembolization. *Radiol Artif Intell* 2019; **1**: e180021 [PMID: [31858078](#) DOI: [10.1148/ryai.2019180021](#)]
- 39 **Peng J**, Kang S, Ning Z, Deng H, Shen J, Xu Y, Zhang J, Zhao W, Li X, Gong W, Huang J, Liu L. Residual convolutional neural network for predicting response of transarterial chemoembolization in hepatocellular carcinoma from CT imaging. *Eur Radiol* 2020; **30**: 413-424 [PMID: [31332558](#) DOI: [10.1007/s00330-019-06318-1](#)]
- 40 **Liu D**, Liu F, Xie X, Su L, Liu M, Xie X, Kuang M, Huang G, Wang Y, Zhou H, Wang K, Lin M, Tian J. Accurate prediction of responses to transarterial chemoembolization for patients with hepatocellular carcinoma by using artificial intelligence in contrast-enhanced ultrasound. *Eur Radiol* 2020; **30**: 2365-2376 [PMID: [31900703](#) DOI: [10.1007/s00330-019-06553-6](#)]
- 41 **Abajian A**, Murali N, Savic LJ, Laage-Gaupp FM, Nezami N, Duncan JS, Schlachter T, Lin M, Geschwind JF, Chapiro J. Predicting Treatment Response to Image-Guided Therapies Using Machine Learning: An Example for Trans-Arterial Treatment of Hepatocellular Carcinoma. *J Vis Exp* 2018; **140**: e58382 [PMID: [30371657](#) DOI: [10.3791/58382](#)]
- 42 **Mähringer-Kunz A**, Wagner F, Hahn F, Weinmann A, Brodehl S, Schotten S, Hinrichs JB, Düber C, Galle PR, Pinto Dos Santos D, Kloockner R. Predicting survival after transarterial chemoembolization for hepatocellular carcinoma using a neural network: A Pilot Study. *Liver Int* 2020; **40**: 694-703 [PMID: [31943703](#) DOI: [10.1111/liv.14380](#)]
- 43 **Sieghart W**, Hücke F, Pinter M, Graziadei I, Vogel W, Müller C, Heinzl H, Trauner M, Peck-Radosavljevic M. The ART of decision making: retreatment with transarterial chemoembolization in patients with hepatocellular carcinoma. *Hepatology* 2013; **57**: 2261-2273 [PMID: [23316013](#) DOI: [10.1002/hep.26256](#)]
- 44 **Adhoue X**, Penaranda G, Naude S, Raoul JL, Perrier H, Bayle O, Monnet O, Beaurain P, Bazin C, Pol B,

- Folgoc GL, Castellani P, Bronowicki JP, Bourlière M. Retreatment with TACE: the ABCR SCORE, an aid to the decision-making process. *J Hepatol* 2015; **62**: 855-862 [PMID: 25463541 DOI: 10.1016/j.jhep.2014.11.014]
- 45 **Kim BK**, Shim JH, Kim SU, Park JY, Kim DY, Ahn SH, Kim KM, Lim YS, Han KH, Lee HC. Risk prediction for patients with hepatocellular carcinoma undergoing chemoembolization: development of a prediction model. *Liver Int* 2016; **36**: 92-99 [PMID: 25950442 DOI: 10.1111/liv.12865]
- 46 **Ziv E**, Yarmohammadi H, Boas FE, Petre EN, Brown KT, Solomon SB, Solit D, Reidy D, Erinjeri JP. Gene Signature Associated with Upregulation of the Wnt/ β -Catenin Signaling Pathway Predicts Tumor Response to Transarterial Embolization. *J Vasc Interv Radiol* 2017; **28**: 349-355.e1 [PMID: 28126478 DOI: 10.1016/j.jvir.2016.11.004]
- 47 **Liang JD**, Ping XO, Tseng YJ, Huang GT, Lai F, Yang PM. Recurrence predictive models for patients with hepatocellular carcinoma after radiofrequency ablation using support vector machines with feature selection methods. *Comput Methods Programs Biomed* 2014; **117**: 425-434 [PMID: 25278224 DOI: 10.1016/j.cmpb.2014.09.001]
- 48 **Xu RH**, Wei W, Krawczyk M, Wang W, Luo H, Flagg K, Yi S, Shi W, Quan Q, Li K, Zheng L, Zhang H, Caughey BA, Zhao Q, Hou J, Zhang R, Xu Y, Cai H, Li G, Hou R, Zhong Z, Lin D, Fu X, Zhu J, Duan Y, Yu M, Ying B, Zhang W, Wang J, Zhang E, Zhang C, Li O, Guo R, Carter H, Zhu JK, Hao X, Zhang K. Circulating tumour DNA methylation markers for diagnosis and prognosis of hepatocellular carcinoma. *Nat Mater* 2017; **16**: 1155-1161 [PMID: 29035356 DOI: 10.1038/nmat4997]
- 49 **Villanueva A**, Portela A, Sayols S, Battiston C, Hoshida Y, Méndez-González J, Imbeaud S, Letouzé E, Hernandez-Gea V, Cornella H, Pinyol R, Solé M, Fuster J, Zucman-Rossi J, Mazzaferro V, Esteller M, Llovet JM; HEPTROMIC Consortium. DNA methylation-based prognosis and epidrivers in hepatocellular carcinoma. *Hepatology* 2015; **61**: 1945-1956 [PMID: 25645722 DOI: 10.1002/hep.27732]
- 50 **Yeh CC**, Goyal A, Shen J, Wu HC, Strauss JA, Wang Q, Gurvich I, Safyan RA, Manji GA, Gamble MV, Siegel AB, Santella RM. Global Level of Plasma DNA Methylation is Associated with Overall Survival in Patients with Hepatocellular Carcinoma. *Ann Surg Oncol* 2017; **24**: 3788-3795 [PMID: 28593503 DOI: 10.1245/s10434-017-5913-4]
- 51 **Dong RZ**, Yang X, Zhang XY, Gao PT, Ke AW, Sun HC, Zhou J, Fan J, Cai JB, Shi GM. Predicting overall survival of patients with hepatocellular carcinoma using a three-category method based on DNA methylation and machine learning. *J Cell Mol Med* 2019; **23**: 3369-3374 [PMID: 30784182 DOI: 10.1111/jcmm.14231]
- 52 **Azer SA**. Deep learning with convolutional neural networks for identification of liver masses and hepatocellular carcinoma: A systematic review. *World J Gastrointest Oncol* 2019; **11**: 1218-1230 [PMID: 31908726 DOI: 10.4251/wjgo.v11.i12.1218]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA
Telephone: +1-925-3991568
E-mail: bpgoffice@wjgnet.com
Help Desk: <https://www.f6publishing.com/helpdesk>
<https://www.wjgnet.com>

