

1

**Name of Journal:** *World Journal of Hepatology*

**ESPS Manuscript NO:** 29360

**Manuscript Type:** Original Article

*Case Control Study*

**Predictors for Advanced Fibrosis in Morbidly Obese non-alcoholic  
Fatty Liver Patients**

**Zelber-Sagi Shira**<sup>1,2</sup> PhD, **Shoham Dafna**<sup>2</sup> MPH, **Zvibel Isabel**<sup>3,5</sup> PhD,  
**Abu-Abeid Subhi**<sup>4</sup> MD, **Shibolet Oren**<sup>1,5</sup> MD Prof. and **Fishman  
Sigal**<sup>3, 5</sup> MD.

## Match Overview

1	<b>Crossref</b> 108 words Shira Zelber-Sagi. "Sampling variability of transient elastography according to probe location :", European Journal of Gastroenterology and Hepatology	3%
2	<b>Internet</b> 48 words crawled on 29-Jun-2016 <a href="http://www.wjgnet.com">www.wjgnet.com</a>	1%
3	<b>Crossref</b> 42 words Zelber-Sagi, S. "Predictors for incidence and remission of NAFLD in the general population during a seven-year follow-up"	1%
4	<b>Crossref</b> 41 words Boursier, Jérôme, Julien Vergniol, Anne Guillet, Jean-Baptiste Hiriart, Adrien Lannes, Brigitte Le Bail, Sophie Micard	1%
5	<b>Internet</b> 30 words crawled on 15-Oct-2015 <a href="http://www.ncbi.nlm.nih.gov">www.ncbi.nlm.nih.gov</a>	1%
6	<b>Crossref</b> 30 words Croci, L. N. M. Byrne, S. Choquette, A. P. Hills, V. S. Chachay, A. D. Clouston, T. M. O'Moore-Sullivan, G. A. Macdonald	1%

学术搜索

找到约 3,280 条结果 (用时0.16秒)

文章

我的图书馆

时间不限

2016以来

2015以来

2012以来

自定义范围...

按相关性排序

按日期排序

搜索所有网页

中文网页

简体中文网页

☒ 包括专利

☐ 包含引用

创建快讯

小提示: 只搜索中文(简体)结果, 可在 学术搜索设置 指定搜索语言

### Independent predictors of liver fibrosis in patients with nonalcoholic steatohepatitis

P Angulo, JC Keach, KP Batts, KD Lindor - Hepatology, 1999 - Wiley Online Library

... size, no factors that determine increasing liver fibrosis and histologically advanced disease have been ... 4) and transferrin saturation (P .4) were not selected as independent predictors in multivariate ... however, only BMI (P .003) was selected as an independent predictor of the ...

被引用次数: 1559 相关文章 所有 8 个版本 引用 保存

### Nonalcoholic fatty liver disease: predictors of nonalcoholic steatohepatitis and liver fibrosis in the severely obese

JB Dixon, PS Bhathal, PE O'brien - Gastroenterology, 2001 - Elsevier

... The predictors for the 26 patients with NASH are shown in Table 3. Using logistic regression, waist-to-hip ratio was the most significant predictor from the ... Raised fasting plasma glucose and fasting insulin levels had independent predictive value for NASH, with the IR ...

被引用次数: 1289 相关文章 所有 8 个版本 引用 保存

### [HTML] Systematic review: the epidemiology and natural history of non - alcoholic fatty liver disease and non - alcoholic steatohepatitis in adults

G Vernon, A Baranova... - Alimentary pharmacology ..., 2011 - Wiley Online Library

... of follow-up, patients with steatosis developed both inflammation and ballooning, and mild fibrotic changes. ... During a median follow-up of 44 months in NAFLD patients with advanced fibrosis (F3 and F4 ... The metabolic syndrome as a predictor of non-alcoholic fatty liver disease. ...

被引用次数: 1170 相关文章 所有 7 个版本 引用 保存

### Predictors of nonalcoholic steatohepatitis and advanced fibrosis in morbidly obese patients

JP Ong, H Elariny, R Collantes, A Younoszai... - Obesity surgery, 2005 - Springer

alanine aminotransferase, insulin resistance, and hypertension were found to be predictive



学术搜索

找到约 3,330 条结果 (用时0.14秒)

文章

我的图书馆

时间不限

2016以来

2015以来

2012以来

自定义范围...

按相关性排序

按日期排序

搜索所有网页

中文网页

简体中文网页

☒ 包括专利

☐ 包含引用

创建快讯

小提示: 只搜索中文(简体)结果, 可在 [学术搜索设置](#) 指定搜索语言

### Nonalcoholic fatty liver disease: predictors of nonalcoholic steatohepatitis and liver fibrosis in the severely obese

[JB Dixon](#), [PS Bhathal](#), [PE O'Brien](#) - [Gastroenterology](#), 2001 - Elsevier

... The predictors for the 26 patients with NASH are shown in Table 3. Using logistic regression, waist-to-hip ratio was the most significant predictor from the ... Raised fasting plasma glucose and fasting insulin levels had independent predictive value for NASH, with the IR ...

被引用次数: 1304 [相关文章](#) [所有 8 个版本](#) [引用](#) [保存](#)

### [HTML] Systematic review: the epidemiology and natural history of non - alcoholic fatty liver disease and non - alcoholic steatohepatitis in adults

[G Vernon](#), [A Baranova](#)... - [Alimentary pharmacology](#) ..., 2011 - Wiley Online Library

... of follow-up, patients with steatosis developed both inflammation and ballooning, and mild fibrotic changes. ... During a median follow-up of 44 months in NAFLD patients with advanced fibrosis (F3 and F4 ... The metabolic syndrome as a predictor of non-alcoholic fatty liver disease. ...

被引用次数: 1253 [相关文章](#) [所有 7 个版本](#) [引用](#) [保存](#)

### Independent predictors of liver fibrosis in patients with nonalcoholic steatohepatitis

[P Angulo](#), [JC Keach](#), [KP Batts](#), [KD Lindor](#) - [Hepatology](#), 1999 - Wiley Online Library

... of Association Between Severe Liver Fibrosis (stages 3-4) and Independent Predictors ... with increased values of transferrin saturation correlated positively with more advanced liver fibrosis in univariate ... trans- ferrin saturation was not selected as an independent predictor of more ...

被引用次数: 1584 [相关文章](#) [所有 8 个版本](#) [引用](#) [保存](#)

### Predictors of nonalcoholic steatohepatitis and advanced fibrosis in morbidly obese patients

[JP Ong](#), [H Elariny](#), [R Collantes](#), [A Younoszai](#)... - [Obesity surgery](#), 2005 - Springer

[全部](#)[新闻](#)[图片](#)[视频](#)[更多 ▾](#)[搜索工具](#)

找到约 85,800 条结果（用时 0.74 秒）

## Google 学术: Predictors for advanced fibrosis in morbidly obese non-alcoholic fatty liver patients

Nonalcoholic fatty liver disease: predictors of ... - Dixon - 被引用次数: 1308

... of non-alcoholic fatty liver disease and non-alcoholic ... - Vernon - 被引用次数: 1265

Independent predictors of liver fibrosis in patients with ... - Angulo - 被引用次数: 1591

## Prevalence and Predictors of Non-Alcoholic Fatty Liver Disease in ...

<https://www.ncbi.nlm.nih.gov/pubmed/25835982> - 翻译此页

作者: P Praveenraj - 2015 - 被引用次数: 7 - 相关文章

Prevalence and Predictors of Non-Alcoholic Fatty Liver Disease in Morbidly Obese ... METHODS: There were 134 morbidly obese patients who underwent ... non-alcoholic steatohepatitis (NASH), fibrosis, and advanced fibrosis were identified.

## Predictors of nonalcoholic steatohepatitis and advanced fibrosis in ...

<https://www.ncbi.nlm.nih.gov/pubmed/15826462> - 翻译此页

作者: JP Ong - 2005 - 被引用次数: 253 - 相关文章

BACKGROUND: Nonalcoholic fatty liver disease (NAFLD) is a common form of ... well as predictors of NASH and advanced fibrosis in morbidly obese patients.

## Predictors of Non-alcoholic Fatty Liver Disease in Obese and ... - NCBI

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - 翻译此页

作者: HM El-Karakasy - 2011 - 被引用次数: 9 - 相关文章

Pediatric non-alcoholic fatty liver disease (NAFLD) is a global problem which ... fibrosis to advanced fibrosis and cirrhosis when fat may no longer be present. .... [13] Ong et al[14] found that 93% of the 212 morbidly obese patients had NAFLD