

World Journal of *Clinical Cases*

World J Clin Cases 2024 January 16; 12(2): 236-465



EDITORIAL

- 236 Use of artificial intelligence in the field of pain medicine
Chang MC

ORIGINAL ARTICLE**Retrospective Study**

- 240 Ultrasound blood flow characteristics changes in fetal umbilical artery thrombosis: A retrospective analysis
Hong SJ, Hong LW, He XQ, Zhong XH
- 249 Electroencephalogram findings in 10 patients with post-stroke epilepsy: A retrospective study
Wen LM, Li R, Wang YL, Kong QX, Xia M
- 256 Exploration of cardiac rehabilitation nursing for elderly patients with myocardial infarction based on individualized cardiac rehabilitation
Liu HN, Gao B
- 267 Survival benefit of concurrent chemoradiotherapy for advanced ampulla of Vater cancer
Kwon CH, Seo HI, Kim DU, Han SY, Kim S, Lee NK, Hong SB, Ahn JH, Park YM, Noh BG
- 276 Utility of plasma D-dimer for diagnosis of venous thromboembolism after hepatectomy
Miyake T, Yanagimoto H, Tsugawa D, Akita M, Asakura R, Arai K, Yoshida T, So S, Ishida J, Urade T, Nanno Y, Fukushima K, Gon H, Komatsu S, Asari S, Toyama H, Kido M, Ajiki T, Fukumoto T
- 285 Lenvatinib combined with sintilimab plus transarterial chemoembolization as first-line treatment for advanced hepatocellular carcinoma
Sun SS, Guo XD, Li WD, Chen JL

Observational Study

- 293 Timing theory integrated nursing combined behavior change integrated theory of nursing on primiparous influence
He YX, Lv Y, Lan TT, Deng F, Zhang YY
- 302 Inverse relationship between platelet Akt activity and hippocampal atrophy: A pilot case-control study in patients with diabetes mellitus
Tokuda H, Hori T, Mizutani D, Hioki T, Kojima K, Onuma T, Enomoto Y, Doi T, Matsushima-Nishiwaki R, Ogura S, Iida H, Iwama T, Sakurai T, Kozawa O

Randomized Controlled Trial

- 314 Impact of continuous care on cardiac function in patients with lung cancer complicated by coronary heart disease
Gao T, Luo JL, Guo P, Hu XW, Wei XY, Hu Y

- 322 Use of cognitive-behavioral career coaching to reduce work anxiety and depression in public employees
Otu MS, Sefotho MM

META-ANALYSIS

- 335 Efficacy and safety of Yangxue Qingnao Granules in treatment of migraine: A systematic review and meta-analysis
Zhou B, Wang GS, Yao YN, Hao T, Li HQ, Cao KG

CASE REPORT

- 346 Use of MLC901 in cerebral venous sinus thrombosis: Three case reports
Arsovska AA, Venketasubramanian N
- 354 Primary biliary cholangitis presenting with granulomatous lung disease misdiagnosed as lung cancer: A case report
Feng SL, Li JY, Dong CL
- 361 Asymptomatic low-grade appendiceal mucinous neoplasm: A case report
Yao MQ, Jiang YP, Wang YY, Mou YP, Fan JX
- 367 Surgically treating a rare and asymptomatic intraductal papillary neoplasm of the bile duct: A case report
Zhu SZ, Gao ZF, Liu XR, Wang XG, Chen F
- 374 Absence of enhancement in a lesion does not preclude primary central nervous system T-cell lymphoma: A case report
Kim CS, Choi CH, Yi KS, Kim Y, Lee J, Woo CG, Jeon YH
- 383 Mental retardation, seizures and language delay caused by new SETD1B mutations: Three case reports
Ding L, Wei LW, Li TS, Chen J
- 392 Three cancers in the renal pelvis, bladder, and colon: A case report
Chen J, Huang HY, Zhou HC, Liu LX, Kong CF, Zhou Q, Fei JM, Zhu YM, Liu H, Tang YC, Zhou CZ
- 399 Severe aconite poisoning successfully treated with veno-arterial extracorporeal membrane oxygenation: A case report
Kohara S, Kamijo Y, Kyan R, Okada I, Hasegawa E, Yamada S, Imai K, Kaizaki-Mitsumoto A, Numazawa S
- 405 Chemotherapy combined with bevacizumab for small cell lung cancer with brain metastases: A case report
Yang HY, Xia YQ, Hou YJ, Xue P, Zhu SJ, Lu DR
- 412 Diagnostic challenges and individualized treatment of cervical adenocarcinoma metastases to the breast: A case report
Akers A, Read S, Feldman J, Gooden C, English DP
- 418 Subsequent bilateral acute carpal tunnel syndrome due to tophaceous infiltration: A case report
Yeoh SC, Wu WT, Shih JT, Su WC, Yeh KT

- 425 Uniportal video-assisted thoracoscopic fissureless right upper lobe anterior segmentectomy for inflammatory myofibroblastic tumor: A case report
Ahn S, Moon Y
- 431 Hybrid treatment of varied orthodontic appliances for a patient with skeletal class II and temporomandibular joint disorders: A case report and review of literature
Lu T, Mei L, Li BC, Huang ZW, Li H
- 443 Significant improvement after sensory tricks and trunk strength training for Parkinson's disease with antecollis and camptocormia: A case report
Wang JR, Hu Y
- 451 Granulomatous mastitis in a 50-year-old male: A case report and review of literature
Cui LY, Sun CP, Li YY, Liu S
- 460 Double-chambered left ventricle with a thrombus in an asymptomatic patient: A case report
Kim N, Yang IH, Hwang HJ, Sohn IS

ABOUT COVER

Editorial Board Member of *World Journal of Clinical Cases*, Xin Ye, MD, Professor, Department of Oncology, The First Affiliated Hospital of Shandong First Medical University, Jinan 250014, Shandong Province, China.
yexintaian2020@163.com

AIMS AND SCOPE

The primary aim of *World Journal of Clinical Cases* (*WJCC*, *World J Clin Cases*) is to provide scholars and readers from various fields of clinical medicine with a platform to publish high-quality clinical research articles and communicate their research findings online.

WJCC mainly publishes articles reporting research results and findings obtained in the field of clinical medicine and covering a wide range of topics, including case control studies, retrospective cohort studies, retrospective studies, clinical trials studies, observational studies, prospective studies, randomized controlled trials, randomized clinical trials, systematic reviews, meta-analysis, and case reports.

INDEXING/ABSTRACTING

The *WJCC* is now abstracted and indexed in Science Citation Index Expanded (SCIE, also known as SciSearch®), Journal Citation Reports/Science Edition, Current Contents®/Clinical Medicine, PubMed, PubMed Central, Reference Citation Analysis, China Science and Technology Journal Database, and Superstar Journals Database. The 2023 Edition of Journal Citation Reports® cites the 2022 impact factor (IF) for *WJCC* as 1.1; IF without journal self cites: 1.1; 5-year IF: 1.3; Journal Citation Indicator: 0.26; Ranking: 133 among 167 journals in medicine, general and internal; and Quartile category: Q4.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: *Hua-Ge Yu*; Production Department Director: *Xu Guo*; Editorial Office Director: *Jin-Lei Wang*.

NAME OF JOURNAL

World Journal of Clinical Cases

ISSN

ISSN 2307-8960 (online)

LAUNCH DATE

April 16, 2013

FREQUENCY

Thrice Monthly

EDITORS-IN-CHIEF

Bao-Gan Peng, Salim Surani, Jerzy Tadeusz Chudek, George Kontogorgos, Maurizio Serati

EDITORIAL BOARD MEMBERS

<https://www.wjgnet.com/2307-8960/editorialboard.htm>

PUBLICATION DATE

January 16, 2024

COPYRIGHT

© 2024 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjgnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>

Retrospective Study

Survival benefit of concurrent chemoradiotherapy for advanced ampulla of Vater cancer

Chae Hwa Kwon, Hyung Il Seo, Dong Uk Kim, Sung Yong Han, Suk Kim, Nam Kyung Lee, Seung Baek Hong, Ji Hyun Ahn, Young Mok Park, Byeong Gwan Noh

Specialty type: Medicine, research and experimental

Provenance and peer review: Unsolicited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's scientific quality classification

Grade A (Excellent): 0
Grade B (Very good): 0
Grade C (Good): C
Grade D (Fair): D
Grade E (Poor): 0

P-Reviewer: Wang K, China; Yang F, China

Received: October 31, 2023

Peer-review started: October 31, 2023

First decision: November 28, 2023

Revised: December 7, 2023

Accepted: December 27, 2023

Article in press: December 27, 2023

Published online: January 16, 2024



Chae Hwa Kwon, BioMedical Research Institute, Pusan National University Hospital, Busan 49241, South Korea

Hyung Il Seo, Young Mok Park, Byeong Gwan Noh, Department of Surgery, Pusan National University School of Medicine and Biomedical Research Institute, Pusan National University Hospital, Busan 49241, South Korea

Dong Uk Kim, Sung Yong Han, Department of Internal Medicine, Pusan National University School of Medicine and Biomedical Research Institute, Pusan National University Hospital, Busan 49241, South Korea

Suk Kim, Nam Kyung Lee, Seung Baek Hong, Department of Radiology, Pusan National University School of Medicine and Biomedical Research Institute, Pusan National University Hospital, Busan 49241, South Korea

Ji Hyun Ahn, Department of Pathology and Biomedical Research Institute, Pusan National University Hospital, Busan 49241, South Korea

Corresponding author: Hyung Il Seo, MD, PhD, Surgeon, Department of Surgery, Pusan National University School of Medicine and Biomedical Research Institute, Pusan National University Hospital, 179, Gudeok-ro, Seo-gu, Busan 49241, South Korea. seohi71@pusan.ac.kr

Abstract

BACKGROUND

Currently, there is no standard adjuvant therapy for patients with resected ampulla of Vater (AoV) cancer.

AIM

To evaluate the effectiveness of adjuvant concurrent chemoradiotherapy (CCRT) in patients with advanced AoV cancer who underwent curative resection.

METHODS

This single-centered, retrospective study included 29 patients with advanced AoV cancer who underwent pancreaticoduodenectomy between 2006 and 2018. The impact of CCRT on advanced AoV cancer was analyzed.

RESULTS

The 1-, 3-, and 5-yr recurrence-free survival (RFS) rates for patients with advanced AoV cancer were 82.8%, 48.3%, and 40.8%, respectively, and the overall survival (OS) rates were 89.7%, 62.1%, and 51.7%, respectively. Lymphovascular invasion was found to be a significant risk factor for RFS and OS in patients with advanced AoV cancer in the univariate analysis, whereas T stage and lymph node metastasis were significantly associated with OS in the multivariate analysis. Compared to the patients who did not receive adjuvant CCRT, those who received adjuvant CCRT did not show statistically significant improvements in the RFS and OS, although they had a significantly lower average age and significantly higher platelet-to-lymphocyte ratio.

CONCLUSION

Adjuvant CCRT did not improve survival outcomes in patients with advanced AoV cancer. These findings contribute to existing knowledge on the effectiveness of CCRT in this patient population and provide important insights for clinical decision-making.

Key Words: Advanced ampulla of Vater cancer; Adjuvant concurrent chemoradiotherapy; Recurrence; Survival; Vater cancer

©The Author(s) 2024. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: We examined the potential survival benefits of adjuvant concurrent chemoradiotherapy (CCRT). Our findings indicated that adjuvant CCRT may not provide any survival advantage to patients with ampulla of Vater (AoV) cancer who had T3/T4 or lymph node-positive tumors. Therefore, the use of adjuvant CCRT as a standard approach in the treatment of advanced AoV cancer patients should be reconsidered.

Citation: Kwon CH, Seo HI, Kim DU, Han SY, Kim S, Lee NK, Hong SB, Ahn JH, Park YM, Noh BG. Survival benefit of concurrent chemoradiotherapy for advanced ampulla of Vater cancer. *World J Clin Cases* 2024; 12(2): 267-275

URL: <https://www.wjgnet.com/2307-8960/full/v12/i2/267.htm>

DOI: <https://dx.doi.org/10.12998/wjcc.v12.i2.267>

INTRODUCTION

Although ampulla of Vater (AoV) cancer has a relatively good prognosis among periampullary cancers, it still has a 5-yr survival rate of only 37%-51% after resection[1,2]. Advanced stage and lymph node involvement were identified as significant prognostic factors[3,4]. Thus, adjuvant treatment is required for patients at advanced stages with a high risk of recurrence. However, previous studies have suggested that radiotherapy alone has no survival benefit[5,6]. Further, Miura *et al*[6] reported minor benefits of radiotherapy in patients with lymph node (LN) metastases. Consequently, the most commonly used adjuvant treatment is chemotherapy or concurrent chemoradiotherapy (CCRT).

Several trials have demonstrated that adjuvant chemotherapy is associated with improved survival compared with observation alone, with a benefit greater in patients with advanced T3/T4 lesions or LN metastases[4,7,8]. However, some studies have reported contrasting opinions[9,10]. Despite the controversy surrounding the benefits of adjuvant chemotherapy after resection for AoV cancer and the lack of consensus guidelines, adjuvant chemotherapy tends to be used clinically[11]. Currently, the combination of adjuvant chemotherapy and radiation is the standard approach. However, evidence supporting the use of CCRT in AoV cancer is limited, and further research is needed to determine the best treatment approach. Therefore, this study aimed to retrospectively evaluate the efficacy of adjuvant CCRT in patients with advanced AoV carcinoma who underwent curative-intent resection.

MATERIALS AND METHODS

Patients

A total of 92 patients with AoV cancer, confirmed by histological examination, underwent pancreaticoduodenectomy between January 2006 and December 2018 at a single tertiary hospital by a single surgeon. Of these, 11 patients with advanced AoV cancer (T3/T4 or LN metastases) who underwent adjuvant CCRT, and 18 patients who did not receive adjuvant therapy and who did not meet the following exclusion criteria were included: exclusive adjuvant chemotherapy or radiotherapy, treatment-associated mortality, significant decline in post-surgery physical strength, variants of adenocarcinoma, neuroendocrine carcinoma, regional LN metastasis beyond LN, total pancreatectomy, incomplete medical records, and cause of death not recurrence but a different medical cause. The clinicopathologic characteristics of the patients are showed in [Supplementary Table 1](#). TNM staging was performed according to the 8th edition of the American Joint Committee Staging System. Demographic, adjuvant treatment, and survival outcome data were retrospectively obtained from the medical records. This study was approved by the Institutional Review Board of Clinical Trial Center in our hospital (IRB No. 2303-007-124).




Figure 1 Kaplan-Meier curves of 29 AoV cancer patients. A: Recurrence-free survival; B: Overall survival. The dashed line indicates the median survival time.

Adjuvant CCRT

Adjuvant CCRT was recommended for patients with LN metastasis regardless of the T stage and was administered 4-8 wk after surgery according to the patient's voluntary decision. The chemotherapy regimen consisted of six cycles of fluoropyridine- or gemcitabine-based chemotherapy [5-FU plus leucovorin (FLv) or gemcitabine plus cisplatin (GP)] for 6 cycles, with a radiation dose of 50.4 Gy. One patient died of gastric infarction and was excluded from the study.

Follow-up

Patients underwent tumor marker (CA19-9 and CEA) testing every 3 mo, with additional imaging tests (abdominal and chest computed tomography) performed if abnormalities were detected. If there were no abnormalities, imaging tests were performed every 6 mo. Palliative treatment was administered to patients who experienced recurrence. One patient died from complications due to radiofrequency ablation and was excluded from the study.

Statistical analysis

The clinical and pathological characteristics of the CCRT and non-CCRT groups were compared and analyzed using the Mann-Whitney *U* and chi-square tests for continuous and categorical variables, respectively. Univariate and multivariate analyses were performed using Cox proportional hazard models. Statistical analyses were performed using the R software (version 4.2.1). The R packages "moonbook," "survminer" and "survival," were used. Statistical significance was set at $P < 0.05$. The statistical methods of this study were reviewed by Kim JM from Biomedical Research Institute, Pusan National University Hospital.

RESULTS

The 1-, 3-, and 5-yr recurrence-free survival (RFS) rates for advanced AoV cancer were 82.8%, 48.3%, and 40.8%, respectively, and the overall survival (OS) rates were 89.7%, 62.1%, and 51.7%, respectively. The median RFS and OS durations were 34 and 64 mo, respectively (Figure 1). Tables 1 and 2 show the results of the univariate and multivariate analyses for RFS and OS in patients with advanced AoV cancer. Patients with positive lymphovascular invasion (LVI) had a significantly higher risk of RFS [hazard ratio (HR): 2.971, confidence interval (CI): 1.123-7.861, $P = 0.028$] and OS (HR: 3.35, CI: 1.226-9.153, $P = 0.018$, respectively). However, other factors such as age; tumor size; T stage; LN metastasis; perineural invasion (PNI); differentiation, and CCRT; and biochemical markers such as, bilirubin, CEA, CA19.9, CRP, albumin, platelet-to-lymphocyte ratio (PLR), and neutrophil-to-lymphocyte ratio (NLR), were not significantly associated with RFS or OS in the univariate analysis. In the multivariate analysis, while LVI did not have a statistically significant impact on RFS or OS, T stage and LN metastasis were significant prognostic factors for OS (HR: 3.015, CI: 0.989-9.095, $P = 0.052$; HR: 3.702, CI: 1.116-12.283, $P = 0.032$, respectively).

Patients with advanced AoV cancer were divided into two groups according to whether they received CCRT: patients who received adjuvant CCRT (CCRT, $n = 11$) and those who did not (non-CCRT, $n = 18$). The clinical features were then compared between these two groups. As shown in Figure 2A, the 3-yr and 5-yr RFS rates were 55.6% and 50.0%, respectively, in the non-CCRT group, and 36.4% and 27.3%, respectively, in the CCRT group. The median RFS in the non-CCRT and CCRT groups was 43 and 32 mo, respectively. The 3-yr and 5-yr OS rates were 55.6% and 55.6% in the non-

Table 1 Univariate and multivariate analyses for recurrence-free survival

	Univariate		Multivariate	
	HR (95%CI)	P value	HR (95%CI)	P value
Sex (male)	1.988 (0.568-6.955)	0.282		
Age	0.976 (0.939-1.014)	0.215		
Size	0.816 (0.530-1.255)	0.354		
T stage	1.808 (0.485-6.743)	0.378	1.952 (0.499-7.628)	0.326
LN metastasis (positive)	2.026 (0.747-5.498)	0.166	1.848 (0.522-6.541)	0.341
LVI (positive)	2.971 (1.123-7.861)	0.028 ^a	2.002 (0.589-6.806)	0.266
PNI (positive)	1.026 (0.395-2.663)	0.958		
Differentiation (poor)	1.366 (0.504-3.706)	0.540		
CCRT	1.543 (0.594-4.006)	0.373		
Bilirubin	0.935 (0.864-1.01)	0.089		
CEA (≥ 5 ng/mL)	0.429 (0.056-3.288)	0.416		
CA19.9 (≥ 39 U/L)	0.635 (0.225-1.796)	0.392		
CRP (≥ 0.5 mg/mL)	1.139 (0.367-3.536)	0.822		
Albumin (< 3.3 or > 5.2 mg/dL)	0.903 (0.119-6.831)	0.922		
PLR	1.001 (0.997-1.005)	0.661		
NLR	0.999 (0.871-1.144)	0.983		

^aP < 0.05.

LN: Lymph node; LVI: Lymphovascular invasion; PNI: Perineural invasion; CCRT: Concurrent chemoradiotherapy; CEA: Carcinoembryonic antigen; CA19.9: Carbohydrate antigen 19-9; CRP: C-reactive protein; PLR: Platelet-to-lymphocyte ratio; NLR: Neutrophil-to-lymphocyte ratio; HR: Hazard ratio; CI: Confidence interval.




Figure 2 Kaplan-Meier curves between concurrent chemoradiotherapy (n = 11) and non-concurrent chemoradiotherapy groups (n = 18). A: Recurrence-free survival; **B:** Overall survival. CCRT: Concurrent chemoradiotherapy.

Table 2 Univariate and multivariate analyses for overall survival

	Univariate		Multivariate	
	HR (95%CI)	P value	HR (95%CI)	P value
Sex (male)	1.752 (0.5667-5.416)	0.33		
Age	0.992 (0.9545-1.031)	0.693		
Size	0.924 (0.6395-1.334)	0.672		
T stage	2.102 (0.6376-6.927)	0.222	3.015 (0.989-9.159)	0.052
LN metastasis (positive)	2.596 (0.967-6.968)	0.058	3.702 (1.116-12.283)	0.032 ^a
LVI (positive)	3.35 (1.226-9.153)	0.018 ^a	1.806 (0.549-5.941)	0.33
PNI (positive)	1.344 (0.5331-3.39)	0.531		
Differentiation (poor)	1.398 (0.5406-3.617)	0.489		
CCRT	1.073 (0.423-2.724)	0.881		
Bilirubin	0.966 (0.8975-1.04)	0.359		
CEA (≥ 5 ng/mL)	0.423 (0.0555-3.226)	0.407		
CA19.9 (≥ 39 U/L)	1.372 (0.475-3.963)	0.559		
CRP (≥ 0.5 mg/mL)	3.503 (0.7991-15.35)	0.096		
Albumin (< 3.3 or > 5.2 mg/dL)	3.088 (0.6807-14.01)	0.144		
PLR	1.003 (0.9991-1.007)	0.128		
NLR	1.061 (0.9529-1.18)	0.282		

^a*P* < 0.05.

LN: Lymph node; LVI: Lymphovascular invasion; PNI: Perineural invasion; CCRT: Concurrent chemoradiotherapy; CEA: Carcinoembryonic antigen; CA19-9: Carbohydrate antigen 19-9; CRP: C-reactive protein; PLR: Platelet-to-lymphocyte ratio; NLR: Neutrophil-to-lymphocyte ratio; HR: Hazard ratio; CI: Confidence interval.

CCRT group and 72.7% and 45.5% in the CCRT group, respectively. The median OS durations of the non-CCRT and CCRT groups were 76 and 46 mo, respectively (Figure 2B). The patients who had received CCRT showed worse RFS and OS than those who did not receive CCRT, but the difference was not statistically significant (*P* = 0.37, *P* = 0.89, respectively). Table 3 shows the patient characteristic in the CCRT and non-CCRT groups. The average of age was significantly lower in the CCRT group than in the non-CCRT group (59.0 ± 8.3 vs 71.22 ± 12.17, *P* = 0.007). Additionally, the PLR was significantly higher in the CCRT group (*P* = 0.044). However, there were no significant differences in sex, tumor size, LN metastasis, LVI, PNI, CEA, CA19.9, CRP, albumin, NLR, American Society of Anesthesiologists grade, or transfusion between the two groups.

DISCUSSION

After tumor resection, the prognosis of patients with AoV cancer is generally better than that of patients with pancreatic cancer; however, poor survival rates have been observed in patients with advanced-stage cancer. Adjuvant treatment is often used in these cases, and this study evaluated the effectiveness of adjuvant CCRT in patients with AoV cancer with poor prognostic factors, such as high T stage or LN metastases. However, our findings suggest that adjuvant CCRT may not provide any survival benefits in patients with advanced AoV cancer.

While some studies suggest that adjuvant chemotherapy (ACT) provides a survival benefit for patients with advanced AoV cancer, studies such as the ESPAC-3[7] and Al Abbas *et al*[12] have some limitations, including heterogeneity in the study population and variations in the chemotherapy drugs used. To date, most studies have used 5-FU- or gemcitabine-based chemotherapy as ACT. In this study, FLv or GP regimens were used, but no difference in the survival benefit was observed. Some studies have compared the efficacy of 5-FU- and gemcitabine-based regimens. In a subgroup analysis of the ESPAC-3 study population, the patients who received gemcitabine showed a survival benefit, whereas no benefit was observed among those who received 5-FU with folinic acid[7]. Conversely, in a study by Al Abbas *et al*[12], 5-FU led to better survival rates than gemcitabine in patients with advanced-stage disease. Therefore, an optimal treatment regimen for patients with AoV cancer is yet to be established.

Recent studies have evaluated the different effects of ACT regimens on survival outcomes based on histological subtypes of AoV cancer[9,12,13]. While 5-FU- and gemcitabine-based regimens may be useful for certain histological subtypes, Al Abbas *et al*[12] reported that 5-FU-based ACT improved survival regardless of the subtype. However, Ecker

Table 3 Baseline characteristics between the ampulla of Vater cancer patients with and without concurrent chemoradiotherapy

Characteristics	CCRT (n = 11)	Non-CCRT (n = 18)	Total (n = 29)	P value
Sex (%)				0.647
Male	9 (81.8)	12 (66.7)	21 (72.4)	
Female	2 (18.2)	6 (33.3)	8 (27.6)	
Age, median (IQR)	60 (52; 63)	75.5 (63.3; 79.5)	65 (60; 76)	0.007 ^b
Size, median (IQR)	2.9 (1.9; 3.8)	2.7 (2.2; 3.0)	2.7 (2.1; 3.2)	0.928
T stage (%)				0.200
2	3 (27.3)	2 (11.1)	5 (17.2)	
3	7 (63.6)	16 (88.9)	23 (79.3)	
4	1 (9.1)	0 (0.0)	1 (3.4)	
LN metastasis (%)				1.000
Negative	5 (45.5)	9 (50.0)	14 (48.3)	
Positive	6 (54.5)	9 (50.0)	15 (51.7)	
LVI (%)				1.000
Negative	7 (63.6)	12 (66.7)	19 (65.5)	
Positive	4 (36.4)	6 (33.3)	10 (34.5)	
PNI (%)				0.661
Negative	5 (45.5)	11 (61.1)	16 (55.2)	
Positive	6 (54.5)	7 (38.9)	13 (44.8)	
Differentiation (%)				0.076
Well-to-moderate	4 (36.4)	14 (77.8)	18 (62.1)	
Poor	6 (54.5)	3 (16.7)	9 (31.0)	
Unknown	1 (9.1)	1 (5.6)	2 (6.9)	
Bilirubin	4.6 (1.3; 11.9)	2.8 (0.5; 13.7)	4.1 (0.9; 12.9)	0.515
CEA (%)				1.000
< 5 ng/mL	8 (88.9)	14 (87.5)	22 (88.0)	
≥ 5 ng/mL	1 (11.1)	2 (12.5)	3 (12.0)	
CA19.9 (%)				0.866
< 39 U/L	3 (30.0)	7 (41.2)	10 (37.0)	
≥ 39 U/L	7 (70.0)	10 (58.8)	17 (63.0)	
CRP (%)				0.297
< 0.5 mg/dL	1 (11.1)	7 (38.9)	8 (29.6)	
≥ 0.5 mg/dL	8 (88.9)	11 (61.1)	19 (70.4)	
Albumin (%)				1.000
3.3-5.2 g/dL	10 (90.9)	17 (94.4)	27 (93.1)	
< 3.3 or > 5.2 g/dL	1 (9.1)	1 (5.6)	2 (6.9)	
PLR, median (IQR)	277.0 (215.1; 291.7)	159.8 (124.1; 237.9)	202.0 (133.9; 287.5)	0.044 ^a
NLR, median (IQR)	4.1 (3.3; 4.5)	2.7 (2.2; 4.7)	3.6 (2.3; 4.6)	0.334
ASA				0.105
I	3 (27.3)	0 (0.0)	3 (10.3)	
II	6 (54.5)	11 (61.1)	17 (58.6)	
III	2 (18.2)	6 (33.3)	8 (27.6)	

IV	0 (0.0)	1 (5.6)	1 (3.4)
----	---------	---------	---------

^a $P < 0.05$.

^b $P < 0.01$.

LN: Lymph node; LVI: Lymphovascular invasion; PNI: Perineural invasion; CCRT: Concurrent chemoradiotherapy; CEA: Carcinoembryonic antigen; CA19-9: Carbohydrate antigen 19-9; CRP: C-reactive protein; PLR: Platelet-to-lymphocyte ratio; NLR: Neutrophil-to-lymphocyte ratio; ASA: American Society of Anesthesiologists.

et al[9] found no survival benefit of ACT regardless of the histological type or advanced stage. The lack of an association between chemotherapeutic agents and histological type presents a challenge in evaluating the efficacy of ACT. Although the benefits of different chemotherapy regimens according to histologic type have been suggested, the small number of patients in each subset and poor reproducibility of histologic typing are considerable limitations. In this study, most of the patients had not undergone phenotypic classification when receiving adjuvant therapy; therefore, treatment decisions were not based on histological subtypes.

The effectiveness of CCRT in patients with advanced-stage AoV cancer is controversial, despite some studies demonstrating its benefits. The Mayo Clinic reported improved disease relapse and survival rates following CCRT[4]. In addition, Nassour *et al*[11] showed that CCRT was associated with improved survival compared with observation only. In contrast, studies by Ecker *et al*[9] and Kim *et al*[10,14] found no significant differences in survival or recurrence rates, regardless of the addition of radiotherapy. Therefore, the use of CCRT in clinical practice should be carefully considered, as this study highlights that CCRT does not have survival benefits for patients with advanced AoV cancer.

Our study found that age and PLR were correlated with the administration of adjuvant CCRT, with older patients and those with a low PLR being less likely to receive this treatment. These findings are consistent with those reported by the Mayo Clinic regarding age. However, these factors should not be used as the criteria for selecting patients for CCRT, as our study results did not show a significant difference in the survival outcomes between the CCRT and non-CCRT groups.

This study had limitations. First, this was a single-center, retrospective, non-randomized study with a small sample size, which should be taken into consideration when interpreting the results. The limited sample size is attributed to the specificity of the disease, and particularly in advanced stages, the target group inevitably diminishes. Therefore, further confirmation of the findings is warranted through multi-institutional research. Second, the chemotherapy regimens used were not standardized and the histological type was not evaluated. Third, a large proportion of the patient population received palliative treatment, which may have influenced the OS outcomes.

CONCLUSION

In conclusion, our study provides some evidence that CCRT may not provide a survival benefit to patients with advanced AoV cancer. Therefore, further research is required to address the limitations of the current study and provide definitive answers regarding the role of CCRT in the treatment of advanced AoV cancer.

ARTICLE HIGHLIGHTS

Research background

Although ampulla of Vater (AoV) cancer has a relatively favorable prognosis among periampullary cancers, the 5-yr survival rate after resection remains poor. The benefits of adjuvant treatment for AoV cancer is still controversial, leading to the prevalent use of chemotherapy or concurrent chemoradiotherapy (CCRT).

Research motivation

Despite clinical use of adjuvant treatment, there are no consensus guidelines for patients with AoV carcinoma. This study aims to contribute valuable insights into the survival benefits of CCRT in patients with advanced AoV cancer, providing evidence to guide treatment decisions.

Research objectives

The study aims to retrospectively assess the efficacy of adjuvant CCRT in patients with advanced AoV carcinoma who underwent curative resection.

Research methods

Eleven patients with advanced AoV cancer [T3/T4 or lymph node (LN) metastases] who underwent adjuvant CCRT, and 18 patients who did not receive adjuvant therapy were retrospectively reviewed.

Research results

The study found 1-, 3-, and 5-yr RFS rates of 82.8%, 48.3%, and 40.8%, and OS rates of 89.7%, 62.1%, and 51.7%, respectively, for advanced AoV cancer. T stage and LN metastasis were significantly associated with OS in the multivariate analysis. However, CCRT did not show a statistically significant survival advantage.

Research conclusions

This study suggests that adjuvant CCRT may not provide survival benefits for patients with advanced AoV cancer.

Research perspectives

Additional multi-institutional studies with larger sample sizes, standardized regimens, and histological evaluations are recommended to identify the optimal options for patients with advanced AoV cancer.

FOOTNOTES

Author contributions: Seo HI contributed design of the study; Kwon CH contributed analysis of data; Seo HI and Kwon CH contributed interpretation of data and drafting the article; Kim DU and Han SY contributed review the manuscript; Kim S, Lee NK, Hong SB, Ahn JH, Par YM and Noh BG contributed acquisition of data; all authors have read and approved the final manuscript.

Institutional review board statement: This study was approved by the Institutional Review Board of Clinical Trial Center in Pusan National University hospital (IRB No. 2303-007-124).

Informed consent statement: All study participants, or their legal guardian, provided informed written consent prior to study enrollment.

Conflict-of-interest statement: The authors have no financial relationships to disclose.

Data sharing statement: The data that support the findings of this study are available from the corresponding author.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Country/Territory of origin: South Korea

ORCID number: Chae Hwa Kwon 0000-0003-2277-1570; Hyung Il Seo 0000-0002-4132-7662; Dong Uk Kim 0000-0002-7208-7753; Sung Yong Han 0000-0002-0256-9781; Suk Kim 0000-0003-3268-1763; Nam Kyung Lee 0000-0003-1972-2719; Seung Baek Hong 0000-0002-1731-0430; Ji Hyun Ahn 0000-0002-3312-788X; Young Mok Park 0000-0002-4165-3054; Byeong Gwan Noh 0000-0002-7764-9516.

S-Editor: Gao CC

L-Editor: A

P-Editor: Cai YX

REFERENCES

- 1 **Berberat PO**, Künzli BM, Gulbinas A, Ramanauskas T, Kleeff J, Müller MW, Wagner M, Friess H, Büchler MW. An audit of outcomes of a series of periampullary carcinomas. *Eur J Surg Oncol* 2009; **35**: 187-191 [PMID: 18343082 DOI: 10.1016/j.ejso.2008.01.030]
- 2 **O'Connell JB**, Maggard MA, Manunga J Jr, Tomlinson JS, Reber HA, Ko CY, Hines OJ. Survival after resection of ampullary carcinoma: a national population-based study. *Ann Surg Oncol* 2008; **15**: 1820-1827 [PMID: 18369675 DOI: 10.1245/s10434-008-9886-1]
- 3 **Sakata J**, Shirai Y, Wakai T, Yokoyama N, Sakata E, Akazawa K, Hatakeyama K. Number of positive lymph nodes independently affects long-term survival after resection in patients with ampullary carcinoma. *Eur J Surg Oncol* 2007; **33**: 346-351 [PMID: 17097846 DOI: 10.1016/j.ejso.2006.10.008]
- 4 **Jin Z**, Hartgers ML, Sanhueza CT, Shubert CR, Alberts SR, Truty MJ, Muppa P, Nagorney DM, Smyrk TC, Hassan M, Mahipal A. Prognostic factors and benefits of adjuvant therapy after pancreatoduodenectomy for ampullary adenocarcinoma: Mayo Clinic experience. *Eur J Surg Oncol* 2018; **44**: 677-683 [PMID: 29506768 DOI: 10.1016/j.ejso.2018.02.008]
- 5 **Zhao W**, Wang B, Zhao A, Tian Q, Zhang L, Wang L, Zhao X, Yang J, Dong D. The role of radiotherapy in patients with resected ampullary carcinoma: findings based on the SEER database. *HPB (Oxford)* 2019; **21**: 1535-1540 [PMID: 31056288 DOI: 10.1016/j.hpb.2019.03.369]
- 6 **Miura JT**, Jayakrishnan TT, Amini A, Johnston FM, Tsai S, Erickson B, Quebbeman EJ, Christians KK, Evans DB, Gamblin TC, Turaga KK. Defining the role of adjuvant external beam radiotherapy on resected adenocarcinoma of the ampulla of Vater. *J Gastrointest Surg* 2014; **18**: 2003-2008 [PMID: 25159502 DOI: 10.1007/s11605-014-2629-7]
- 7 **Neoptolemos JP**, Moore MJ, Cox TF, Valle JW, Palmer DH, McDonald AC, Carter R, Tebbutt NC, Dervenis C, Smith D, Glimelius B, Charnley RM, Lacaine F, Scarfe AG, Middleton MR, Anthony A, Ghaneh P, Halloran CM, Lerch MM, Oláh A, Rawcliffe CL, Verbeke CS, Campbell F, Büchler MW; European Study Group for Pancreatic Cancer. Effect of adjuvant chemotherapy with fluorouracil plus folinic acid or gemcitabine vs observation on survival in patients with resected periampullary adenocarcinoma: the ESPAC-3 periampullary cancer

- randomized trial. *JAMA* 2012; **308**: 147-156 [PMID: 22782416 DOI: 10.1001/jama.2012.7352]
- 8 **Kwon J**, Kim BH, Kim K, Chie EK, Ha SW. Survival Benefit of Adjuvant Chemoradiotherapy in Patients With Ampulla of Vater Cancer: A Systematic Review and Meta-analysis. *Ann Surg* 2015; **262**: 47-52 [PMID: 25775067 DOI: 10.1097/SLA.0000000000001182]
- 9 **Ecker BL**, Vollmer CM Jr, Behrman SW, Allegrini V, Aversa J, Ball CG, Barrows CE, Berger AC, Cagigas MN, Christein JD, Dixon E, Fisher WE, Freedman-Weiss M, Guzman-Pruneda F, Hollis RH, House MG, Kent TS, Kowalsky SJ, Malleo G, Salem RR, Salvia R, Schmidt CR, Seykora TF, Zheng R, Zureikat AH, Dickson PV. Role of Adjuvant Multimodality Therapy After Curative-Intent Resection of Ampullary Carcinoma. *JAMA Surg* 2019; **154**: 706-714 [PMID: 31141112 DOI: 10.1001/jamasurg.2019.1170]
- 10 **Kim HS**, Jang JY, Yoon YS, Park SJ, Kwon W, Kim SW, Han HS, Han SS, Park JS, Yoon DS. Does adjuvant treatment improve prognosis after curative resection of ampulla of Vater carcinoma? A multicenter retrospective study. *J Hepatobiliary Pancreat Sci* 2020; **27**: 721-730 [PMID: 32652820 DOI: 10.1002/jhbp.801]
- 11 **Nassour I**, Hynan LS, Christie A, Minter RM, Yopp AC, Choti MA, Mansour JC, Porembka MR, Wang SC. Association of Adjuvant Therapy with Improved Survival in Ampullary Cancer: A National Cohort Study. *J Gastrointest Surg* 2018; **22**: 695-702 [PMID: 29127604 DOI: 10.1007/s11605-017-3624-6]
- 12 **Al Abbas AI**, Falvello V, Zenati M, Mani A, Hogg ME, Zeh HJ 3rd, Singhi A, Bahary N, Zureikat AH. Impact of adjuvant chemotherapy regimen on survival outcomes in immunohistochemical subtypes of ampullary carcinoma. *J Surg Oncol* 2020; **121**: 322-329 [PMID: 31840257 DOI: 10.1002/jso.25808]
- 13 **Chang DK**, Jamieson NB, Johns AL, Scarlett CJ, Pajic M, Chou A, Pinese M, Humphris JL, Jones MD, Toon C, Nagrial AM, Chantrill LA, Chin VT, Pinho AV, Rooman I, Cowley MJ, Wu J, Mead RS, Colvin EK, Samra JS, Corbo V, Bassi C, Falconi M, Lawlor RT, Crippa S, Sperandio N, Bersani S, Dickson EJ, Mohamed MA, Oien KA, Foulis AK, Musgrove EA, Sutherland RL, Kench JG, Carter CR, Gill AJ, Scarpa A, McKay CJ, Biankin AV. Histomolecular phenotypes and outcome in adenocarcinoma of the ampulla of vater. *J Clin Oncol* 2013; **31**: 1348-1356 [PMID: 23439753 DOI: 10.1200/JCO.2012.46.8868]
- 14 **Kim JH**, Jeong JH, Ryoo BY, Kim KP, Chang HM, Oh D, Song TJ, Lee SS, Seo DW, Lee SK, Kim MH, Park Y, Kwon JW, Hwang DW, Lee JH, Lee W, Kim SC, Yoo C, Song KB. Adjuvant Chemotherapy for Resected Ampulla of Vater Carcinoma: Retrospective Analysis of 646 Patients. *Cancer Res Treat* 2021; **53**: 424-435 [PMID: 33171024 DOI: 10.4143/crt.2020.953]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA
Telephone: +1-925-3991568
E-mail: bpgoffice@wjgnet.com
Help Desk: <https://www.f6publishing.com/helpdesk>
<https://www.wjgnet.com>

