

Minimal deviation mucinous adenocarcinoma ('adenoma ...<https://onlinelibrary.wiley.com/doi/10.1111/j.1440-1827.2009.02473.x>

Imaging demonstrated thickened myometrium due to adenomyosis. Serum CA19-9 levels were elevated to 486 U/mL. Microscopic examination of hysterectomy specimens indicated highly differentiated...

Cited by: 17 Author: Kaoru Abiko, Tsukasa Baba, Madoka Og...

Publish Year: 2010

Minimal deviation mucinous adenocarcinoma ('adenoma ...https://www.researchgate.net/publication/40893939_Minimal_deviation_mucinous...

Serum CA19-9 levels were elevated to 486 U/mL. ... overlapping imaging characteristics-case report and literature review. ... case of minimal deviation adenocarcinoma of endometrioid type ...

Minimal deviation mucinous adenocarcinoma of the uterine ...<https://www.spandidos-publications.com/ol/8/6/2481>

Minimal deviation adenocarcinoma (MDA), also known as adenoma malignum of the uterine cervix, accounts for only ~1% of uterine cervical adenocarcinomas. ... Case report. A 65-year-old female...

Minimal deviation mucinous adenocarcinoma ('adenoma ...<https://onlinelibrary.wiley.com/doi/abs/10.1111/j.1440-1827.2009.02473.x>

Dec 28, 2009 · Minimal deviation mucinous adenocarcinoma ('adenoma malignum') of the uterine corpus. ... A case is described of a 56-year-old postmenopausal woman who presented with mucinous...

Cited by: 17 Author: Kaoru Abiko, Tsukasa Baba, Madoka Og...

Publish Year: 2010

Minimal deviation mucinous adenocarcinoma ('adenoma ...<https://www.deepdyve.com/lp/wiley/minimal...>

Jan 01, 2010 · Primary mucinous adenocarcinomas of the uterine corpus are typically low grade and frequently associated with endometrial hyperplasia and/or ordinary endometrioid adenocarcinoma, bu...

A case of CA19-9-producing adenocarcinoma of the remnant ...<https://www.researchgate.net/publication/229550102...>

We report a case of early gastric cancer with a high level of serum CA19-9 (400IU/ml). A 62-year-old

ALL IMAGES VIDEOS

86,000 Results Any time

Minimal deviation mucinous adenocarcinoma ('adenoma ...
https://www.researchgate.net/publication/40893939...
Serum CA19-9 levels were elevated to 486 U/mL... overlapping imaging characteristics-case report and literature review. ... case of minimal deviation adenocarcinoma of endometrioid type ...

Minimal deviation mucinous adenocarcinoma ('adenoma ...
https://www.deepdyve.com/lp/wiley/minimal...
Jan 01, 2010 T2-weighted imaging showed an ill-defined myometrial thickening with decreased intensity, suggesting adenomyosis. Serum CA19-9 levels were 486.2 U/mL (normal, <37 U/mL), while CEA and CA125 levels were within the normal range. Vaginal discharge contained elevated CA19-9, measuring 211 ...

Minimal deviation mucinous adenocarcinoma ('adenoma ...
https://onlinelibrary.wiley.com/doi/abs/10.1111/j.1440-1827.2009.02473.x
Dec 28, 2009 Minimal deviation mucinous adenocarcinoma ('adenoma malignum') of the uterine corpus ... A case is described of a 56-year-old postmenopausal woman who presented with mucinous vaginal discharge. Imaging demonstrated thickened myometrium due to adenomyosis. ... Serum CA19-9 levels were elevated to 486 U/mL. Microscopic examination of ...
Cited by: 17 Author: Kaoru Abiko, Tsukasa Baba, Madoka Ogaw...
Publish Year: 2010

Minimal deviation mucinous adenocarcinoma ('adenoma ...
https://onlinelibrary.wiley.com/doi/10.1111/j.1440-1827.2009.02473.x
Primary mucinous adenocarcinomas of the uterine corpus are typically low grade and frequently associated with endometrial hyperplasia and/or ordinary endometrioid adenocarcinoma, but ...
Cited by: 17 Author: Kaoru Abiko, Tsukasa Baba, Madoka Ogaw...
Publish Year: 2010

Minimal deviation mucinous adenocarcinoma of the uterine ...
https://www.spandidos-publications.com/ol/8/6/2481
Minimal deviation adenocarcinoma (MDA), also known as adenoma malignum of the uterine cervix, accounts for only ~1% of uterine cervical adenocarcinomas. ... Case report. A 65-year-old female (gravida 2, para 0) visited a local clinic due to abnormal vaginal bleeding. ... 125 levels were 153 U/ml (normal, <35 U/ml) and CA19-9 levels were 44 U ...

A case of CA19-9-producing adenocarcinoma of the remnant ...
https://www.researchgate.net/publication/229550102...
We report a case of early gastric cancer with a high level of serum CA19-9 (400IU/ml). A 62-year-old man was admitted to the hospital because of abnormality of the stomach by screening.

PDF Diagnostic challenges in minimal deviation adenocarcinoma ...
https://www.spandidos-publications.com/10.3892/mco.2013.144/download
Minimal deviation adenocarcinoma (MDA) of the uterine cervix, otherwise known as adenoma malignum, is a rare variant of cervical adenocarcinoma, which represents a diagnostic challenge in the field of gynecologic oncology, due to its benign-resembling histological characteristics. To achieve a better understanding of this disease, we present

PDF Ho Journal of Cytology & Histology Kanomata and Muramaki ...
https://www.hilarispublisher.com/open-access/well...
case of nested variant of urothelial carcinoma with minimal deviation adenocarcinoma-like features [3]. Herein, we report a pure primary adenocarcinoma of the urinary bladder with minimal deviation adenocarcinoma-like features. Case Report A 64-year-old Japanese woman presented complaining of hematuria and lower abdominal pain.

Elevated CA 125 level in a mucinous cystadenoma and a ...
https://medicalcasereports.biomedcentral.com/...
Sep 03, 2020 The presence of a suspicious ovarian cyst with elevated cancer antigen 125 level in a woman of reproductive age poses a serious therapeutic dilemma. Mature cystic teratomas and mucinous cystadenomas may also cause an increase in cancer antigen 125. A 43-year-old Sinhalese woman with a history of anovulatory subfertility for 5 years presented with heavy menstrual bleeding and secondary ...

Difficulties in diagnosis of a minimal deviation ...
https://link.springer.com/article/10.1007/s00404-019-05286-7
Sep 16, 2019 Rare minimal deviation adenocarcinoma (MDA) diagnosed postoperatively as incidental finding of a suspicious cervical lesion at laparoscopy, emphasizing it represents a diagnostic challenge mimicking both benign and malignant cervical lesions with often overlapping imaging characteristics—case report and literature review. 35-year-old Gravida with primary infertility presented ...
Author: Alexandra Stoehr, Dominik Nann, Annet... Publish Year: 2019

Adenocarcinoma

Adenocarcinoma is a type of cancer that starts in the glandular cells of the body. It is defined as neoplasia of epithelial tissue that has glandular origin, glandular characteristics, or both. Adenocarcinomas are part of the larger grouping of carcinomas, but are also sometimes called by more precise terms omitting the word, where these exist. Thus invasive ductal carcinoma, the most common form of breast cancer, is adenocarcinoma but does not use the term in its name—however, esophageal adenocarcinoma does to distinguish it from the other common type of esophageal cancer, esophageal squamous cell carcinoma. Several of the most common forms of cancer are adenocarcinomas, and the various sorts of adenocarcinoma vary greatly in all their aspects, so that few useful generalizations can be made about them.

Wikipedia

People also search for

- Prostate Cancer
Carcinoma
Colon Cancer
Tumor
Non-small cell lung cancer
See more

Data from: Wikipedia
Suggest an edit

See results for

Endometrial Cancer (Medical Condition)
A cancer that begins in the lining of the uterus (endometrium). This causes vaginal bleeding after the menopause or frequent bleeding between the ...

Search Tools

Turn off Hover Translation (关闭取读)

Name of Journal: *World Journal of Clinical Cases*

Manuscript NO: 63051

Manuscript Type: CASE REPORT

Minimal deviation adenocarcinoma with elevated CA19-9: A case report

Dong Y *et al.* Minimal deviation adenocarcinoma

Yan Dong, Ying Lv, Jing Guo, Lin Sun

Abstract

BACKGROUND

Minimal deviation adenocarcinoma is a rare malignant, high rate of misdiagnosis, high aggressiveness, diagnosis of disease according to histopathology. Surgical resection is the preferred and most effective treatment, but the outcomes are often unsatisfactory.

CASE SUMMARY

Match Overview

1	Internet 39 words crawled on 15-Feb-2020 publishing.sib.care.windows.net	2%
2	Internet 17 words crawled on 23-Apr-2020 www.sjgnet.com	1%
3	Internet 14 words crawled on 10-Jun-2019 www.tandfonline.com	1%
4	Crossref 13 words Qin Yang, Wei Jiang, Lei Li, Qin Huang, Kai-xuan Yang. "For khead box profen P1 is a useful marker for the diagnosis r...	1%
5	Crossref 12 words Wenjun Xiong, Haipeng Huang, Yuling Xue, Jin Li, Jin Wan, Wei Wang. "Treatment of gastrointestinal stromal tumor w...	1%
6	Internet 11 words crawled on 27-Apr-2020 pericles.pericles-prod.literatumonline.com	1%
7	Internet 11 words crawled on 17-Mar-2021 www.frontiersin.org	1%
8	Internet 9 words crawled on 28-Jul-2020 world/science.org	1%

国内版 国际版

Minimal deviation adenocarcinoma with elevated CA19-9: A case re



ALL IMAGES VIDEOS

85,300 Results Any time ▾

Minimal deviation mucinous adenocarcinoma ('adenoma ...

https://www.researchgate.net/publication/40893939_Minimal_deviation_mucinous...

Serum CA19-9 levels were elevated to 486 U/mL. ... overlapping imaging characteristics-case report and literature review. ... case of minimal deviation adenocarcinoma of endometrioid type ...

Minimal deviation mucinous adenocarcinoma ('adenoma ...

<https://onlinelibrary.wiley.com/doi/abs/10.1111/j.1440-1827.2009.02473.x>

Dec 28, 2009 · Minimal deviation mucinous adenocarcinoma ('adenoma malignum') of the uterine corpus. ... A case is described of a 56-year-old postmenopausal woman who presented with mucinous...

Cited by: 17 Author: Kaoru Abiko, Tsukasa Baba, Madoka Og...

Publish Year: 2010

A case of CA19-9-producing adenocarcinoma of the remnant ...

<https://www.researchgate.net/publication/229550102...>

We report a case of early gastric cancer with a high level of serum CA19-9 (400IU/ml). A 62-year-old man was admitted to the hospital because of abnormality of the stomach by screening.

Primary adenocarcinoma of the rete testis with elevated ...

<https://link.springer.com/content/pdf/10.1007/s13691-020-00437-z.pdf>

Primary adenocarcinoma of the rete testis is an extremely rare tumor with a poor prognosis. Herein, we report a case of primary adenocarcinoma of the rete testis accompanied by elevated serum...

Author: Hiroyuki Kitano, Kazuhiro Sentani, Ke... Publish Year: 2020

Minimal deviation mucinous adenocarcinoma ('adenoma ...

<https://onlinelibrary.wiley.com/doi/10.1111/j.1440-1827.2009.02473.x>

Primary mucinous adenocarcinomas of the uterine corpus are typically low grade and frequently associated with endometrial hyperplasia and/or ordinary endometrioid adenocarcinoma, but ...

Cited by: 17 Author: Kaoru Abiko, Tsukasa Baba, Madoka Og...

Publish Year: 2010

[PDF] Diagnostic challenges in minimal deviation ...

<https://www.spandidos-publications.com/10.3892/mco.2013.144/download>

Minimal deviation adenocarcinoma (MDA) of the uterine cervix, otherwise known as adenoma