



23035-Review

BY JU YUP LEE

Quotes Excluded  
Bibliography Excluded24%  
SIMILAR

Name of journal: *World Journal of Gastrointestinal Endoscopy*

ESPS Manuscript NO: 23035

Manuscript Type: ORIGINAL ARTICLE

Retrospective Cohort Study

**Risk factors for local recurrence after en bloc endoscopic submucosal dissection for early gastric cancer**

**Ju Yup Lee, Kwang Bum Cho, Eun Soo Kim, Kyung Sik Park, Yoo Jin Lee, Yoon Suk Lee, Byoung Kuk Jang, Woo Jin Chung, and Jae Seok Hwang**

### Match Overview

|   |   |    |
|---|---|----|
| 1 | <b>CrossCheck</b> 104 words<br>"Poster - Endoscopy : Poster - Endoscopy", Journal of Gastroenterology and Hepatology, 2015.                       | 2% |
| 2 | <b>CrossCheck</b> 98 words<br>Takada, Jun, Hiroshi Araki, Fumito Onogi, Takayuki Nakamishi, Masaya Kubota, Takashi Ibuka, Masahito Shimizu        | 2% |
| 3 | <b>CrossCheck</b> 70 words<br>Min, Yang Won. "Endoscopic treatment for early gastric cancer", World Journal of Gastroenterology, 2014.            | 2% |
| 4 | <b>CrossCheck</b> 52 words<br>Matsumura, Tomoaki, Makoto Arai, Daisuke Maruoka, Kenichiro Okimoto, Shoko Minemura, Hideaki Ishigami, Kei          | 1% |
| 5 | <b>CrossCheck</b> 51 words<br>Ken Ohnita. "Factors related to the curability of early gastric cancer with endoscopic submucosal dissection", Surg | 1% |
| 6 | <b>CrossCheck</b> 46 words<br>Kwon, Young Min. "Factors associated with use of gastric cancer screening services in Korea", World Journal of G    | 1% |



Risk factors for local recurrence after en bloc endoscopic submucosal dis

[全部](#)[新闻](#)[图片](#)[视频](#)[更多 ▾](#)[搜索工具](#)

找到约 3,850 条结果 (用时 0.75 秒)

### The Impact of Human T-Cell Lymphotropic Virus I Infection ...

[www.ncbi.nlm.nih.gov](http://www.ncbi.nlm.nih.gov) > ... > [PubMed Central \(PMC\)](#) ▾ [翻译此页](#)

作者: F Bahia - 2011 - 被引用次数: 9 - 相关文章

HIV, hepatitis C (HCV), and human T-cell lymphotropic virus I (HTLV-1) are ... of triply infected patients (Group 2) reported any history of injection drug use ... HIV/HTLV-1 and HIV/HCV coinfections may worsen clinical related outcomes, but virologic ... HIV and HTLV-1 coinfection has been found in Bahia with worse morbidity ...

### [PDF] Prevention and control of infectious diseases among peo...

[ecdc.europa.eu/en/.../111012\\_Guidance\\_ECDC-EMCDDA.pdf](http://ecdc.europa.eu/en/.../111012_Guidance_ECDC-EMCDDA.pdf) ▾ [翻译此页](#)

interventions to prevent infections among people who inject drugs. .... a large burden on European health systems, significant individual suffering, .... with more than one type of hepatitis, or co-infection with HIV and hepatitis C ... In addition to HIV and hepatitis B and C, infections caused by human T-lymphotropic virus type II ...

### Burden of hepatitis C virus disease and access to hepatitis ...

[www.lancet.com/journals/laninf/article/PIIS1473...X/fulltext](http://www.lancet.com/journals/laninf/article/PIIS1473...X/fulltext) ▾ [翻译此页](#)

2014年12月2日 - 90% of individuals infected with hepatitis C virus (HCV) worldwide reside in ... Weighted HCV prevalence was 5777 (37.2%) of 14 447; HIV/HCV co-infection ... Of the 5777 people who inject drugs that were HCV antibody positive, 440



Risk factors for local recurrence after en bloc endoscopic submucosal dis

[全部](#)[新闻](#)[图片](#)[视频](#)[更多](#)[搜索工具](#)

找到约 3,850 条结果 (用时 0.75 秒)

### The Impact of Human T-Cell Lymphotropic Virus I Infection ...

[www.ncbi.nlm.nih.gov](http://www.ncbi.nlm.nih.gov) > ... > [PubMed Central \(PMC\)](#) ▾ [翻译此页](#)

作者: F Bahia - 2011 - 被引用次数: 9 - 相关文章

HIV, hepatitis C (HCV), and human T-cell lymphotropic virus I (HTLV-1) are ... of triply infected patients (Group 2) reported any history of injection drug use ... HIV/HTLV-1 and HIV/HCV coinfections may worsen clinical related outcomes, but virologic ... HIV and HTLV-1 coinfection has been found in Bahia with worse morbidity ...

### [PDF] Prevention and control of infectious diseases among peo...

[ecdc.europa.eu/en/.../111012\\_Guidance\\_ECDC-EMCDDA.pdf](http://ecdc.europa.eu/en/.../111012_Guidance_ECDC-EMCDDA.pdf) ▾ [翻译此页](#)

interventions to prevent infections among people who inject drugs. .... a large burden on European health systems, significant individual suffering, .... with more than one type of hepatitis, or co-infection with HIV and hepatitis C ... In addition to HIV and hepatitis B and C, infections caused by human T-lymphotropic virus type II ...

### Burden of hepatitis C virus disease and access to hepatitis ...

[www.lancet.com/journals/laninf/article/PIIS1473...X/fulltext](http://www.lancet.com/journals/laninf/article/PIIS1473...X/fulltext) ▾ [翻译此页](#)

2014年12月2日 - 90% of individuals infected with hepatitis C virus (HCV) worldwide reside in ... Weighted HCV prevalence was 5777 (37.2%) of 14 447; HIV/HCV co-infection ... Of the 5777 people who inject drugs that were HCV antibody positive, 440

human immunodeficiency, and human T-lymphotropic viruses



[全部](#)[图片](#)[新闻](#)[视频](#)[更多 ▾](#)[搜索工具](#)

找到约 28,600 条结果 (用时 1.19 秒)

## Google 学术: Risk factors for local recurrence after en bloc endoscopic submucosal dissection for early gastric cancer

Advantage of endoscopic submucosal dissection ... - Oka - 被引用次数: 538

Endoscopic submucosal dissection for early gastric ... - Isomoto - 被引用次数: 379

... of early gastric cancers after endoscopic submucosal ... - Takenaka - 被引用次数: 112

### [PDF] PDF (741 KB) - Advances in Digestive Medicine

[aidm-online.com/article/S2351-9797\(15\)00008-0/pdf](http://aidm-online.com/article/S2351-9797(15)00008-0/pdf) ▾ 翻译此页

作者: OB Panel - 被引用次数: 96 - 相关文章

2015年4月7日 - endoscopic submucosal dissection (ESD) for early gastric cancer and precancer lesion in Taiwan. In this long-term ... lesions) were analyzed for local recurrence after ESD. Results and ... analyzed the risk factors that affect the local recurrence .... CR Z curative resection; E(κ) Z en bloc resection; E(A) Z no ...

### Outcomes of endoscopic submucosal dissection for early ...

[www.sciencedirect.com/science/article/pii/S2351979715000080](http://www.sciencedirect.com/science/article/pii/S2351979715000080) - 翻译此页

作者: YY Hsieh - 2015 - 被引用次数: 1 - 相关文章

Outcomes of endoscopic submucosal dissection for early gastric cancer and ... In previous long-term follow-up studies, risk factors for local recurrence, such as EGC .... Characteristics of the 48 early gastric cancer lesions after curative resection. ... E(-) = no en-bloc resection; EGC = early gastric cancer; ESD = endoscopic ...

### Sa1638 Risk Factors for Local Recurrence After En Bloc ...

[... \(2013-05-13\) ...](#) 翻译此页