$\underline{Gastrointestinal\ Complications\ (PDQ@)-Health\ Professional\ ...}$ $\underline{https://www.cancer.gov/about-cancer/treatment/side-effects/constipation/gi... \bullet$ Apr 22, 2020 - With diarrhea, the digestive and absorptive functions of the gastrointestinal (GI) tract are altered or lost, resulting in malabsorption of fat, lactose, bile salts, and vitamin B 12. Symptoms of proctitis—including mucoid rectal discharge, rectal pain, and rectal bleeding (if mucosal ulceration is present)-may result from radiation damage to the anus or rectum.

Endoscopic Management of Tumor Bleeding from Inoperable ...

https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4381138 Mar 27, 2015 · INTRODUCTION. Tumor bleeding accounts for up to 5% of upper gastrointestinal bleeding (UGIB) cases.1,2,3,4,5 Of these, primary gastric cancer is the most common cause of tumor bleeding, accounting for between 36% and 58% of bleeding cases resulting from upper gastrointestinal malignancies.1,3,4 However, little information is available regarding the current management of gastric cancer bleeding.

Cited by: 24 Author: Young-II Kim, II Ju Choi Publish Year: 2015


People also ask

177,000 Results Any time *

What is the percentage of lower gastrointestinal bleeding? What are the risk factors for upper GI bleeding? How are gastrointestinal problems related to cancer treatment?	`
	8

Search Tools

Turn off Hover Translation (关闭取词)



Gastrointestinal hemorrhage in the setting of gastrointestinal cano







ALL

IMAGES

VIDEOS

175,000 Results

Any time ▼

Gastrointestinal Complications (PDQ®)-Health Professional ...

https://www.cancer.gov/about-cancer/treatment/side-effects/constipation/gi... •

Apr 22, 2020 · With diarrhea, the **digestive** and absorptive functions of the **gastrointestinal** (GI) tract are altered or lost, resulting in malabsorption of fat, lactose, bile salts, and vitamin B 12. Symptoms of proctitis—including mucoid rectal discharge, rectal pain, and rectal **bleeding** (if mucosal ulceration is present)—may result from radiation damage to the anus or rectum.

Endoscopic Management of Tumor Bleeding from ...

https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4381138

Mar 27, 2015 · INTRODUCTION. Tumor bleeding accounts for up to 5% of upper gastrointestinal bleeding (UGIB) cases:1,2,3,4,5 Of these, primary gastric cancer is the most common cause of tumor bleeding, accounting for between 36% and 58% of bleeding cases resulting from upper gastrointestinal malignancies:1,3,4 However, little information is available regarding the current management of gastric cancer bleeding.

Cited by: 24 Author: Young-Il Kim, Il Ju Choi

Publish Year: 2015



Feedback

An Evidence-Based Review of Gastrointestinal Bleeding ... https://www.reliasmedia.com/articles/144557-an-evidence-based-review-of...

Jun 15, 2019 · Compared to those with upper GI bleeding, lower GI bleeding patients are more likely to have higher levels of hemoglobin (84% vs. 61%) and less likely to experience shock (19% vs. 35%). 1 Bleeds that originate from the colon require fewer blood transfusions than those that originate from the