



Baishideng Publishing Group Co., Limited

Flat C, 23/F., Lucky Plaza,
315-321 Lockhart Road,
Wan Chai, Hong Kong, China

ESPS Peer-review Report

Name of Journal: World Journal of Gastroenterology

ESPS Manuscript NO: 3469

Title: Transcatheter arterial chemoembolisation followed by immediate radiofrequency ablation for solitary large hepatocellular carcinoma

Reviewer code: 01752190

Science editor: Wang, Jin-Lei

Date sent for review: 2013-05-02 16:43

Date reviewed: 2013-05-10 09:17

CLASSIFICATION	LANGUAGE EVALUATION	RECOMMENDATION	CONCLUSION
<input type="checkbox"/> Grade A (Excellent)	<input type="checkbox"/> Grade A: Priority Publishing	Google Search:	<input type="checkbox"/> Accept
<input type="checkbox"/> Grade B (Very good)	<input type="checkbox"/> Grade B: minor language polishing	<input type="checkbox"/> Existed	<input type="checkbox"/> High priority for publication
<input type="checkbox"/> Grade C (Good)	<input type="checkbox"/> Grade C: a great deal of language polishing	<input type="checkbox"/> No records	<input type="checkbox"/> Rejection
<input type="checkbox"/> Grade D (Fair)	<input type="checkbox"/> Grade D: rejected	<input type="checkbox"/> Existed	<input type="checkbox"/> Minor revision
<input type="checkbox"/> Grade E (Poor)		<input type="checkbox"/> No records	<input type="checkbox"/> Major revision

COMMENTS TO AUTHORS

Transcatheter arterial chemoembolisation (TACE) is one of the measures for treating unresectable HCC; however, its low complete tumour necrosis rate results in tumour recurrence and metastasis and influences long-term efficacy. TACE immediately followed by radiofrequency ablation under digital subtraction angiography-computed tomography is first used to treat large hepatocellular carcinoma. In this manuscript, the authors summarised 18 patients between January 2010 and June 2012 to assess the technical safety and efficacy of TACE combined with immediate synchronous RFA as a means of treating HCC. The manuscript is very interesting.



Baishideng Publishing Group Co., Limited

Flat C, 23/F., Lucky Plaza,
315-321 Lockhart Road,
Wan Chai, Hong Kong, China

ESPS Peer-review Report

Name of Journal: World Journal of Gastroenterology

ESPS Manuscript NO: 3469

Title: Transcatheter arterial chemoembolisation followed by immediate radiofrequency ablation for solitary large hepatocellular carcinoma

Reviewer code: 01220970

Science editor: Wang, Jin-Lei

Date sent for review: 2013-05-02 16:43

Date reviewed: 2013-05-14 16:48

CLASSIFICATION	LANGUAGE EVALUATION	RECOMMENDATION	CONCLUSION
<input type="checkbox"/> Grade A (Excellent)	<input type="checkbox"/> Grade A: Priority Publishing	Google Search:	<input type="checkbox"/> Accept
<input checked="" type="checkbox"/> Grade B (Very good)	<input checked="" type="checkbox"/> Grade B: minor language polishing	<input type="checkbox"/> Existed	<input checked="" type="checkbox"/> High priority for publication
<input type="checkbox"/> Grade C (Good)	<input type="checkbox"/> Grade C: a great deal of language polishing	<input type="checkbox"/> No records	<input type="checkbox"/> Rejection
<input type="checkbox"/> Grade D (Fair)		BPG Search:	<input type="checkbox"/> Minor revision
<input type="checkbox"/> Grade E (Poor)	<input type="checkbox"/> Grade D: rejected	<input type="checkbox"/> Existed	<input type="checkbox"/> Major revision
		<input type="checkbox"/> No records	

COMMENTS TO AUTHORS

Very good study about the technical safety and efficacy of transcatheter arterial chemoembolisation combined with immediate radiofrequency ablation for large hepatocellular carcinoma.