

Microsoft Bing

Analysis of influencing factors for hepatic fat accumulation diagnosis

Sign in

国内版 国际版

ALL IMAGES VIDEOS

51,500 Results Any time Open links in new tab

Nonalcoholic fatty liver disease: cause or consequence of ...

<https://onlinelibrary.wiley.com/doi/full/10.1111/liv.13185>

Furthermore, genetic studies suggest that excess hepatic fat is associated with progressive liver disease, but does not always increase the risk of incident type 2 diabetes. Here, we will briefly review the epidemiological, pathophysiological and molecular evidence linking NAFLD to the development of type 2 diabetes.

Cited by: 104 Author: Luca Valenti, Luca Valenti, Elisabetta Bug... Publish Year: 2016

Quantification of Liver Fat by MRI-PDFF Imaging in ...

<https://www.sciencedirect.com/science/article/pii/S0973688320301778>

Dec 01, 2020 - Four MR-based diagnostic methods are currently available for liver fat quantification: (1) Dixon MRI technique; (2) modified Dixon type (mDixon); (3) single proton-MR spectroscopy; and 4) PDFF. 16 There are a number of studies that have reported excellent diagnostic performance of MRS to detect and quantify the hepatic steatosis.17, 18, 19 ...

Author: Gajanan A. Rodge, Mahesh K. Goenk... Publish Year: 2020

Search Tools

Turn off Hover Translation (关闭翻译)

PEOPLE ALSO ASK

How is liver fat measured in MRI - PDFF?

How is insulin resistance associated with fatty liver disease?

What are the serum markers of hepatic fibrosis?



435,000 Results

Any time ▾

The **influencing factors** for **type 2 diabetes mellitus** and non-alcoholic fatty liver disease include increased the serum fetuin B level in patients with newly diagnosed **type 2 diabetes mellitus** and resulted in a much higher serum fetuin B level in the group complicated with NAFLD. 2.

[Serum Fetuin B Levels and Factors Influencing its ...](#)

juniperpublishers.com/crdoj/CRDOJ.MS.ID.555843.php

Was this helpful?



PEOPLE ALSO ASK

How is fatty liver disease related to diabetes mellitus? ▾

How often does hepatic fat accumulation occur in diabetes? ▾

Is there a link between hepatic steatohepatitis and diabetes? ▾

Which is associated with fat accumulation type 1 or 2? ▾

Feedback

Fatty liver disease in diabetes mellitus


<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4405411>

Jan 03, 2015 · Non-alcoholic fatty liver disease (NAFLD) is highly prevalent in **type 2 diabetes mellitus (T2DM)**, likely reflecting the frequent occurrence of obesity and insulin resistance in T2DM. **NAFLD** also can occur in **type 1 DM (T1DM)**, but must be distinguished from the more common glycogen hepatopathy

22-Jul-2021 03:41 PM

3705 words • 12 matches • 10 sources

FAQ

iThenticate®

66199-Review-check.docx

Quotes Excluded
Bibliography Excluded

5%
Similarity

Name of Journal: *World Journal of Clinical Cases*

Manuscript NO: 66199

Manuscript Type: ORIGINAL ARTICLE

Retrospective Study

Influencing factors for hepatic fat accumulation in patients with type 2 diabetes mellitus

Wu MJ *et al.* T2DM and hepatic fat accumulation

Meng-Jiao Wu, Qiong-Lei Fang, Sheng-Yi Zou, Yan Zhu, Wen Lu, Xuan Du, Bi-Min Shi

Match Overview

1	Internet 34 words crawled on 05-Oct-2020 pesquisa.bvsalud.org	1%
2	Internet 27 words crawled on 24-Dec-2019 www.wjgnet.com	1%
3	Internet 23 words crawled on 21-Apr-2021 medrxiv1885journals.com	1%
4	Crossref 15 words "APASL Seoul 2008 Meeting", <i>Hepatology International</i> , 2008	<1%
5	Internet 15 words crawled on 16-Jul-2020 lipidworld.biomedcentral.com	<1%
6	Internet 15 words crawled on 08-Oct-2020 www.researchsquare.com	<1%

PAGE 1 OF 13

Text-Only Report

国内版

国际版

Influencing factors for hepatic fat accumulation in patients with typ



ALL

IMAGES

VIDEOS

431,000 Results

Any time ▾

The **influencing factors** for **type 2 diabetes mellitus** and non-alcoholic fatty liver disease include increased the serum fetuin B level in patients with newly diagnosed type **2 diabetes mellitus** and resulted in a much higher serum fetuin B level in the group complicated with NAFLD. 2.

[Serum Fetuin B Levels and Factors Influencing its ...
juniperpublishers.com/crdoj/CRDOJ.MS.ID.555843.php](https://juniperpublishers.com/crdoj/CRDOJ.MS.ID.555843.php)

Was this helpful?

PEOPLE ALSO ASK

How is fatty liver disease related to diabetes mellitus? ▾

How often does hepatic fat accumulation occur in diabetes? ▾

Is there a link between hepatic steatohepatitis and diabetes? ▾

Which is associated with fat accumulation type 1 or 2? ▾

Feedback

Fatty liver disease in diabetes mellitus

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4405411>

Jan 03, 2015 · Non-alcoholic fatty liver disease (NAFLD) is highly prevalent in **type 2 diabetes mellitus (T2DM)**, likely reflecting the frequent occurrence of obesity and insulin resistance in T2DM. NAFLD also can occur in **type 1 DM (T1DM)**, but must be distinguished from the more common glycogen hepatopathy as a cause of hepatomegaly and liver function abnormalities in T1DM.

Cited by: 140

Author: Harikrashna B. Bhatt, Robert J. Smith

Publish Year: 2015

Nonalcoholic fatty liver disease: a frequent condition in ...

<https://pubmed.ncbi.nlm.nih.gov/24652019>

Conclusion: We observed an elevated prevalence of nonalcoholic **fatty liver disease** (87.1 %) in type 2