

World Journal of *Clinical Cases*

World J Clin Cases 2022 August 16; 10(23): 8057-8431



Contents

Thrice Monthly Volume 10 Number 23 August 16, 2022

OPINION REVIEW

- 8057** Invasive intervention timing for infected necrotizing pancreatitis: Late invasive intervention is not late for collection

Xiao NJ, Cui TT, Liu F, Li W

- 8063** Clinical utility of left atrial strain in predicting atrial fibrillation recurrence after catheter ablation: An up-to-date review

Yu ZX, Yang W, Yin WS, Peng KX, Pan YL, Chen WW, Du BB, He YQ, Yang P

MINIREVIEWS

- 8076** Gut microbiota and COVID-19: An intriguing pediatric perspective

Valentino MS, Esposito C, Colosimo S, Caprio AM, Puzone S, Guarino S, Marzuillo P, Miraglia del Giudice E, Di Sessa A

- 8088** Beta receptor blocker therapy for the elderly in the COVID-19 era

Santillo E, Migale M

ORIGINAL ARTICLE

Retrospective Cohort Study

- 8097** Nonselective beta-blocker use is associated with increased hepatic encephalopathy-related readmissions in cirrhosis

Fallahzadeh MA, Asrani SK, Tapper EB, Saracino G, Rahimi RS

Retrospective Study

- 8107** Different squatting positions after total knee arthroplasty: A retrospective study

Li TJ, Sun JY, Du YQ, Shen JM, Zhang BH, Zhou YG

- 8115** Outcomes of seromuscular bladder augmentation compared with standard bladder augmentation in the treatment of children with neurogenic bladder

Sun XG, Li YX, Ji LF, Xu JL, Chen WX, Wang RY

- 8124** Distinctive clinical features of spontaneous pneumoperitoneum in neonates: A retrospective analysis

Kim SH, Cho YH, Kim HY

- 8133** Cognitive training for elderly patients with early Alzheimer's disease in the Qinghai-Tibet Plateau: A pilot study

Wang XH, Luo MQ

- 8141** Diagnostic value of elevated serum carbohydrate antigen 125 level in sarcoidosis

Zhang Q, Jing XY, Yang XY, Xu ZJ

- 8152** Evaluation of progressive early rehabilitation training mode in intensive care unit patients with mechanical ventilation

Qie XJ, Liu ZH, Guo LM

- 8161** Comparison of demographic features and laboratory parameters between COVID-19 deceased patients and surviving severe and critically ill cases

Wang L, Gao Y, Zhang ZJ, Pan CK, Wang Y, Zhu YC, Qi YP, Xie FJ, Du X, Li NN, Chen PF, Yue CS, Wu JH, Wang XT, Tang YJ, Lai QQ, Kang K

Clinical Trials Study

- 8170** Role of H₂receptor blocker famotidine over the clinical recovery of COVID-19 patients: A randomized controlled trial

Mohiuddin Chowdhury ATM, Kamal A, Abbas MKU, Karim MR, Ali MA, Talukder S, Hamidullah Mehedi H, Hassan H, Shahin AH, Li Y, He S

Observational Study

- 8186** Short-term prognostic factors for hepatitis B virus-related acute-on-chronic liver failure

Ye QX, Huang JF, Xu ZJ, Yan YY, Yan Y, Liu LG

- 8196** Three-dimensional psychological guidance combined with evidence-based health intervention in patients with liver abscess treated with ultrasound

Shan YN, Yu Y, Zhao YH, Tang LL, Chen XM

- 8205** Role of serum β 2-microglobulin, glycosylated hemoglobin, and vascular endothelial growth factor levels in diabetic nephropathy

Yang B, Zhao XH, Ma GB

SYSTEMATIC REVIEWS

- 8212** Gallbladder neuroendocrine carcinoma diagnosis, treatment and prognosis based on the SEER database: A literature review

Cai XC, Wu SD

CASE REPORT

- 8224** Sepsis complicated with secondary hemophagocytic syndrome induced by giant gouty tophi rupture: A case report

Lai B, Pang ZH

- 8232** Spontaneous remission of autoimmune pancreatitis: Four case reports

Zhang BB, Huo JW, Yang ZH, Wang ZC, Jin EH

- 8242** Epstein-Barr-virus-associated hepatitis with aplastic anemia: A case report

Zhang WJ, Wu LQ, Wang J, Lin SY, Wang B

- 8249** Aspiration as the first-choice procedure for airway management in an infant with large epiglottic cysts: A case report

Zheng JQ, Du L, Zhang WY

- 8255** Sequential multidisciplinary minimally invasive therapeutic strategy for heart failure caused by four diseases: A case report
Zhao CZ, Yan Y, Cui Y, Zhu N, Ding XY
- 8262** Primary ascending colon cancer accompanying skip metastases in left shoulder skin and left neck lymph node: A case report
Zhou JC, Wang JJ, Liu T, Tong Q, Fang YJ, Wu ZQ, Hong Q
- 8271** Clinical and genetic study of ataxia with vitamin E deficiency: A case report
Zhang LW, Liu B, Peng DT
- 8277** Complete resection of large-cell neuroendocrine and hepatocellular carcinoma of the liver: A case report
Noh BG, Seo HI, Park YM, Kim S, Hong SB, Lee SJ
- 8284** Immunotherapy combined with antiangiogenic agents in patients with advanced malignant pleural mesothelioma: A case report
Xuan TT, Li GY, Meng SB, Wang ZM, Qu LL
- 8291** Bladder malacoplakia: A case report
Wang HK, Hang G, Wang YY, Wen Q, Chen B
- 8298** Delayed inflammatory response evoked in nasal alloplastic implants after COVID-19 vaccination: A case report
Seo MG, Choi EK, Chung KJ
- 8304** Phosphoglyceride crystal deposition disease requiring differential diagnosis from malignant tumors and confirmed by Raman spectroscopy: A case report
Ohkura Y, Uruga H, Shiiba M, Ito S, Shimoyama H, Ishihara M, Ueno M, Udagawa H
- 8312** Vulvovaginal myeloid sarcoma with massive pelvic floor infiltration: A case report and review of literature
Wang JX, Zhang H, Ning G, Bao L
- 8323** Femoral neck stress fracture and medial tibial stress syndrome following high intensity interval training: A case report and review of literature
Tan DS, Cheung FM, Ng D, Cheung TLA
- 8330** Periosteal chondroma of the rib: A case report
Gao Y, Wang JG, Liu H, Gao CP
- 8336** Papillary thyroid carcinoma occurring with undifferentiated pleomorphic sarcoma: A case report
Lee YL, Cheng YQ, Zhu CF, Huo HZ
- 8344** Laparoscopic treatment of bilateral duplex kidney and ectopic ureter: A case report
Wang SB, Wan L, Wang Y, Yi ZJ, Xiao C, Cao JZ, Liu XY, Tang RP, Luo Y
- 8352** Incontinentia pigmenti with intracranial arachnoid cyst: A case report
Li WC, Li ML, Ding JW, Wang L, Wang SR, Wang YY, Xiao LF, Sun T

- 8360** Relapsing polychondritis causing breathlessness: Two case reports
Zhai SY, Zhang YH, Guo RY, Hao JW, Wen SX
- 8367** Endodontic management of a fused left maxillary second molar and two paramolars using cone beam computed tomography: A case report
Mei XH, Liu J, Wang W, Zhang QX, Hong T, Bai SZ, Cheng XG, Tian Y, Jiang WK
- 8375** Infant biliary cirrhosis secondary to a biliary inflammatory myofibroblastic tumor: A case report and review of literature
Huang Y, Shu SN, Zhou H, Liu LL, Fang F
- 8384** Metastatic low-grade endometrial stromal sarcoma with variable morphologies in the ovaries and mesentery: A case report
Yu HY, Jin YL
- 8392** Bronchogenic cysts with infection in the chest wall skin of a 64-year-old asymptomatic patient: A case report
Ma B, Fu KW, Xie XD, Cheng Y, Wang SQ
- 8400** Incidental accumulation of Technetium-99m pertechnetate in subacute cerebral infarction: A case report
Han YH, Jeong HJ, Kang HG, Lim ST
- 8406** Metal stent combined with ileus drainage tube for the treatment of delayed rectal perforation: A case report
Cheng SL, Xie L, Wu HW, Zhang XF, Lou LL, Shen HZ
- 8417** Using ketamine in a patient with a near-occlusion tracheal tumor undergoing tracheal resection and reconstruction: A case report
Xu XH, Gao H, Chen XM, Ma HB, Huang YG

LETTER TO THE EDITOR

- 8422** Reflections on the prevalence of human leukocyte antigen-B27 and human leukocyte antigen-B51 co-occurrence in patients with spondylarthritis
Gonçalves Júnior J, Sampaio-Barros PD, Shinjo SK
- 8425** Comment on "Disease exacerbation is common in inflammatory bowel disease patients treated with immune checkpoint inhibitors for malignancy"
Argyriou K, Kotsakis A
- 8428** Intranasal sufentanil combined with intranasal dexmedetomidine: A promising method for non-anesthesiologist sedation during endoscopic ultrasonography
Wang Y, Ge ZJ, Han C

ABOUT COVER

Editorial Board Member of *World Journal of Clinical Cases*, Peng Liang, MD, Associate Professor, Day Surgery Center, Department of Anesthesiology, West China Hospital of Sichuan University, Chengdu 610041, Sichuan Province, China. 39485572@qq.com

AIMS AND SCOPE

The primary aim of *World Journal of Clinical Cases* (WJCC, *World J Clin Cases*) is to provide scholars and readers from various fields of clinical medicine with a platform to publish high-quality clinical research articles and communicate their research findings online.

WJCC mainly publishes articles reporting research results and findings obtained in the field of clinical medicine and covering a wide range of topics, including case control studies, retrospective cohort studies, retrospective studies, clinical trials studies, observational studies, prospective studies, randomized controlled trials, randomized clinical trials, systematic reviews, meta-analysis, and case reports.

INDEXING/ABSTRACTING

The WJCC is now abstracted and indexed in Science Citation Index Expanded (SCIE, also known as SciSearch®), Journal Citation Reports/Science Edition, Current Contents®/Clinical Medicine, PubMed, PubMed Central, Scopus, Reference Citation Analysis, China National Knowledge Infrastructure, China Science and Technology Journal Database, and Superstar Journals Database. The 2022 Edition of Journal Citation Reports® cites the 2021 impact factor (IF) for WJCC as 1.534; IF without journal self cites: 1.491; 5-year IF: 1.599; Journal Citation Indicator: 0.28; Ranking: 135 among 172 journals in medicine, general and internal; and Quartile category: Q4. The WJCC's CiteScore for 2021 is 1.2 and Scopus CiteScore rank 2021: General Medicine is 443/826.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: Hua-Ge Yin; Production Department Director: Xiang Li; Editorial Office Director: Jin-Lei Wang.

NAME OF JOURNAL

World Journal of Clinical Cases

ISSN

ISSN 2307-8960 (online)

LAUNCH DATE

April 16, 2013

FREQUENCY

Thrice Monthly

EDITORS-IN-CHIEF

Bao-Gan Peng, Jerzy Tadeusz Chudek, George Kontogeorgos, Maurizio Serati, Ja Hyeon Ku

EDITORIAL BOARD MEMBERS

<https://www.wjnet.com/2307-8960/editorialboard.htm>

PUBLICATION DATE

August 16, 2022

COPYRIGHT

© 2022 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>



Retrospective Cohort Study

Nonselective beta-blocker use is associated with increased hepatic encephalopathy-related readmissions in cirrhosis

Mohammad Amin Fallahzadeh, Sumeet K Asrani, Elliot B Tapper, Giovanna Saracino, Robert S Rahimi

Specialty type: Gastroenterology and hepatology

Provenance and peer review: Unsolicited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's scientific quality classification

Grade A (Excellent): 0
Grade B (Very good): 0
Grade C (Good): C, C
Grade D (Fair): 0
Grade E (Poor): 0

P-Reviewer: Xu CF, China; Zhang LL, China

Received: March 7, 2022

Peer-review started: March 7, 2022

First decision: April 5, 2022

Revised: April 13, 2022

Accepted: July 11, 2022

Article in press: July 11, 2022

Published online: August 16, 2022



Mohammad Amin Fallahzadeh, Department of Internal Medicine, Baylor University Medical Center, Dallas, TX 75246, United States

Sumeet K Asrani, Giovanna Saracino, Robert S Rahimi, Division of Hepatology, Baylor University Medical Center, Dallas, TX 75246, United States

Elliot B Tapper, Division of Hepatology, University of Michigan, Ann Arbor, MI 48109, United States

Corresponding author: Mohammad Amin Fallahzadeh, MD, Doctor, Department of Internal Medicine, Baylor University Medical Center, 3500 Gaston Ave, Dallas, TX 75246, United States. aminfa91@gmail.com

Abstract

BACKGROUND

Hepatic encephalopathy (HE) is a neurocognitive condition in cirrhosis leading to frequent hospitalizations. Nonselective beta-blockers (NSBBs) are the mainstay of pharmacologic treatment in cirrhotic patients. We hypothesized that since NSBBs decrease cardiac output and portal flow, the decreased metabolic filtering process of liver parenchyma may lead to increased HE-related hospitalizations.

AIM

To evaluate the impact of NSBB administration on HE-related readmissions in cirrhotic patients.

METHODS

In this retrospective cohort study, we included 393 patients admitted to Baylor University Medical Center for liver-related portal hypertension indications between January 2013 and July 2018. Independent predictors of the first HE-related readmissions were identified using Cox proportional hazards analysis. The cumulative incidence of the first HE-related readmissions between patients receiving NSBBs and not receiving NSBBs was examined using Fine-Gray modeling to account for the competing risk of death or liver transplantation.

RESULTS

The mean age was 58.1 ± 10.2 years and most patients fell into Child class C (49.1%) or B (43.8%). The median Model for End-Stage Liver Disease-Sodium score was 22 (IQR: 11). The cumulative incidence of the first HE-related

readmissions was significantly higher in patients taking NSBBs compared to patients not receiving NSBBs (71.8% *vs* 41.8%, $P < 0.0001$). In multivariate analysis, after adjusting for demographics, markers of liver disease severity, selective beta-blocker, lactulose and rifaximin use, NSBB use [Hazard ratio: 1.74 (95% CI: 1.29-2.34)] was independently associated with the first HE-related readmissions over a median follow-up of 3.8 years.

CONCLUSION

NSBB use is independently associated with increased HE-related readmissions in patients with cirrhosis, regardless of liver disease severity.

Key Words: Altered mental status; Ascites; Esophageal varices; Liver disease; Portal hypertension; Hospitalization

©The Author(s) 2022. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: In this study, we evaluated the impact of nonselective beta-blocker (NSBB) administration on hepatic encephalopathy (HE)-related readmissions in patients with Child B or C cirrhosis. After adjusting for markers of liver disease severity, NSBB use was independently associated with the first HE-related readmissions. NSBB use was also an independent predictor of HE-related admissions per person-month.

Citation: Fallahzadeh MA, Asrani SK, Tapper EB, Saracino G, Rahimi RS. Nonselective beta-blocker use is associated with increased hepatic encephalopathy-related readmissions in cirrhosis. *World J Clin Cases* 2022; 10(23): 8097-8106

URL: <https://www.wjgnet.com/2307-8960/full/v10/i23/8097.htm>

DOI: <https://dx.doi.org/10.12998/wjcc.v10.i23.8097>

INTRODUCTION

Hepatic encephalopathy (HE) is a reversible neurocognitive disorder seen in patients with advanced liver disease[1,2]. It is observed in up to 60% of patients with cirrhosis and is associated with frequent hospitalizations and decreased survival[3-5]. The incidence of HE-related hospitalizations is rising in the United States and imposes a significant economic burden on the healthcare system[6]. In a large population-based cohort, nonselective beta-blocker (NSBB) use was independently linked to HE development[7]. However, the mechanism is unclear.

NSBBs are the mainstay of pharmacologic treatment for portal hypertension and in the prevention of variceal bleeding in cirrhosis[8]. NSBB administration results in reduced cardiac output through inhibition of β_1 receptor and splanchnic vasoconstriction *via* antagonism of the β_2 receptor, leading to decreased portal inflow[9]. NSBB use may be associated with decreased survival in patients with refractory ascites, increased risk of acute kidney injury, and decreased transplant-free survival in patients with prior spontaneous bacterial peritonitis[10-13]. However, its role in the development of HE-related complications is not known.

We hypothesized that NSBB use contributes to decreased metabolic filtering process of the liver parenchyma, by way of decreased portal inflow (*i.e.*, similar to spontaneous portosystemic shunting), resulting in a secondary increase in HE-related hospitalizations independent of liver disease severity.

MATERIALS AND METHODS

Study population

In this observational, retrospective, single-center study, we examined all adults with cirrhosis following a liver-related hospitalization between January 2013 and July 2018. A hospitalization was considered liver-related if the primary or secondary cause of hospitalization was a portal hypertension-related complication such as HE, ascites, variceal bleeding, or hepatorenal syndrome. Patients were considered to have HE on hospital admission only if they had signs of overt HE (*i.e.*, HE grades II-IV) according to the International Society for Hepatic Encephalopathy and Nitrogen Metabolism and West Haven criteria, respectively[14,15].

Aims

Our primary aim was to examine the association between the use of NSBBs and subsequent HE-related readmissions. Our secondary aim was to identify factors that were predictive of HE-related admissions.

Study variables

We collected information about demographics (age, gender), liver disease [etiology of liver disease, history of hepatocellular carcinoma, esophageal varices (EV) and transjugular intrahepatic portosystemic shunt (TIPS)], physical examination findings in particular heart rate, presence of HE (with or without lactulose +/- rifaximin) and ascites as well as biochemical values including serum creatinine, total bilirubin, international normalized ratio, serum albumin, aspartate aminotransferase, alanine aminotransferase, platelet count and white blood cell count during each admission. The patients were divided into two groups according to whether they were receiving NSBB or not on the first liver-related hospitalization. We also gathered data about selective beta-blocker (SBB) use on the first liver-related hospitalization. Model for End-Stage Liver Disease (MELD), MELD-Sodium (MELD-Na) and Child-Turcotte-Pugh (CTP) scores were calculated for all patients during their first admission. Hospital course and outcome variables (*i.e.*, recurrent HE, death or liver transplantation) were determined during the follow-up period. Patients with a change in their NSBB or SBB status after the first liver-related hospitalization were not included in our study.

Statistical analysis

Continuous data with normal distribution are reported as mean \pm SD while continuous data with non-normal distribution are reported as median and ranges (minimum to maximum) or interquartile ranges (IQRs). Independent-samples *t*-test and Mann-Whitney *U* test were used for group comparisons for variables with normal and non-normal distribution, respectively. Categorical data are reported as counts and percentages. Group comparisons for categorical data were made with the χ^2 test.

All analysis began at the landmark time of discharge from the index hospitalization. Cumulative incidence function using Fine-Gray modeling was used to compare the incidence of first HE-related readmissions between the NSBB and no-NSBB groups while taking competing risk of death or liver transplant into account.

Univariate and multivariate Cox regression analyses were done to identify independent predictors of the first HE-related readmissions. Backward elimination technique with $P < 0.10$ for entering the model and $P < 0.05$ for staying in the model was utilized.

To include all of the HE-related admissions, we also determined independent predictors of HE-related hospitalizations per person-month. Due to the overdispersion and right-skewed distribution of this outcome variable, a negative binomial generalized regression model was employed. In this model, total person-months of follow-up was implemented as the offset variable and the follow-up period ended with death, liver transplantation, or end of the study period. The results for the negative binomial generalized regression model were reported as adjusted incidence rate ratios (IRRs) with 95% CIs that represent the relationship between HE-related admissions per person-month and a predictor while considering other covariates.

Subset analysis

To further explore the association of NSBB use and HE-related readmissions, multivariate Cox regression analysis and negative binomial generalized regression model were performed in different subgroups including NSBB *vs* SBB, ascites, EV, MELD-Na score, lactulose and rifaximin subgroups.

A P value < 0.05 was considered to be statistically significant. We performed the statistical analyses using SPSS 21 (SPSS Inc) and R statistical software, version 3.6.1 (R Foundation for Statistical Computing, Vienna, Austria). The study was approved by the institutional review board. The statistical review of the study was performed by a biomedical statistician Giovanna Saracino, PhD.

RESULTS

Baseline characteristics

There were 393 patients with a mean age of 58.1 ± 10.2 years and 144 (36.7%) had ascites. The median MELD-Na score was 22 (IQR: 11) and most patients fell into CTP class C (49.1%) or B (43.8%) (Table 1). A total of 143 patients (36.4%) were treated with NSBBs (nadolol, propranolol, or carvedilol) for either prevention of gastrointestinal bleeding or when used for cardiac indications. Nadolol was the most common NSBB used in this study (46.2%) followed by propranolol (43.4%) and carvedilol (10.5%) (Table 2). In the SBB comparison group, 39 patients were given either metoprolol (89.7%) or atenolol (10.3%).

Outcome and follow-up of all patients

The median follow-up time was 3.8 years (IQR: 4.1 years). There were 187 patients (47.6%) who had HE-

Table 1 Baseline characteristics of the patients on the first liver-related hospitalization

	Whole group (n = 393)	NSBB group (n = 143)	No-NSBB group (n = 250)	P value
Age, yr	58.1 ± 10.2	57.5 ± 8.8	58.4 ± 11.0	0.39 ²
Gender, male	229 (58.3)	88 (61.5)	141 (56.4)	0.32 ³
Heart rate, bpm	84 (44-161)	76 (44-122)	87 (50-161)	< 0.001 ⁴
Etiology of liver disease ¹				0.32 ³
Hepatitis C virus	129 (32.8)	55 (38.5)	74 (29.6)	
Alcoholic	124 (31.6)	38 (26.6)	86 (34.4)	
NASH	49 (12.5)	17 (11.9)	32 (12.8)	
Cryptogenic	51 (13.0)	21 (14.7)	30 (12.0)	
Other causes	56 (14.2)	23 (16.1)	33 (13.2)	
CTP score				0.054 ³
A	28 (7.1)	13 (9.1)	15 (6.0)	
B	172 (43.8)	71 (49.7)	101 (40.4)	
C	193 (49.1)	59 (41.3)	134 (53.6)	
MELD score	19 (6-40)	17 (6-39)	19 (6-40)	0.02 ⁴
MELD-Na score	22 (6-40)	20 (6-39)	22.5 (6-40)	0.005 ⁴
History of hepatocellular carcinoma	29 (7.4)	8 (5.6)	21 (8.4)	0.31 ³
History of esophageal varices	154 (39.2)	84 (58.7)	70 (28.0)	< 0.001 ³
History of TIPS	47 (12.0)	17 (11.9)	30 (12.0)	0.97 ³
Presence of HE	323 (82.2)	113 (79.0)	210 (84.0)	0.22 ³
Lactulose use	285 (72.5)	107 (74.8)	178 (71.2)	0.44 ³
Rifaximin use	208 (52.9)	81 (56.6)	127 (50.8)	0.26 ³
Presence of ascites	144 (36.7)	50 (35.0)	94 (37.8)	0.58 ³
International normalized ratio	1.5 (1-14)	1.4 (1-4)	1.6 (1-14)	0.003 ⁴
Platelet count, × 10 ³ /mm ³	84 (4-515)	72 (15-280)	95 (4-515)	0.002 ⁴
White cell count, × 10 ³ /mm ³	6.8 (0.2-51.9)	5.7 (1.3-43.7)	7.7 (0.2-51.9)	< 0.001 ⁴
Creatinine, mg/dL	1.3 (0.3-33.0)	1.4 (0.4-33.0)	1.3 (0.3-9.3)	0.59 ⁴
Total bilirubin, mg/dL	2.7 (0.2-137.0)	2.5 (0.2-43.0)	3.1 (0.3-137.0)	0.03 ⁴
Serum Albumin, g/dL	2.7 (1.0-5.0)	2.8 (2.0-5.0)	2.6 (1.0-5.0)	0.02 ⁴
Aspartate aminotransferase, U/L	59 (3-4048)	51 (8-677)	67 (3-4048)	< 0.001 ⁴
Alanine aminotransferase, U/L	39 (10-1180)	34 (13-538)	41 (10-1180)	0.02 ⁴

Data are presented as *n* (%), mean ± SD or median (range).

¹The cumulative percent exceeds 100% as some patients had multiple etiologies of liver disease.

²*P* calculated by Independent-samples *t*-test.

³*P* calculated by χ^2 test.

⁴*P* calculated by Mann-Whitney *U* test.

CTP: Child-Turcotte-Pugh; HE: Hepatic encephalopathy; NSBB: Nonselective beta-blocker; MELD: Model for End-Stage Liver Disease; Na: Sodium; NASH: Non-alcoholic steatohepatitis; TIPS: Transjugular intrahepatic portosystemic shunt.

related readmissions during the follow-up period. The median time between the first admission and future readmission was 1.9 mo (IQR:4.7 mo). Ninety-six patients (24.4%) died and 50 patients (12.7%) received a liver transplant during the study period. The leading causes of death was sepsis [*n* = 32 (33.3%)] followed by cirrhosis and its complications [*n* = 24 (25.0%)], respiratory failure [*n* = 17 (17.7%)] and multi-organ failure [*n* = 8 (8.33%)]. The remaining 15 (15.6%) patients died of other causes.

Table 2 Type, dose and number of patients on nonselective or selective beta-blockers

	Number of patients	Total daily dose (mg)
Nonselective beta-blocker (<i>n</i> = 143)		
Nadolol	66 (46.2)	20 (20-80)
Propranolol	62 (43.4)	30 (10-80)
Carvedilol	15 (10.5)	12.50 (6.25-50.00)
Selective beta-blocker (<i>n</i> = 39)		
Metoprolol	35 (89.7)	50.00 (12.50-200.00)
Atenolol	4 (10.3)	50 (25-100)

Data are presented as *n* (%) or median (range).

Outcomes according to NSBB therapy

The NSBB group had significantly lower heart rate, MELD-Na score, and platelet count compared with the group not receiving NSBB therapy (Table 1). Further, patients on NSBB therapy had significantly higher rates of presence of EV in comparison with patients not receiving treatment with NSBBs (58.7% *vs* 28%, respectively).

Ninety-one patients (63.6%) in the NSBB group and 96 patients (38.4%) in the no-NSBB group experienced HE-related readmission ($P < 0.001$). The cumulative incidence of the first HE-related readmissions within 5.5 years was significantly higher in patients taking NSBB compared with patients who were not prescribed NSBB (71.8% *vs* 41.8%, respectively; $P < 0.0001$) (Figure 1). The mean time to the first HE-related readmission was not significantly different between the two groups (5.3 mo *vs* 4.6 mo in NSBB and no-NSBB groups, respectively, $P = 0.5$). Furthermore, no significant difference was observed between the two groups regarding the mortality rate [$n = 32$ (22.4%) in the NSBB *vs* $n = 64$ (25.6%) in the no-NSBB groups, $P = 0.5$]. Stratifying the patients according to SBB use, 18 patients (46.2%) with SBB use and 169 patients (47.7%) without SBB use experienced HE-related readmission ($P = 0.9$).

Factors associated with the first HE-related readmissions

Results of the univariate analysis of factors associated with HE-related rehospitalization are shown in Table 3. After adjustment of demographic characteristics and surrogate markers of liver disease severity, NSBB use was the only independent predictor of the first HE-related readmissions [HR: 1.74 (95%CI: 1.29-2.34)]. This effect was not seen in patients taking SBBs. To further explore this, multivariate Cox regression model was employed in different subgroups of our patients including NSBB *vs* SBB, ascites, EV, MELD-Na score, lactulose, and rifaximin subgroups. NSBB use remained an independent predictor of the first HE-related readmissions in all of these subgroups (Supplementary Tables 1-9).

Factors associated with HE-related admissions per person-month

To further explore our findings, we examined the association of NSBB use with all HE-related admissions. NSBB use was an independent predictor of HE-related admissions per person-month [IRR: 1.50 (95%CI: 1.08-2.07)] alongside other variables including MELD-Na score, history of TIPS, lactulose use and platelet count (Table 4). Similar findings were present with liver-related admissions as the outcome variable. This effect was not observed in patients on SBBs. To further investigate this, negative binomial generalized regression model was performed in different subgroups of our patients including NSBB *vs* SBB, ascites, EV, MELD-Na score, lactulose and rifaximin subgroups. NSBB use remained an independent predictor of HE-related admissions per person-month in all of these subgroups (Supplementary Tables 10-18).

DISCUSSION

Our results demonstrate that patients treated with NSBBs experienced a significantly higher rate of the first HE-related readmissions compared to patients who did not receive NSBBs. Additionally, NSBB group patients had significantly higher cumulative incidence of the first HE-related readmissions compared to patients in the no-NSBB group. Finally, NSBB use was an independent predictor of HE-related admissions per person-month. These findings were persistent even after adjustment for markers of liver disease severity. NSBB use is associated with incident HE[16,17]; however, these data now extend prior research to show that NSBB use is associated with an increased burden of HE-related

Table 3 Univariate and multivariate Cox regression predicting the first hepatic encephalopathy-related readmission

Variable	Unadjusted		Adjusted	
	HR (95%CI)	P value	HR (95%CI)	P value
Age, yr	1.00 (0.99-1.02)	0.93		
Gender, male	1.03 (0.77-1.38)	0.82		
MELD-Na score (reference: MELD-Na score < 15)				
15 ≤ MELD-Na score ≤ 24	1.26 (0.88-1.79)	0.21		
25 ≤ MELD-Na score ≤ 34	0.89 (0.58-1.36)	0.60		
MELD-Na score > 34	0.38 (0.09-1.55)	0.18		
History of EV, presence of	1.24 (0.93-1.66)	0.15		
History of TIPS, presence of	1.34 (0.89-2.01)	0.16	1.48 (0.98-2.25)	0.065
NSBB use, presence of	1.81 (1.35-2.41)	< 0.001	1.74 (1.29-2.34)	< 0.001
SBB use, presence of	0.90 (0.55-1.46)	0.66		
Lactulose use, presence of	1.28 (0.89-1.82)	0.18		
Rifaximin use, presence of	0.88 (0.66-1.18)	0.40		
Ascites, presence of	1.10 (0.81-1.48)	0.55		
Platelet count, × 10 ³ /mm ³	0.996 (0.994-0.999)	0.008	0.997 (0.995-1.000)	0.07

EV: Esophageal varices; HR: Hazard ratio; MELD-Na: Model for End-Stage Liver Disease-Sodium; NSBB: Nonselective beta-blocker; SBB: Selective beta-blocker; TIPS: Transjugular intrahepatic portosystemic shunt.

Table 4 Negative binomial generalized regression model predicting hepatic encephalopathy-related admissions per person-month

Variable	IRR (95%CI)	P value	B value
Age, yr	0.99 (0.98-1.01)	0.19	-0.010
Gender, male	1.10 (0.82-1.48)	0.54	0.094
MELD-Na score	1.05 (1.03-1.08)	< 0.001	0.052
History of EV, presence of	0.98 (0.72-1.35)	0.92	-0.016
History of TIPS, presence of	1.93 (1.24-3.01)	0.003	0.660
NSBB use, presence of	1.50 (1.08-2.07)	0.015	0.403
SBB use, presence of	0.81 (0.50-1.31)	0.40	-0.208
Lactulose use, presence of	1.47 (1.00-2.15)	0.048	0.384
Rifaximin use, presence of	0.73 (0.53-1.00)	0.050	-0.319
Ascites, presence of	1.26 (0.93-1.72)	0.14	0.233
Platelet count, × 10 ³ /mm ³	0.997 (0.995-1.000)	0.03	-0.003

EV: Esophageal varices; IRR: Incidence rate ratio; MELD-Na: Model for End-Stage Liver Disease-Sodium; NSBB: Nonselective beta-blocker; SBB: Selective beta-blocker; TIPS: Transjugular intrahepatic portosystemic shunt.

readmissions.

Do NSBBs lead to HE?

There is controversy in the literature regarding the effect that NSBB use has on HE development. Prior studies in early-stage cirrhotic patients reported beneficial to no effect of NSBB on HE development. This includes a cohort of 28 responders to propranolol ± isosorbide mononitrate therapy[18], a randomized trial of propranolol in 20 CTP class A patients[19], and a cohort of 82 patients with cirrhosis [20]. Conversely, other reports indicate an increased risk of HE in NSBB users. In a prospective cohort study of 218 patients with cirrhosis, both NSBB/SBB use was independently associated with a higher




Figure 1 Cumulative incidence of death or liver transplant in cirrhotic patients with a hepatic encephalopathy-related admission that received nonselective beta-blockers vs those who did not (A) and cumulative incidence of the first hepatic encephalopathy-related readmissions in the same groups after adjusting for competing risk of death or liver transplant (B). HE: Hepatic encephalopathy; LT: Liver transplant; NSBB: Nonselective beta-blocker.

rate of minimal HE diagnosis[17]. In a population-based cohort study of 1979 cirrhotic patients, NSBB use was a significant risk factor for incident HE[16]. They hypothesized that NSBB use is a proxy for high-risk varices and severe portal hypertension. The possible explanations for these contradictory research reports of NSBB on HE development are small patient populations, different demographic data, severity of the liver disease, and duration of follow up.

Our data make two major contributions. First, we showed that adjusting for disease severity with granular patient-level data, NSBB use was clearly associated with recurrent HE. Second, our data raise the complexity of the discussion substantially. Our competing risk regression shows that NSBBs are associated with a slightly lower risk of death, confirming the benefits of NSBB use[21,22]. While longer survival avails patients of more opportunity for readmission, our data also shows that NSBBs are associated with a higher burden of readmissions per person-month. Taken together, these data clarify the tradeoffs of NSBB therapy. Therefore, physicians need to be vigilant about NSBB prescription and intensifying therapy.

Reconciling data

We showed that NSBB use in decompensated cirrhotic patients (CTP classes B and C) with high MELD-Na scores increases the risk of HE-related admissions during long-term follow-up. Although no clear explanation for the increase in HE-related admissions with NSBB use can be made, specific comments can be described. Krag *et al*[23] proposed the 'window hypothesis', a certain time frame during the natural course of cirrhosis that only within which NSBB use has a beneficial effect on mortality. The same concept can be true regarding the effect of NSBBs on HE development. In cirrhotic patients with mild to moderate portal hypertension, NSBBs counteract the hyperdynamic cardiovascular state and decrease portal hypertension[23,24]. This likely results in a beneficial effect on HE as it counteracts the most likely pathophysiologic mechanism of HE development; shunting of ammonia towards the brain. This can also be the explanation for the slightly increased survival of the patients taking NSBBs in our study.

With cirrhosis progression, patients develop severe portal hypertension that results in increased cardiac output and decreased systemic vascular resistance. NSBB administration in this stage compromises the systemic perfusion pressure that can ultimately decrease hepatic perfusion[13,23-25]. This will result in increased blood ammonia level shunting systemically to the brain in the context of severe portal hypertension, resulting in the development of HE and increased HE-related readmissions over time.

Our study has certain limitations. Although retrospective in nature, having a relatively large sample size with long-term follow up mitigates the study design. We did not know whether any of the patients had a liver-related hospitalization or a previous HE episode before inclusion in the study. However, adjusting for disease severity did not change the results. Although the exact start and end dates and

compliance with NSBB and SBB use in all cases and hence the association between duration of NSBB use and risk of HE-related readmissions were not determined, stable estimates across a variety of subsets showed similar results. Furthermore, blood pressure, ammonia levels, precipitant factors for HE, association of NSBB use with different overt HE grades, indications/contraindications for NSBB use and diuretic/proton pump inhibitor use were not explored in our study. Due to limited sample size for an individual NSBB medication and lack of a universal dose-conversion guideline for different NSBBs, a meaningful analysis exploring the association of different doses of NSBBs and risk of HE-related readmissions could not be performed in our study. Although HVPG measurements would have been useful, obtaining HVPG data in a retrospective fashion on our patients was impractical. However, future trials could help delineate the exact HVPG level at which HE readmissions occur in relation to heart rate and NSBB use. Therefore, we believe that our study provides a foundation to guide future prospective trials, allowing pharmacologic comparisons to further delineate the association between NSBB, SBB and HE-related readmissions.

CONCLUSION

In conclusion, we show that NSBB use is independently associated with increased HE-related readmissions in patients with cirrhosis, regardless of liver disease severity or biochemical abnormalities. Further prospective studies are needed to determine the impact of NSBBs on HE and other portal hypertension complications.

ARTICLE HIGHLIGHTS

Research background

Hepatic encephalopathy (HE) is a cirrhosis complication leading to frequent hospitalizations and imposes a significant economic burden on the healthcare system. Nonselective beta-blockers (NSBBs) are the mainstay of pharmacologic treatment for portal hypertension and in the prevention of variceal bleeding in cirrhosis. The role of NSBBs in the development of HE-related complications is not known.

Research motivation

We hypothesized that since NSBBs decrease cardiac output and portal flow, the decreased metabolic filtering process of liver parenchyma may lead to increased HE-related hospitalizations. If there is a signal that NSBB use is associated with HE-related hospitalizations, further multicenter trials are warranted to explore the impact of NSBBs on HE and other portal hypertension complications.

Research objectives

The main objective of this study was to evaluate the impact of NSBB administration on HE-related readmissions in cirrhotic patients.

Research methods

We performed an observational, retrospective, single-center cohort study including 393 patients with cirrhosis admitted to Baylor University Medical Center for liver-related portal hypertension indications between January 2013 and July 2018. Independent predictors of the first HE-related readmissions were identified using Cox proportional hazards analysis. The cumulative incidence of the first HE-related readmissions between patients receiving NSBBs and not receiving NSBBs was examined using Fine-Gray modeling to account for the competing risk of death or liver transplantation.

Research results

In a cohort of patient with mostly Child class C (49.1%) or B (43.8%) cirrhosis, the cumulative incidence of the first HE-related readmissions was significantly higher in patients taking NSBBs compared to patients not receiving NSBBs (71.8% *vs* 41.8%, $P < 0.0001$). In multivariate analysis, after adjusting for demographics, markers of liver disease severity, selective beta-blocker, lactulose and rifaximin use, NSBB use [Hazard ratio: 1.74 (95%CI: 1.29-2.34)] was independently associated with the first HE-related readmissions over a median follow-up of 3.8 years. These results warrant further multicenter clinical trials to explore the impact of NSBBs on HE and other portal hypertension complications.

Research conclusions

NSBB use in patients with advanced cirrhosis is independently associated with increased HE-related readmissions, regardless of liver disease severity or biochemical abnormalities. This can be due to the role of NSBB use in decreasing the systemic perfusion pressure that can ultimately lead to a decrease in hepatic perfusion in advanced cirrhosis that will result in an increased blood ammonia level shunting

systemically to the brain.

Research perspectives

As this study was a retrospective study, future prospective cohort and randomized clinical trials are warranted to explore the impact of NSBBs on HE and other portal hypertension complications.

ACKNOWLEDGEMENTS

We thank Mr. Daniel Bizzarri for his help with data gathering.

FOOTNOTES

Author contributions: Fallahzadeh MA, Asrani SK and Rahimi RS designed the research study; Fallahzadeh MA, Asrani SK, Tapper EB, Saracino G and Rahimi RS performed the acquisition, analysis and interpretation of the data; Fallahzadeh MA, Asrani SK and Rahimi RS drafted the manuscript and all authors contributed to revising the manuscript critically; all authors have read and approved the final manuscript.

Institutional review board statement: The study was reviewed and approved for publication by our Institutional Reviewer.

Informed consent statement: According to the IRB protocol and policy, all participants of the study provided informed consent indirectly about personal and medical data collection prior to study enrollment.

Conflict-of-interest statement: All the authors report no relevant conflicts of interest for this article.

Data sharing statement: Technical appendix, statistical code, and dataset available from the corresponding author at aminfa91@gmail.com. Consent was not obtained but the presented data are anonymized and risk of identification is low. No additional data are available.

STROBE statement: The authors have read the STROBE Statement—checklist of items, and the manuscript was prepared and revised according to the STROBE Statement—checklist of items.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Country/Territory of origin: United States

ORCID number: Mohammad Amin Fallahzadeh 0000-0002-4308-9856; Sumeet K Asrani 0000-0001-9174-5670; Elliot B Tapper 0000-0002-0839-1515; Giovanna Saracino 0000-0001-9323-5623; Robert S Rahimi 0000-0002-2595-1852.

Corresponding Author's Membership in Professional Societies: American Association for the Study of Liver Diseases, No. 224919.

S-Editor: Gong ZM

L-Editor: A

P-Editor: Gong ZM

REFERENCES

- 1 Ferenci P. Hepatic encephalopathy. *Gastroenterol Rep (Oxf)* 2017; **5**: 138-147 [PMID: 28533911 DOI: 10.1093/gastro/gox013]
- 2 Fallahzadeh MA, Rahimi RS. Hepatic Encephalopathy and Nutrition Influences: A Narrative Review. *Nutr Clin Pract* 2020; **35**: 36-48 [PMID: 31872484 DOI: 10.1002/ncp.10458]
- 3 Stepanova M, Mishra A, Venkatesan C, Younossi ZM. In-hospital mortality and economic burden associated with hepatic encephalopathy in the United States from 2005 to 2009. *Clin Gastroenterol Hepatol* 2012; **10**: 1034-41.e1 [PMID: 22642955 DOI: 10.1016/j.cgh.2012.05.016]
- 4 Alsahhar JS, Rahimi RS. Updates on the pathophysiology and therapeutic targets for hepatic encephalopathy. *Curr Opin Gastroenterol* 2019; **35**: 145-154 [PMID: 30893082 DOI: 10.1097/MOG.0000000000000527]
- 5 Bajaj JS, Reddy KR, Tandon P, Wong F, Kamath PS, Garcia-Tsao G, Maliakkal B, Biggins SW, Thuluvath PJ, Fallon MB,

- Subramanian RM, Vargas H, Thacker LR, O'Leary JG; North American Consortium for the Study of End-Stage Liver Disease. The 3-month readmission rate remains unacceptably high in a large North American cohort of patients with cirrhosis. *Hepatology* 2016; **64**: 200-208 [PMID: 26690389 DOI: 10.1002/hep.28414]
- 6 **Hirode G**, Vittinghoff E, Wong RJ. Increasing Burden of Hepatic Encephalopathy Among Hospitalized Adults: An Analysis of the 2010-2014 National Inpatient Sample. *Dig Dis Sci* 2019; **64**: 1448-1457 [PMID: 30863953 DOI: 10.1007/s10620-019-05576-9]
- 7 **Tapper EB**, Parikh ND, Sengupta N, Mellinger J, Ratz D, Lok AS, Su GL. A risk score to predict the development of hepatic encephalopathy in a population-based cohort of patients with cirrhosis. *Hepatology* 2018; **68**: 1498-1507 [PMID: 29091289 DOI: 10.1002/hep.29628]
- 8 **Qi XS**, Bai M, Fan DM. Nonselective β -blockers may induce development of portal vein thrombosis in cirrhosis. *World J Gastroenterol* 2014; **20**: 11463-11466 [PMID: 25170238 DOI: 10.3748/wjg.v20.i32.11463]
- 9 **Giannelli V**, Lattanzi B, Thalheimer U, Merli M. Beta-blockers in liver cirrhosis. *Ann Gastroenterol* 2014; **27**: 20-26 [PMID: 24714633]
- 10 **Kim SG**, Larson JJ, Lee JS, Therneau TM, Kim WR. Beneficial and harmful effects of nonselective beta blockade on acute kidney injury in liver transplant candidates. *Liver Transpl* 2017; **23**: 733-740 [PMID: 28187503 DOI: 10.1002/lt.24744]
- 11 **Mandorfer M**, Bota S, Schwabl P, Bucsics T, Pfisterer N, Kruzik M, Hagmann M, Blacky A, Ferlitsch A, Sieghart W, Trauner M, Peck-Radosavljevic M, Reiberger T. Nonselective β blockers increase risk for hepatorenal syndrome and death in patients with cirrhosis and spontaneous bacterial peritonitis. *Gastroenterology* 2014; **146**: 1680-90.e1 [PMID: 24631577 DOI: 10.1053/j.gastro.2014.03.005]
- 12 **Sersté T**, Melot C, Francoz C, Durand F, Rautou PE, Valla D, Moreau R, Lebre C. Deleterious effects of beta-blockers on survival in patients with cirrhosis and refractory ascites. *Hepatology* 2010; **52**: 1017-1022 [PMID: 20583214 DOI: 10.1002/hep.23775]
- 13 **Téllez L**, Ibáñez-Samaniego L, Pérez Del Villar C, Yotti R, Martínez J, Carrión L, Rodríguez de Santiago E, Rivera M, González-Mansilla A, Pastor Ó, Bermejo J, Bañares R, Albillos A. Non-selective beta-blockers impair global circulatory homeostasis and renal function in cirrhotic patients with refractory ascites. *J Hepatol* 2020; **73**: 1404-1414 [PMID: 32446716 DOI: 10.1016/j.jhep.2020.05.011]
- 14 **Weissenborn K**. Hepatic Encephalopathy: Definition, Clinical Grading and Diagnostic Principles. *Drugs* 2019; **79**: 5-9 [PMID: 30706420 DOI: 10.1007/s40265-018-1018-z]
- 15 **Bajaj JS**, Lauridsen M, Tapper EB, Duarte-Rojo A, Rahimi RS, Tandon P, Shawcross DL, Thabut D, Dhiman RK, Romero-Gomez M, Sharma BC, Montagnese S. Important Unresolved Questions in the Management of Hepatic Encephalopathy: An ISHEN Consensus. *Am J Gastroenterol* 2020; **115**: 989-1002 [PMID: 32618647 DOI: 10.14309/ajg.0000000000000603]
- 16 **Tapper EB**, Parikh ND. Mortality due to cirrhosis and liver cancer in the United States, 1999-2016: observational study. *BMJ* 2018; **362**: k2817 [PMID: 30021785 DOI: 10.1136/bmj.k2817]
- 17 **Acharya C**, Thacker L, Fagan A, White MB, Sterling RK, Stravitz RT, Sanyal AJ, Puri P, Heuman D, Fuchs M, John B, Lee H, Matherly S, Siddiqui MS, Bajaj JS. 422 - Beta Blockers Use in Cirrhosis is Independently Associated with Minimal Hepatic Encephalopathy. *Gastroenterology* 2018; **154**: S-1089 [DOI: 10.1016/s0016-5085(18)33627-8]
- 18 **Abrales JG**, Tarantino I, Turnes J, Garcia-Pagan JC, Rodés J, Bosch J. Hemodynamic response to pharmacological treatment of portal hypertension and long-term prognosis of cirrhosis. *Hepatology* 2003; **37**: 902-908 [PMID: 12668985 DOI: 10.1053/jhep.2003.50133]
- 19 **Dunk AA**, Moore J, Symon A, Dickie A, Sinclair TS, Mowat NA, Brunt PW. The effects of propranolol on hepatic encephalopathy in patients with cirrhosis and portal hypertension. *Aliment Pharmacol Ther* 1988; **2**: 143-151 [PMID: 2979240 DOI: 10.1111/j.1365-2036.1988.tb00681.x]
- 20 **Rajini K**, Solomon TR, Arumugham A, Ananthavadivelu M, Rangachari B, Kalyanasundaram M, Govindarajan R, Vaishnavi Priya C, Kavitha S. Influence of Beta Blockers on Complications of Cirrhosis. *J Clin Exp Hepatol* 2016; **6**: S52-S53 [DOI: 10.1016/j.jceh.2016.06.102]
- 21 **Gluud LL**, Krag A. Banding ligation versus beta-blockers for primary prevention in oesophageal varices in adults. *Cochrane Database Syst Rev* 2012; CD004544 [PMID: 22895942 DOI: 10.1002/14651858.CD004544.pub2]
- 22 **Rodrigues SG**, Mendoza YP, Bosch J. Beta-blockers in cirrhosis: Evidence-based indications and limitations. *JHEP Rep* 2020; **2**: 100063 [PMID: 32039404 DOI: 10.1016/j.jhepr.2019.12.001]
- 23 **Krag A**, Wiest R, Albillos A, Gluud LL. The window hypothesis: haemodynamic and non-haemodynamic effects of β -blockers improve survival of patients with cirrhosis during a window in the disease. *Gut* 2012; **61**: 967-969 [PMID: 22234982 DOI: 10.1136/gutjnl-2011-301348]
- 24 **Kockerling D**, Nathwani R, Forlano R, Manousou P, Mullish BH, Dhar A. Current and future pharmacological therapies for managing cirrhosis and its complications. *World J Gastroenterol* 2019; **25**: 888-908 [PMID: 30833797 DOI: 10.3748/wjg.v25.i8.888]
- 25 **Iwakiri Y**. The molecules: mechanisms of arterial vasodilatation observed in the splanchnic and systemic circulation in portal hypertension. *J Clin Gastroenterol* 2007; **41** Suppl 3: S288-S294 [PMID: 17975478 DOI: 10.1097/MCG.0b013e3181468b4c]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

Telephone: +1-925-3991568

E-mail: bpgoffice@wjgnet.com

Help Desk: <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

