

16

Name of Journal: *World Journal of Gastrointestinal Oncology***ESPS Manuscript NO:** 21759**Manuscript Type:** REVIEW

5

Chromodomain-Helicase-DNA Binding Protein 5, Chromodomain-Helicase-DNA Binding Protein 7, and Pronecrotic Mixed Lineage Kinase Domain- Like Protein Serve as Potential Prognostic Biomarkers in Patients With Resected Pancreatic Adenocarcinomas

Crystal S. Seldon, Lauren E. Colbert, William A. Hall, Sarah B. Fisher, David S. Yu, Jerome C. Landry

Match Overview

1

CrossCheck 188 words
Colbert, Lauren E., Sarah B. Fisher, Claire W. Hardy, William A. Hall, Burcu Saka, Joseph W. Shelton, Aleksandra

3%

2

CrossCheck 137 words
Hall, W A, A V Petrova, L E Colbert, C W Hardy, S B Fisher, B Saka, J W Shelton, M D Warren, B G Pantazides, K G

2%

3

Internet 48 words
crawled on 20-Sep-2014
effilop.com

1%

4

CrossCheck 47 words
Zhao, Rui, Fanyi Meng, Nisha Wang, Wenli Ma, and Qiao Yan. "Silencing of CHD5 Gene by Promoter Methylation

1%

5

Internet 47 words
crawled on 09-Apr-2014
zfin.org

1%

6

Internet 46 words
crawled on 09-Sep-2015
www.science.gov

1%

7

Internet 24 words
crawled on 17-Sep-2009

<1%

Chromodomain-Helicase-DNA Binding Protein 5, Chromodomain-Helicase-DNA Binding Protein 7, and Pro Necrotic Mixed Lineage Kinase Domain-Like Protein Serve as Potential Prognostic Biomarkers in Patients With Resected Pancreatic Adenocarcinomas

Did you mean: Chromodomain-Helicase-DNA Binding Protein 5, Chromodomain-Helicase-DNA Binding Protein 7, and *Pro Necrotic Mixed Lineage Kinase Domain-Like Protein* Serve as Potential Prognostic Biomarkers in Patients With Resected Pancreatic Adenocarcinomas

Low CHD5 expression activates the DNA damage response and predicts poor outcome in patients undergoing adjuvant therapy for resected pancreatic cancer

WA Hall, AV Petrova, LE Colber, CW Hardy, SB Fisher... - *Oncogene*, 2013 - nature.com

... resulting from activated oncogenes and inactivated tumor suppressor proteins induces activation of ... 12 CHD5 binds to histone H3 for its tumor suppressor function. ... Decreased expression of chromodomain helicase DNA-binding protein 5 is an unfavorable prognostic marker in ...

Cited by 5 Related articles All 4 versions Cite Save

CHD7 expression predicts survival outcomes in patients with resected pancreatic cancer

LE Colber, AV Petrova, SB Fisher, BG Pantazides... - *Cancer research*, 2014 - AACR

... Chromodomain helicase DNA binding protein 7 (CHD7) is a member of a family of chromodomain enzymes that belong to the ... by centrifugation at 16 × g for 10 minutes and 150 µg of protein for each ... with 0.5% Triton X-100 (Fisher Scientific), and blocked in a 5% bovine serum ...

Cited by 3 Related articles All 7 versions Cite Save

Create alert

Did you mean to search for: Chromodomain-Helicase-DNA Binding Protein 5, Chromodomain-Helicase-DNA Binding Protein 7, and *Pro Necrotic Mixed Lineage Kinase Domain-Like Protein* Serve as Potential Prognostic Biomarkers in Patients With Resected Pancreatic Adenocarcinomas

About Google Scholar Privacy Terms Provide feedback

全部 图片 新闻 视频 更多 ▾ 搜索工具

找到约 26 条结果 (用时 0.84 秒)

您是不是要找: [Chromodomain-helicase-DNA binding protein 5, chromodomain-helicase-DNA binding protein 7 and pro](#)
[necrotic mixed lineage kinase domain - like protein serve as potential prognostic biomarkers in patients with resected pancreatic adenocarcinomas](#)

[CHD7 Expression Predicts Survival Outcomes in Patients ...](#)

www.ncbi.nlm.nih.gov > ... > Literature > PubMed Central (PMC) - 翻译此页

2014年3月13日 - Pancreatic adenocarcinoma (PAC) has a poor prognosis with a five-year overall survival ... agent used to treat patients with PAC in the adjuvant setting (2, 5).

... Chromodomain helicase DNA binding protein 7 (CHD7), is a member of

Pronecrotic mixed lineage kinase domain-like protein expression is a ...

[Oncogene - Low CHD5 expression activates the DNA ...](#)

www.nature.com > Journal home > Archive > Original Articles - 翻译此页

2013年11月25日 - On Kaplan-Meier log-rank survival analysis, patients with low CHD5

... CHD5; DNA damage response; biomarker; pancreatic cancer; tumor ...

Chromodomain helicase DNA-binding protein 5 (CHD5) is a Pronecrotic mixed lineage kinase domain-like protein expression is Clinical Practice & Research.

[Aleksandra Petrova - Publications - ResearchGate](#)

www.researchgate.net/profile/Aleksandra.../publications ▾ 翻译此页

Article: Pronecrotic mixed lineage kinase domain-like protein expression is a prognostic biomarker in patients with early-stage resected pancreatic adenocarcinoma ... pilot clinical trial of chromodomain-helicase-DNA-binding protein 7 (CHD7) expression as a prognostic and predictive biomarker in patients with early-stage ...