

找到约 1,510 条结果 (用时 0.62 秒)

Minimally Invasive Ivor-Lewis Esophagectomy (MIILE): A Single ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5549044/> - 翻译此页

作者: J Wang - 2017 - 相关文章

2016年7月12日 - Since the introduction of minimally invasive esophagectomy, which was reported by However, the difference in blood loss, postoperative first day drain volume, ... injury of recurrent nerve, delayed gastric emptying, incisional hernia, and ... Some factors that influence these results include the following: (1) ...

Outcomes After Minimally Invasive Esophagectomy - NCBI - NIH

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4103614/> - 翻译此页

作者: JD Luketich - 2012 - 被引用次数: 508 - 相关文章

Outcomes after diagnosis of esophageal cancer are suboptimal, with a ... modality —esophagectomy with lesser morbidity for both early-stage disease and ... Schematic representation of the construction of minimally invasive Ivor Lewis anastomosis. in 200 consecutive patients: factors influencing postoperative outcomes.

缺少字词: delayed dge miile

Factors influencing delayed gastric emptying after pylorus-preserving ...

<https://www.ncbi.nlm.nih.gov/pubmed/12788421> - 翻译此页

作者: YC Park - 2003 - 被引用次数: 165 - 相关文章

BACKGROUND: Delayed gastric emptying (DGE) is the most frequent postoperative ... to determine which factors influence the development of DGE after PPPD. ... affect gastric emptying after PPPD during the early postoperative period.

缺少字词: minimally invasive ivor lewis esophagectomy miile

ivor lewis esophagectomy: Topics by Science.gov

<https://www.science.gov/topicpages/i/ivor+lewis+esophagectomy.html> - 翻译此页

No significant influence of the D₅₀ on the proximal esophagus was noted on the ... OBJECTIVES Ivor Lewis minimally invasive oesophagectomy (ILMIE) is a complex Methods Patients undergoing Ivor-Lewis esophagectomy for early stage Gastric emptying was significantly delayed after esophagectomy, and ...

postoperative gastrointestinal anastomotic: Topics by Science.gov

www.science.gov/topicpages/p/postoperative+gastrointestinal+anastomotic.html

Early detection of gastrointestinal leakage is important for successful treatment. ... Morbidity and mortality after esophagectomy are often related to anastomotic delayed gastric emptying, gastric outlet obstruction, and gastrogastic fistula. mobilization with cervical anastomosis and the post-operative procedure.

Name of Journal: *World Journal of Clinical Cases*

Manuscript NO: 44646

Manuscript type: ORIGINAL ARTICLE

Observational Study

Influencing factors of postoperative early delayed gastric emptying after minimally invasive Ivor-Lewis esophagectomy

Huang L *et al.* Factors of DGE after MIILE

Lei Huang, Jian-Qiang Wu, Bing Han, Zhi Wen, Pei-Rui Chen, Xiao-Kang Sun, Xiang-Dong Guo, Chang-Ming Zhao

Match Overview

4 matches

1 **Crossref** 40 words
"Abstracts of the 26th Annual Conference of APASL, February 15–19, 2017, Shanghai, China", *Hepatology International*

2%

[全部](#)[图片](#)[新闻](#)[视频](#)[购物](#)[更多](#)[设置](#)[工具](#)

找到约 4,780 条结果 (用时 0.45 秒)

Google 学术 : Influencing factors of postoperative early delayed gastric emptying after minimally invasive Ivor-Lewis esophagectomyMinimally invasive esophagectomy: lessons learned ... - **Nguyen** - 被引用次数 : 184... anastomotic leaks complicating esophagectomy: a ... - **Urschel** - 被引用次数 : 536... , and survival in patients undergoing Ivor Lewis ... - **Karl** - 被引用次数 : 326**The dynamic of nasogastric decompression after esophagectomy and ...**<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4756234/> ▾ [翻译此页](#)作者 : Y Zhao - 2016 - [相关文章](#)

2015年10月9日 - To investigate the regularity and the influence factors of nasogastric ... The nasogastric decompression of the patients with delayed gastric ... after gastric or esophageal surgery, especially after minimally invasive (MIE) approach (8,9). ... McKeown in 11 patients (3 MIE, 5 open, and 3 hybrid), Ivor-Lewis in 70 ...

Post-discharge complications after esophagectomy account for high ...<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4893471/> ▾ [翻译此页](#)作者 : SY Chen - 2016 - 被引用次数 : 8 - [相关文章](#)

Moreover, readmission rates after esophagectomy range from 5%-25%[8,9], and the ... Given that approximately a third of post-operative surgical complications ... their associated risk factors, and their influence on early hospital outcomes ... the chest was not entered and the stomach was used as a conduit), Ivor-Lewis (if the ...

[PDF]

Functional results of the Ivor Lewis Esophagectomy - Ghent University ...https://lib.ugent.be/fulltxt/.../002/.../RUG01-002163909_2014_0001_AC.pdf ... ▾ [翻译此页](#)作者 : A DELDYCKE - [相关文章](#)

Table 6: Logistic regression for delayed gastric emptying 3. Abstract (Engels). Introduction: Little is known on functional outcome after Ivor Lewis esophagectomy (ILE) ... influence of certain clinical factors on the occurrence of functional disorders and their impact search had to be specified by 'NOT minimally invasive'.

Preoperative endoscopic pyloric balloon dilatation decreases the rate ...https://www.researchgate.net/.../329351187_Preoperative_endoscopic_pylori... - [翻译此页](#)

2018年12月4日 - Delayed gastric emptying (DGE) after Ivor-Lewis esophagectomy ... esophageal tumors or logistic reasons (NDG: non-pyloric dilatation ... can prevent the need for early postoperative

找到约 7,070 条结果 (用时 0.46 秒)

Google 学术 : Influencing factors of postoperative early delayed gastric emptying after minimally invasive Ivor-Lewis esophagectomy

Minimally invasive esophagectomy: lessons learned ... - Nguyen - 被引用次数 : 184

... anastomotic leaks complicating esophagectomy: a ... - Urschel - 被引用次数 : 536

..., and survival in patients undergoing Ivor Lewis ... - Karl - 被引用次数 : 326

The dynamic of nasogastric decompression after esophagectomy and ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4756234/> ▼ [翻译此页](#)

作者 : Y Zhao - 2016 - [相关文章](#)

2015年10月9日 - To investigate the regularity and the influence factors of nasogastric ... The nasogastric decompression of the patients with delayed gastric ... after gastric or esophageal surgery, especially after minimally invasive (MIE) approach (8,9). ... McKeown in 11 patients (3 MIE, 5 open, and 3 hybrid), Ivor-Lewis in 70 ...

Minimally Invasive Ivor-Lewis Esophagectomy (MIILE): A Single ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5549044/> - [翻译此页](#)

作者 : J Wang - 2017 - [相关文章](#)

2016年7月12日 - Since the introduction of minimally invasive esophagectomy, which was reported by However, the difference in blood loss, postoperative first day drain volume, ... injury of recurrent nerve, delayed gastric emptying, incisional hernia, and ... Some factors that influence these results include the following: (1) ...

[PDF]

Functional results of the Ivor Lewis Esophagectomy - Ghent University ...

https://lib.ugent.be/fulltxt/.../002/.../RUG01-002163909_2014_0001_AC.pdf ... ▼ [翻译此页](#)

作者 : A DELDYCKE - [相关文章](#)

Table 6: Logistic regression for delayed gastric emptying ... 3. Abstract (Engels). Introduction: Little is known on functional outcome after Ivor Lewis esophagectomy (ILE) ... influence of certain clinical factors on the occurrence of functional disorders and their impact search had to be specified by 'NOT minimally invasive'.

(PDF) Recent Advances in Minimally Invasive Esophagectomy

https://www.researchgate.net/.../323561005_Recent_Advances_in_Minimall... - [翻译此页](#)

2018年8月1日 - several factors: patient selection, versatility in ... vention in the treatment of postoperative com- ... Key words: esophageal cancer, minimally invasive esophagectomy, open ... by