

Match Overview

1	CrossCheck 118 words Lohsiriwat, V., and D. Lohsiriwat. "Atraumatic O-ring wound retractor reduces postoperative pain". Techniques in Colo...	4%
2	CrossCheck 58 words Solomon, D.H.. "Determinants of selective cyclooxygenase -2 inhibitor prescribing: are patient or physician characte ...	2%
3	Internet 48 words crawled on 22-Jan-2016 www.wjgnet.com	2%
4	CrossCheck 42 words Lohsiriwat, Varut. "Enhanced recovery after surgery vs c... nventional care in emergency colorectal surgery". World J	1%
5	Internet 22 words crawled on 16-Sep-2015 numonthly.com	1%
6	CrossCheck 22 words Vather, Ryash, and Ian P. Bissett. "Risk factors for the dev elopment of prolonged post-operative ileus following elect...	1%

18

Name of Journal: *World Journal of Gastrointestinal Oncology*

ESPS Manuscript NO: 25081

Manuscript Type: ORIGINAL ARTICLE

Retrospective Cohort Study

Opioid-sparing ¹⁴ effect of selective cyclooxygenase-2 inhibitors on ¹⁵ surgical outcomes after open colorectal surgery within an enhanced recovery after surgery protocol

Varut Lohsiriwat

Abstract

AIM: To evaluate the opioid-sparing ¹⁴ effect of selective cyclooxygenase-2 (COX-2) inhibitors on short-term surgical outcomes after open colorectal surgery.

METHODS: Patients undergoing open colorectal resection within an enhanced recovery after surgery (ERAS) protocol from 2011 to 2015 were reviewed. Patients with combined general anesthesia and epidural anesthesia, and those with acute colonic obstruction or perforation were excluded. Patients receiving selective COX-2 inhibitor were compared with well-matched individuals without such a drug. Outcome measures included numeric pain score and morphine milligram equivalent (MME) consumption on postoperative day (POD) 1-3, gastrointestinal recovery (time to tolerate solid diet and time to defecate), complications and length of postoperative

学术搜索

找到约 80 条结果 (用时0.10秒)

文章

我的图书馆

时间不限

2016以来

2015以来

2012以来

自定义范围...

按相关性排序

按日期排序

搜索所有网页

中文网页

简体中文网页

包括专利

包含引用

创建快讯

小提示: 只搜索中文(简体)结果, 可在 学术搜索设置 指定搜索语言

Complications of the COX-2 inhibitors parecoxib and valdecoxib after cardiac surgery

NA Nussmeier, AA Whelton, MT Brown... - ... England Journal of ..., 2005 - Mass Medical Soc
... Although a hint of this adverse effect was noted in an earlier trial of parecoxib and valdecoxib
in ... Parecoxib sodium, a parenteral cyclooxygenase 2 selective inhibitor, improves morphine
analgesia and is opioid-sparing following total hip arthroplasty. ... (2015) Effects of sexually ...
被引用次数: 1019 相关文章 所有 19 个版本 引用 保存

researchgate.net 中的 [PDF]

Fast-track protocols in colorectal surgery

CL Donohoe, M Nguyen, J Cook, SG Murray, N Chen... - the surgeon, 2011 - Elsevier
... critics site bradycardia and hypotension as two possible complications resulting from inhibition
of sympathetic ... angiotensin-aldosterone system and anti-diuretic hormone, with the net effect of
fluid ... 48 However over-hydration may have adverse effects on physiological function. ...
被引用次数: 61 相关文章 所有 6 个版本 引用 保存

researchgate.net 中的 [PDF]

The role of the anesthesiologist in fast-track surgery; from multimodal analgesia to perioperative medical care

..., F Carli, Fast-Track Surgery Study Group - Anesthesia & ..., 2007 - journals.lww.com
BACKGROUND: Improving perioperative efficiency and
throughput has become increasingly important in t.
被引用次数: 369 相关文章 所有 9 个版本 引用 保存

researchgate.net 中的 [PDF]

Risk of anastomotic leakage with non - steroidal anti - inflammatory drugs in colorectal surgery

KJ Gorissen, D Benning, T Berghmans... - ... Journal of Surgery, 2012 - Wiley Online Library
... Addition of NSAIDs to morphine is estimated to have an opioid-sparing effect of 30 per cent, with ...
As this study was not designed to investigate the mode of action of NSAIDs, the ... Effect of diclofenac
on cyclooxygenase-2 levels and early breaking strength of experimental colonic ...
被引用次数: 74 相关文章 所有 6 个版本 引用 保存

researchgate.net 中的 [PDF]

[HTML] A comprehensive review of evidence-based strategies to prevent and treat postoperative ileus

SK Story, RS Chamberlain - Digestive surgery, 2009 - karger.com
... study with 40 patients found evidence that COX-2 inhibitors had opioid-sparing effects and
reduced ... apparent effectiveness of these fast-track approaches, and the negative effect of more

karger.com 中的 [HTML]

[全部](#) [新闻](#) [图片](#) [视频](#) [更多 ▾](#) [搜索工具](#)

找到约 3,390 条结果 (用时 0.67 秒)

[Opioid-sparing effect of selective cyclooxygenase-2 ...](#)

www.clinicalnutritionespen.com/article/S2405...2/abstract ▾ [翻译此页](#)

Opioid-sparing effect of selective cyclooxygenase-2 inhibitors on surgical outcomes after open colorectal surgery within an enhanced recovery after surgery ...

[Fast-track surgery: Toward comprehensive peri-operative care](#)

www.ncbi.nlm.nih.gov > ... > [PubMed Central \(PMC\)](#) ▾ [翻译此页](#)

作者: AJ Nanavati - 2014 - 被引用次数: 7 - [相关文章](#)

The concept of "fast-track surgery" also known as "enhanced recovery after ..." such a protocol targets the patient in each phase before, during, and after surgery [Figure 1]. ... (COX-2) inhibitors have shown to have opioid sparing effects when used ... to reduce post-operative ileus and vomiting after use in colorectal surgery.

[Postoperative Pain Control](#)

www.ncbi.nlm.nih.gov > ... > [PubMed Central \(PMC\)](#) ▾ [翻译此页](#)

作者: V Garimella - 2013 - 被引用次数: 7 - [相关文章](#)

Although traditionally the mainstay of postoperative analgesia is opioid based, ... Enhanced recovery protocols to reduce length of stay in colorectal surgery are ... acute pain is defined as pain present in a surgical patient after a procedure. ... or COX-2 inhibitors are given in addition to PCA morphine after surgery with a ...

[\[PDF\] Enhanced Recovery after Surgery Pain Management ...](#)



全部 新闻 图片 视频 更多 ▾ 搜索工具

找到约 3,280 条结果 (用时 0.74 秒)

[Opioid-sparing effect of selective cyclooxygenase-2 ...](#)

www.clinicalnutritionespen.com/article/S2405...2/abstract ▾ 翻译此页

Opioid-sparing effect of selective cyclooxygenase-2 inhibitors on surgical outcomes after open colorectal surgery within an enhanced recovery after surgery ...

[Fast-track surgery: Toward comprehensive peri-operative care](#)

www.ncbi.nlm.nih.gov > ... > PubMed Central (PMC) ▾ 翻译此页

作者: AJ Nanavati - 2014 - 被引用次数: 7 - 相关文章

The concept of "fast-track surgery" also known as "enhanced recovery after ... such a protocol targets the patient in each phase before, during, and after surgery [Figure 1]. (COX-2) inhibitors have shown to have opioid sparing effects when used ... to reduce post-operative ileus and vomiting after use in colorectal surgery.

[Postoperative Pain Control](#)

www.ncbi.nlm.nih.gov > ... > PubMed Central (PMC) ▾ 翻译此页

作者: V Garimella - 2013 - 被引用次数: 7 - 相关文章

Although traditionally the mainstay of postoperative analgesia is opioid based, ... Enhanced recovery protocols to reduce length of stay in colorectal surgery are ... acute pain is defined as pain present in a surgical patient after a procedure. or COX-2 inhibitors are given in addition to PCA morphine after surgery with a ...

[Recent Clinical Nutrition ESPEN Articles - Elsevier](#)

www.journals.elsevier.com/clinical-nutrition.../recent-articles/ ▾ 翻译此页

Implementation of an Enhanced Recovery After Surgery (ERAS) protocol can improve Opioid-sparing effect of selective cyclooxygenase-2 inhibitors on surgical outcomes after open colorectal surgery within an enhanced recovery after ...

[PDF] Enhanced Recovery after Surgery Pain Management