



All

Images

Videos

翻译成中文

关闭取词

28,900 Results

Any time ▼

## [Peritoneal cancer arising after total abdominal ...](#)

[https://www.researchgate.net/publication/320773239\\_Peritoneal\\_cancer\\_arising\\_after...](https://www.researchgate.net/publication/320773239_Peritoneal_cancer_arising_after...)

Peritoneal cancer arising after total abdominal hysterectomy and bilateral salpingo-oophorectomy for cervical cancer in a patient with right breast cancer and germline mutation of BRCA1 gene: a ...

## [A diagnostic dilemma following risk-reducing surgery for ...](#)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2075500>

We present a case report of a woman with a strong family history of breast cancer who underwent risk-reducing surgery in the form of bilateral salpingo-oophorectomy following a mastectomy for a right-sided breast tumour. Following the finding of a BRCA1 mutation, a prophylactic left-sided mastectomy was performed. After remaining well for ...

Cited by: 7

Author: Manish Chand, Patrick J Moore, Andrew ...

Publish Year: 2007

## [UpToDate](#)

<https://www.uptodate.com/contents/risk-reducing-bilateral-salpingo-oophorectomy-in...>

INTRODUCTION. Risk-reducing bilateral salpingo-oophorectomy (rrBSO) is an important option for decreasing the risk of epithelial ovarian and fallopian tubal cancer in women who have the highest likelihood of developing this malignancy []. Other preventive measures and surveillance measures are available, but are of limited efficacy in reducing the high rates of cancer mortality.

## [Primary peritoneal carcinoma metastasizing to breast: a ...](#)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5069842>

The prognosis of PPC is usually poor with extensive cancer spread at the initial diagnosis. The pathogenesis of PPC remains illusive. In this article, we report a rare case of primary peritoneal cancer with breast metastasis in a patient 10 years after total abdominal hysterectomy and bilateral salpingo-oophorectomy.

Cited by: 2

Author: Ji-Yuan Sun, Wondwossen Gebre, Yi-Min...

Publish Year: 2016

## Match Overview

1	<b>Crossref</b> 19 words Michiko Harao, Jiro Ando, Hiroyuki Kamata, Nobuo Hoshi, Seiji Igarashi, Ryuzo Sekiguchi, Kokichi Sugano. "Peritone	1%
2	<b>Internet</b> 16 words crawled on 12-Sep-2017 <a href="http://link.springer.com">link.springer.com</a>	1%
3	<b>Crossref</b> 12 words "Frontiers in Ovarian Cancer Science", Springer Science and Business Media LLC, 2017	<1%

1 **Name of Journal:** *World Journal of Clinical Cases*

2 **Manuscript NO:** 49978

3 **Manuscript Type:** CASE REPORT

4

5 **Peritoneal cancer after bilateral mastectomy, hysterectomy and bilateral**

6 **salpingo-oophorectomy with poor prognosis: A case report and review of**

7 **literature**

8

9 Ma YN *et al.* Poor prognosis of peritoneal cancer

10

11 Ya-Na Ma, Hua-Lei Bu, Cheng-Juan Jin, Xia Wang, You-Zhong Zhang, Hui

12 Zhang

13



国内版

国际版

Peritoneal cancer after bilateral mastectomy, hysterectomy and bilateral salpingo c



All

Images

Videos

关闭取词

27,100 Results

Any time ▼

## A diagnostic dilemma following risk-reducing surgery for ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2075500>

When considering surgical prophylaxis, bilateral salpingo-oophorectomy and total hysterectomy is the most effective procedure to reduce the risk of development of ovarian and peritoneal carcinoma . The estimated risk reduction is between 80–85% for ovarian, Fallopian tube and peritoneal cancer [ ...

Cited by: 7

Author: Manish Chand, Patrick J Moore, Andrew ...

Publish Year: 2007

## Peritoneal cancer arising after total abdominal ...

[https://www.researchgate.net/publication/320773239\\_Peritoneal\\_cancer\\_arising\\_after...](https://www.researchgate.net/publication/320773239_Peritoneal_cancer_arising_after...)

Peritoneal cancer arising after total abdominal hysterectomy and bilateral salpingo-oophorectomy for cervical cancer in a patient with right breast cancer and germline mutation of BRCA1 gene: a ...

## Risk-reducing bilateral salpingo-oophorectomy in women at ...

<https://www.uptodate.com/contents/risk-reducing-bilateral-salpingo-oophorectomy-in...>

Risk-reducing bilateral salpingo-oophorectomy (rrBSO) is an important option for decreasing the risk of epithelial ovarian and fallopian tubal cancer in women who have the highest likelihood of developing this malignancy [1]. Other preventive measures and surveillance measures are available, but are ...

## Cardiac and gingival metastasis after total abdominal ...

<https://link.springer.com/article/10.1007/s10397-009-0514-2> ▼

We present a unique case of cardiac and gingival metastases developing 4 years after total abdominal hysterectomy with bilateral salpingo-oophorectomy for primary uterine epithelial angiosarcoma. Initial treatment remains total abdominal hysterectomy with bilateral salpingo-oophorectomy.

Author: Olivier Donnez, Etienne Marbaix, Patri...

Publish Year: 2011



All

Images

Videos

关闭取词

27,100 Results

Any time ▾

### Peritoneal cancer arising after total abdominal ...

[https://www.researchgate.net/publication/320773239\\_Peritoneal\\_cancer\\_arising\\_after...](https://www.researchgate.net/publication/320773239_Peritoneal_cancer_arising_after...)

Peritoneal cancer arising after total abdominal hysterectomy and bilateral salpingo-oophorectomy for cervical cancer in a patient with right breast cancer and germline mutation of BRCA1 gene: a ...

### Risk-reducing bilateral salpingo-oophorectomy in women at ...

<https://www.uptodate.com/contents/risk-reducing-bilateral-salpingo-oophorectomy-in...>

Risk-reducing bilateral salpingo-oophorectomy (rrBSO) is an important option for decreasing the risk of epithelial ovarian and fallopian tubal cancer in women who have the highest likelihood of developing this malignancy [1]. Other preventive measures and surveillance measures are available, but are ...

### Ovarian carcinosarcoma: a case report, diagnosis ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4938474>

This report aims to present a rare case of ovarian carcinosarcoma and its current, optimal diagnosis and therapy strategy. Description of case We report the case of a 70-year-old patient, who presented at the Obstetrics and Gynecology Department of the University Hospital of Mainz, with a short history of increased abdominal circumference.

### Surgery for BRCA, TP53 and PALB2: a literature review ...

<https://ecancer.org/.../article/863-surgery-for-brca-tp53-and-palb2-a-literature-review> ▾

Therefore, ovarian cancer in BRCA mutation carriers tends to be discovered at an advanced stage with poor prognosis. Due to the lack of effective screening methods, prophylactic surgery must be considered. Bilateral salpingo-oophorectomy is a risk-reducing surgical method to reduce the risk of ovarian cancer in BRCA mutation carriers.

### Brain metastasis from primary peritoneal carcinoma: Case ...

<https://www.sciencedirect.com/science/article/pii/S0967586816312772>

Brain metastasis from primary peritoneal carcinoma: Case report and literature review. ... serum CA 125 level of 14,400 U/ml (normal range, 0–35 U/ml). She underwent a total abdominal hysterectomy, a bilateral salpingo-oophorectomy, omentectomy and debulking of the tumour. During laparotomy, she was found to have multiple tumour nodules ...

Author: Xuan Liu, Jun Masuoka, Ryo Ebashi, ... Publish Year: 2017

### [PDF] Small cell carcinoma of the ovary with hypercalcemia: ...

<https://www.heighpubs.org/cjog/pdf/cjog-aid1006.pdf>