

Levels and activities of von Willebrand factor and metalloproteinase with thrombo

全部

图片

视频

新闻

地图

图书

找到约 39,500 条结果

时间不限

过去 1 小时内

过去 24 小时内

过去 1 周内

过去 1 个月内

过去 1 年内

所有结果

精确匹配

[ADAMTS13 ADAM metalloproteinase with thrombospondin type 1 ...](https://www.ncbi.nlm.nih.gov/gene/11093)

<https://www.ncbi.nlm.nih.gov/gene/11093>

16 Jan 2017 ... A disintegrin and metalloproteinase with thrombospondin motifs 13, a disintegrin-like ... Reduced ADAMTS13 activity together with increased VWF levels were ... be an important mechanism for ADAMTS13 inactivation during inflammation; Thus, ... ADAMTS13 levels do not appear to be related to no-reflow ...

[Von Willebrand Factor, ADAMTS13 and D-Dimer Are Correlated ...](https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4500451/)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4500451/>

13 Jul 2015 ... ADAMTS13 activity was determined by fluorescence resonance energy transfer assay. Von Willebrand factor levels were increased in patients with mild ... and the associated inflammatory state in diabetic patients may lead to ... and metalloproteinase with thrombospondin type 1 motif, member 13) is an ...

[Von Willebrand Factor and ADAMTS13 | Arteriosclerosis ...](http://atvb.ahajournals.org/content/31/7/1703)

[atvb.ahajournals.org/content/31/7/1703](http://atvb.ahajournals.org/content/31/7/1703)

This association was independent of VWF levels and preeclampsia risk factors but ... and metalloproteinase thrombospondin type 1 motif, member 13 (ADAMTS13 ) ... no association between ADAMTS13 (activity or Ag), VWF:Ag, P-selectin, or TSP-1 ... with inflammatory bowel disease displaying high or low CRP levels (10 mg/L ...

[ADAMTS13 - ADAM metalloproteinase with thrombospondin type 1 ...](https://www.ncbi.nlm.nih.gov/gene/11093)

## Match Overview

1	<b>Crossref</b> 170 words Mazur, P., D. Plicner, J. Zdziarska, J. Sadowski, and A. Undas. "Decreased von Willebrand factor ristocetin cofactor	5%
2	<b>Internet</b> 27 words crawled on 08-Jan-2017 <a href="http://www.wjgnet.com">www.wjgnet.com</a>	1%
3	<b>Crossref</b> 26 words Owczarek, D.. "Activated thrombin activatable fibrinolysis inhibitor (TAFIa) is associated with inflammatory mark...	1%
4	<b>Crossref</b> 21 words Owczarek, Danuta, Dorota Cibor, Kinga Salapa, Mikołaj K. Glowacki, Tomasz Mach, and Anetta Undas. "Reduce	1%
5	<b>Crossref</b> 18 words A. UNDAS. "Altered fibrin clot structure in patients with advanced coronary artery disease: a role of C-reactive pri...	<1%
6	<b>Crossref</b> 17 words Graham, Susan M., Junmei Chen, Dominic W. Chung, Kevin R. Barker, Andrea L. Conroy, Michael T. Hawkes, So	<1%
7	<b>Crossref</b> 15 words "P266. Assessing fatigue in Inflammatory Bowel Disease: a comparison and validation of three fatigue scales: IB...	<1%
8	<b>Internet</b> 15 words crawled on 28-Jul-2015 <a href="http://www.e-sciencecentral.org">www.e-sciencecentral.org</a>	<1%

**Name of journal:** *World Journal of Gastroenterology*

**Manuscript NO:** 33189

**Manuscript Type:** ORIGINAL ARTICLE

**Observational Study**

**Levels and activities of von Willebrand factor and metalloproteinase with thrombospondin type-1 motif, number 13 in inflammatory bowel diseases**

Dorota Cibor, Danuta Owczarek, Saulius Butenas, Kinga Salapa, Tomasz Mach, Anetta Undas

**Abstract**

**AIM**

To evaluate the levels of von Willebrand factor (VWF) and metalloproteinase with thrombospondin type-1 motif, number 13 (ADAMTS13) in inflammatory bowel



Levels and activities of von Willebrand factor and metalloproteinase with thrombo

全部 图片 视频 新闻 地图 图书

找到约 56,100 条结果

时间不限

过去 1 小时内  
过去 24 小时内  
过去 1 周内  
过去 1 个月内  
过去 1 年内

所有结果

精确匹配

[ADAMTS13 ADAM metalloproteinase with thrombospondin type 1 ...](https://www.ncbi.nlm.nih.gov/gene/11093)

<https://www.ncbi.nlm.nih.gov/gene/11093>

3 days ago ... A disintegrin and metalloproteinase with thrombospondin motifs 13, a disintegrin-like ... Reduced ADAMTS13 activity together with increased VWF levels were ... be an important mechanism for ADAMTS13 inactivation during inflammation; Thus, ... ADAMTS13 levels do not appear to be related to no-reflow ...

[ADAMTS13 activity to antigen ratio in physiological and ... - NCBI](https://www.ncbi.nlm.nih.gov/pubmed/17608762)

<https://www.ncbi.nlm.nih.gov/pubmed/17608762>

29 Jun 2007 ... The plasma metalloproteinase ADAMTS13 (A Disintegrin And Metalloproteinase with Thrombospondin type 1 motif 13) cleaves prothrombotic ultralarge multimers of the platelet-adhesive protein von Willebrand factor (ULVWF) into less active ... inflammatory bowel disease (n = 44) and cardiac surgery (n = 30).

[\[PDF\] Von Willebrand Factor and ADAMTS13 - Arteriosclerosis ...](http://atvb.ahajournals.org/content/atvbaha/31/7/1703.full.pdf)

[atvb.ahajournals.org/content/atvbaha/31/7/1703.full.pdf](http://atvb.ahajournals.org/content/atvbaha/31/7/1703.full.pdf)

thrombospondin type 1 motif, member 13 (ADAMTS13) levels and the occurrence of preeclampsia, its characteristics ... quantitative, occurs early, and seems to be dependent on inflammation ... whereas TSP-1 has not yet been measured in a large number ... (except for VWF), ADAMTS13 activity and Ag and VWF Ag levels .

[ADAMTS13 - ADAM metalloproteinase with thrombospondin type 1 ...](http://www.ihop-net.org/UniPub/iHOP/bng/96111.html)

[www.ihop-net.org/UniPub/iHOP/bng/96111.html](http://www.ihop-net.org/UniPub/iHOP/bng/96111.html)

Changes in plasma von Willebrand factor and ADAMTS13 levels associated with left atrial ... and ADAMTS-13 in individuals with type 1 and type 2B von Willebrand disease. ... No discrepancies were noted in liver cirrhosis and inflammatory bowel ... A disintegrin and metalloproteinase with a thrombospondin type 1 motifs 13 ...

[Von Willebrand Factor, ADAMTS13 and D-Dimer Are Correlated ...](#)



All

Images

News

Shopping

Videos

More

Settings

Tools

About 56,500 results (0.96 seconds)

## ADAMTS13 ADAM metalloproteinase with thrombospondin type 1 motif ...

<https://www.ncbi.nlm.nih.gov/gene/11093>

7 days ago - A disintegrin and **metalloproteinase** with **thrombospondin motifs 13**, a disintegrin-like ... Reduced **ADAMTS13 activity** together with increased **VWF levels** were ... be an important mechanism for **ADAMTS13** inactivation during **inflammation**; Thus, .... **ADAMTS13 levels** do not appear to be related to **no-reflow** ...

## ADAMTS13 activity to antigen ratio in physiological and ... - NCBI

<https://www.ncbi.nlm.nih.gov/pubmed/17608762>

by HB Feys - 2007 - Cited by 90 - Related articles

29 Jun 2007 - The plasma **metalloproteinase ADAMTS13** (A Disintegrin And **Metalloproteinase** with **ThromboSpondin type 1 motif 13**) cleaves prothrombotic ultralarge multimers of the platelet-adhesive protein **von Willebrand factor** (ULVWF) into less active ... **inflammatory bowel disease** (n = 44) and cardiac surgery (n = 30).

## Von Willebrand Factor and ADAMTS13 - Arteriosclerosis, Thrombosis ...

[atvb.ahajournals.org/content/atvbaha/31/7/1703.full.pdf](http://atvb.ahajournals.org/content/atvbaha/31/7/1703.full.pdf)

by A Stepanian - 2011 - Cited by 43 - Related articles

**thrombospondin type 1 motif**, member **13 (ADAMTS13) levels** and the occurrence of preeclampsia, its characteristics ... quantitative, occurs early, and seems to be dependent on **inflammation**. ... whereas TSP-1 has not yet been measured in a large **number** .... (except for **VWF**), **ADAMTS13 activity** and Ag and VWF Ag **levels**.

## Von Willebrand Factor, ADAMTS13 and D-Dimer Are Correlated with ...

[journals.plos.org/plosone/article?id=10.1371/journal.pone.0132784](https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0132784) ▼

by CP Domingueti - 2015 - Cited by 5 - Related articles

13 Jul 2015 - **Von Willebrand factor levels** were increased in patients with mild (P=0.001) and ... **ADAMTS13 activity/ADAMTS13 levels** ratio was reduced in patients with mild ... and **metalloproteinase with thrombospondin type 1 motif**, member **13**), and ..... microvascular inflammation in **inflammatory bowel disease**.