



国内版

国际版

Fat clearance and conventional fixation identified ypN0 rectal cancer following int



All

Images

Videos

翻译成中文

关闭取词

14,400 Results

Any time ▼

## Preoperative conventional chemoradiotherapy versus short ...

<https://bmccancer.biomedcentral.com/articles/10.1186/s12885-016-2959-9> ▼

Dec 01, 2016 · Preoperative **conventional** chemoradiotherapy versus short-course **radiotherapy** with delayed surgery for **rectal cancer**: results of a randomized controlled trial ... A major concern is a large group of patients with **intermediate-risk rectal cancer** (T3, >3 mm CRM), for whom **neoadjuvant** chemoradiotherapy appears to be potential overtreatment, but ...

Cited by: 7

Author: Tadas Latkauskas, Henrikas Pauzas, Lau...

Publish Year: 2016

## Evaluation of Lymphadenectomy in Patients Receiving ...

[https://www.researchgate.net/publication/225296525\\_Evaluation\\_of...](https://www.researchgate.net/publication/225296525_Evaluation_of...)

Patients with ypT0-2 **rectal cancer following neoadjuvant** chemoradiation have very small perirectal nodes. Individual metastatic lymph nodes are significantly larger.

## Neoadjuvant chemoradiation and rectal cancer | Journal of ...

[jcp.bmj.com](http://jcp.bmj.com) › Online First ▼

Dec 28, 2018 · Abstract. **Neoadjuvant** chemoradiation (NACR) is now standard of care in stage II and III **rectal cancer**. The advent of this modality of treatment has impacted on the way the pathological evaluation of resection specimens that **have** been subjected to preoperative chemoradiation is ...

Author: Runjan Chetty, Aoife J McCarthy    Publish Year: 2019

## Tumor regression in mesorectal lymphnodes after ...

[https://www.researchgate.net/publication/6471247\\_Tumor\\_regression...](https://www.researchgate.net/publication/6471247_Tumor_regression...)

This study aimed to investigate the application of **fat clearance** in cases of **rectal cancer** after **neoadjuvant** chemoradiation. All patients who underwent proctectomy (R0 resection) from 1998 to 2007 ...

---

**Name of journal:** *World Journal of Gastrointestinal Oncology*

**Manuscript NO:** 47920

**Manuscript Type:** ORIGINAL ARTICLE

*Retrospective Cohort Study*

**Fat clearance and conventional fixation identified ypN0 rectal cancer following intermediate neoadjuvant radiotherapy have similar long-term outcome**

Chen N *et al.* Fat clearance does not bring survival benefit for rectal cancer patients after nCRT

Nan Chen, Ting-Ting Sun, Zhong-Wu Li, Yun-Feng Yao, Lin Wang, Ai-Wen Wu

### Match Overview

2 matches

- 1** **Crossref** 28 words  
Lin Wang, Yong-Heng Li, Yong Cai, Tian-Cheng Zhan, Jin Gu. "Intermediate Neoadjuvant Radiotherapy Combined Wi 1%
- 2** **Crossref** 26 words  
Parsons, Helen M., Todd M. Tuttle, Karen M. Kuntz, James W. Begun, Patricia M. McGovern, and Beth A. Virnig. "Asso ... 1%



All

Images

Videos

翻译成中文

关闭取词

23,700 Results

Any time ▾

## Neoadjuvant chemoradiation and rectal cancer | Journal of ...

<https://jcp.bmj.com/content/72/2/97> ▾

Feb 01, 2019 · **Neoadjuvant** chemoradiation (NACR) is now standard of care in stage II and III **rectal cancer**. The advent of this modality of treatment has impacted on the way the pathological evaluation of resection specimens that have been subjected to preoperative chemoradiation is conducted. The gross description, sectioning and microscopic examination have had to be adapted to accommodate the ...

Author: Runjan Chetty, Aoife J McCarthy    Publish Year: 2019

Online: ISSN 1472-4146

## Evaluation of Lymphadenectomy in Patients Receiving ...

[https://www.researchgate.net/publication/225296525\\_Evaluation\\_of\\_Lymphadenectomy\\_in...](https://www.researchgate.net/publication/225296525_Evaluation_of_Lymphadenectomy_in...)

Evaluation of Lymphadenectomy in Patients Receiving **Neoadjuvant Radiotherapy for Rectal Adenocarcinoma** Article in Annals of Surgical Oncology 19(12):3713-8 · June 2012 with 23 Reads

## Lymph node harvest in colon and rectal cancer: Current ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3277879>

Jan 27, 2012 · The presence of lymph node (LN) metastases in colo-rectal **cancer** is well recognized as one of the most important prognostic factors for **long-term outcome**[1,2]. In clinical practice, the presence of LN metastasis determines those patients most likely to benefit from adjuvant chemotherapy[3 ...

Cited by: 98

Author: James R McDonald, Andrew G Renehan, ...

Publish Year: 2012

## Rectal Cancer, Version 2.2018, NCCN Clinical Practice ...

<https://nccn.org/view/journals/nccn/16/7/article-p874.xml> ▾

Jul 01, 2018 · The ongoing randomized RAPIDO trial (ClinicalTrials.gov identifier: NCT01558921) is assessing DFS at 3 years with the use of preoperative short-course RT followed by 6 cycles of CAPEOX before resection in patients with clinical stage T3–4 **rectal cancer**. 186. Response to **Neoadjuvant Treatment: Following neoadjuvant therapy**, 50% to 60% of patients are downstaged, with ...

Cited by: 27

Author: Al B Benson, Alan P. Venook, Mahmoud ...

Publish Year: 2018

## Evaluating Mesorectal Lymph Nodes in Rectal Cancer Before ...