

85788_Auto_Edited.docx

Name of Journal: *World Journal of Gastroenterology*

Manuscript NO: 85788

Manuscript Type: ORIGINAL ARTICLE

Basic Study

Green tea polyphenols alleviate di-(2-ethylhexyl) phthalate-induced liver injury in mice

GTPs alleviate DEHP-induced liver injury in mice

Heng Shi, Xin-Hai Zhao, Qin Peng, Xian-Ling Zhou, Si-Si Liu, Chuan-Chuan Sun, Qiu-Yu Cao, Shi-Ping Zhu, Sheng-Yun Sun

Abstract

BACKGROUND

Di (2-ethylhexyl) phthalate (DEHP) is a common plasticizer known to cause liver injury. Green tea is reported to exert therapeutic effects on heavy metal exposure-induced organ damage. However, limited studies have examined the therapeutic effects of green tea polyphenols (GTPs) on DEHP-induced liver damage.

AIM

To evaluate the molecular mechanism underlying the therapeutic effects of GTPs on DEHP-induced liver damage.

METHODS

C57BL/6J mice were divided into the following five groups: control, model (DEHP (1500 mg/kg bodyweight)), treatment (DEHP (1500 mg/kg bodyweight) + GTP (70 mg/kg bodyweight), oil, and GTP (70 mg/kg bodyweight) groups. After 8 wk, the liver function, blood lipid profile, and liver histopathology were examined. Differentially expressed miRNAs and mRNAs in the liver tissues were examined using high-throughput sequencing. Additionally, functional enrichment analysis and immune infiltration prediction were performed. The miRNA-mRNA regulatory axis was elucidated using the starBase database. Protein expression was evaluated using immunohistochemistry.

RESULTS

GTPs alleviated DEHP-induced liver dysfunction, blood lipid dysregulation, fatty liver disease, liver fibrosis, and mitochondrial and endoplasmic reticulum lesions in mice. The infiltration of macrophages, mast cells, and natural killer cells varied between the model and treatment groups. *mmu-miR-141-3p* (a differentially expressed miRNA), *Zcchc24* (a differentially expressed mRNA), and *Zcchc24* (a differentially expressed

protein) constituted the miRNA-mRNA-protein regulatory axis involved in mediating the therapeutic effects of GTPs on DEHP-induced liver damage in mice.

CONCLUSION

GTPs alleviated DHEP-induced liver dysfunction, blood lipid dysregulation, fatty liver disease, liver fibrosis, and mitochondrial and endoplasmic reticulum lesions in mice. The infiltration of macrophages, mast cells, and natural killer cells varied between the model and treatment groups. mmu-miR-141-3p (a differentially expressed miRNA), *Zcchc24* (a differentially expressed mRNA), and *Zcchc24* (a differentially expressed protein) constituted the miRNA-mRNA-protein regulatory axis involved in mediating the therapeutic effects of GTPs on DEHP-induced liver damage in mice.

Key Words: Green tea polyphenols; di(2-ethylhexyl) phthalate; liver fibrosis; fatty liver disease; mitochondria; immune

Shi H, Zhao XH, Peng Q, Zhou XL, Liu SS, Sun CC, Cao QY, Zhu SP, Sun SY. Green tea polyphenols alleviate di-(2-ethylhexyl) phthalate-induced liver injury in mice. *World J Gastroenterol* 2023; In press

Core Tip: GTPs alleviated DEHP-induced liver dysfunction, blood lipid dysregulation, fatty liver disease, liver fibrosis, and mitochondrial and endoplasmic reticulum lesions in mice. The infiltration of macrophages, mast cell, and NK cells varied between the model and treatment groups. mmu-miR-141-3p (a differentially expressed miRNA), *Zcchc24* (a differentially expressed mRNA), and *Zcchc24* (a differentially expressed protein) constituted the miRNA-mRNA-protein regulatory axis involved in mediating the therapeutic effects of GTPs on DEHP-induced liver damage in mice.

2 INTRODUCTION

Di-(2-ethylhexyl) phthalate (DEHP), which is the most widely used representative phthalic acid ester, can non-covalently bind to polyolefin plastics and predominantly serves as a plasticizer, increasing the flexibility, transparency, durability, and longevity of plastics. Additionally, DEHP is extensively detected in various daily-life products (including baby toys, food packaging, and cosmetics) and several surgical and medical devices ^[1]. Furthermore, DEHP ² can be continuously released into the environment (air, soil, water, food, etc.), enter living organisms *via* ingestion, inhalation, or skin contact, and subsequently exert toxic effects on health ^[2]. A previous study reported that DEHP and its metabolites were detected in 100% of tested human urine samples, indicating persistent exposure to DEHP ^[3]. The DEHP exposure range of the general population is estimated to be 5.8–19 µg/kg/d, while DEHP exposure in medical environments may exceed 167.9 mg/d ^[4]. The development of modern chemical agriculture has increased the severity of DEHP pollution. DEHP exposure in the Pearl River Delta region of Guangdong Province, China, can reach up to 61 µg/kg/d, which is higher than the tolerable intake of DEHP ^[5]. Additionally, DEHP undergoes rapid degradation upon ingestion in humans. According to ² the US Environmental Protection Agency, the average half-life of DEHP in the human body is 12 h. DEHP and its active metabolite mono-(2-ethylhexyl) phthalate have been detected in various human tissues, including the liver, blood, placenta, amniotic fluid, and early-pregnancy chorionic villus sample ^[6]. Therefore, evaluating the effect of DEHP on the environment and human health has piqued the interest of the scientific community.

The liver, an important organ involved in the synthesis, metabolism, and detoxification processes, is highly susceptible to acute or chronic damage induced by various drugs ^[7]. Recent studies have demonstrated that DEHP adversely affects multiple systems in the body. In particular, epidemiological and animal studies have demonstrated the hepatotoxicity of DEHP ^[8]. The mechanism underlying the hepatotoxic effects of DEHP mainly involves oxidative stress, cell apoptosis, and signaling pathway activation. DEHP induces apoptosis in healthy human liver cells through the mitochondrial signaling pathway and/or the caspase-mediated death receptor pathway ^[9].

2
Additionally, DEHP may adversely affect gap junctional intercellular communication, peroxisome beta-oxidation activity, and DNA replication synthesis, leading to the formation of liver tumors [10]. Currently, the most extensively studied mechanism of DEHP-induced liver damage is oxidative stress in which the regulation of reactive oxygen species (ROS) production plays a critical role [11]. 15
Excessive ROS production leads to peroxidation of the polyunsaturated fatty acids in the cell membrane. DEHP promotes lipid peroxidation in the liver, primarily through the downregulation of superoxide dismutase (SOD) and catalase activity and the upregulation of malondialdehyde (MDA) concentrations [12]. Additionally, DEHP mediates the pathogenesis of non-alcoholic fatty liver disease in a high-fat diet-fed animal model by promoting lipid peroxidation [8].

Green tea is the second most widely consumed beverage worldwide after water [13]. The harvested tea leaves are steamed at high temperatures to inactivate the polyphenol oxidase in green tea, preserving most of the vitamins in tea leaves. Thus, green tea exhibits enhanced antioxidant activity. Epidemiological and clinical studies have reported that phenol compounds in tea extracts, such as green tea polyphenols (GTPs), exert a wide range of beneficial effects on human health, including anti-aging, neuroprotective [14], and therapeutic or preventive effects on various diseases, such as cancer [15], cardiovascular disease [16], and obesity [17]. Moreover, GTPs mitigate the adverse effects of poisoning with various heavy metals [18]. However, limited studies have examined the therapeutic effects of GTPs on DEHP-induced liver diseases.

Micro RNAs (miRNAs), a class of endogenous non-coding small RNAs encoded by mRNA, regulate gene expression by modulating mRNA stability. Several miRNAs are reported to play important roles in the pathogenesis of liver fibrosis [19], cirrhosis [20], and hepatocellular carcinoma [21]. Some miRNAs are diagnostic markers for drug-induced liver injury [22]. This study aimed to identify key protein nodes that may affect the expression level of miRNAs in the regulatory network of miRNA and target genes by analyzing the miRNA regulation network. Hence, this study will provide a

theoretical basis for future studies on the functions and regulatory mechanisms underlying the therapeutic effects of GTPs on DEHP exposure-induced liver damage.

MATERIALS AND METHODS

Reagents

DEHP was purchased from Shanghai Aladdin Biochemical Technology Co., Ltd. (Shanghai, China). GTPs were purchased from Shanghai Yuanye Bio-Technology Co., Ltd. (Shanghai, China). Corn oil was purchased from Hebei Pin Research Biotechnology Co., Ltd. (Baoding, China). Anti-ZCCHC24 antibodies were purchased from Biorbyt Ltd. (Cambridge, UK).

Animal experiments

Animal experiments were performed at the specific pathogen-free grade animal laboratory of the Medical College of Jinan University. C57BL/6J mice ($n = 50$) were purchased from Guangdong Yaokang Biotechnology Co., Ltd. And allowed to acclimatize to the laboratory environment for 1 wk. The animals were maintained under the following conditions: room temperature, 20°C–24°C; relative humidity, 50%–65%; circadian cycle, 12-h light-dark cycle; access to food and water, ad libitum; diet, regular mouse chow. This study was approved by the Institutional Animal Care and Use Committee of Jinan University (ethics approval number: IACUC-20210630-15). All experimental procedures were performed according to the regulations established by the ethical committee.

Previous studies ^[23, 24] have reported that DEHP is soluble in corn oil. Hence, this study used corn oil as the solvent for DEHP. After 1 wk of acclimatization, 50 mice were randomly assigned into the following five groups (10 mice/group): control group, administered with 0.2 mL distilled water; model group, administered with 0.2 mL corn oil and DEHP (1500 mg/kg bodyweight); treatment group, administered with 0.2 mL corn oil and DEHP (1500 mg/kg bodyweight), followed by administration of 0.2 mL GTPs (70 mg/kg bodyweight) after 1 h; oil group, administered with 0.2 mL corn oil; GTP group, administered with 0.2 mL GTP (70 mg/kg bodyweight). Based on our

previous study, DEHP [25] and GTPs [26] were gavaged. The doses were adjusted weekly based on the bodyweight of the mice. The mice were subjected to daily gavage for 8 wk. At the end of the experimental period, the blood and liver samples were obtained under anesthesia after the mice were allowed to fast overnight. Liver index = (liver weight/bodyweight) x 100%.

Evaluation of liver function and blood lipid profile

The blood from mice was collected using the retro-orbital venous plexus method. Next, the whole blood was placed in a 1.5-mL centrifuge tube and left undisturbed at room temperature for 2 h. The sample was centrifuged at 5°C and 3000 rpm for 15 min using a high-speed refrigerated centrifuge to obtain the serum. The serum levels of liver function markers (aspartate aminotransferase (AST), alanine aminotransferase (ALT), and total bilirubin (TBIL)) and lipids (low-density lipoprotein (LDL), total cholesterol, and triglycerides (TGs)) were analyzed using a fully automated biochemical analyzer.

Hematoxylin and eosin (HE) staining

The tissue sections were fixed in 4% neutral buffered formalin, embedded in paraffin, and sectioned into 4 µm-thick sections. The sections were deparaffinized using xylene, rehydrated using a series of graded ethanol solutions, stained with hematoxylin for 5–10 min, washed with distilled water, and stained with eosin for 2–7 min. Next, the sections were dehydrated using a series of graded ethanol solutions, cleared with xylene, mounted with mounting medium, and covered with a cover slip.

Oil red O staining

The frozen sections were fixed in 4% neutral buffered formalin, rinsed with 60% isopropanol, and allowed to dry. The sections were then stained with oil red O for 15–30 min, rinsed with 60% isopropanol, counterstained with hematoxylin for 1–2 min, washed with distilled water, dehydrated, mounted with a mounting medium, and covered with a coverslip.

Periodic acid-Schiff (PAS) staining

The tissue sections were deparaffinized using xylene, rehydrated using a series of graded ethanol solutions, and oxidized with a periodic acid solution for 10–15 min.

Next, the sections were rinsed with distilled water, stained with Schiff reagent for 30–60 min in the dark, counterstained with hematoxylin, washed with distilled water, dehydrated, mounted using a mounting medium, and covered with a cover slip.

Masson's trichrome staining

The tissue sections were deparaffinized using xylene, rehydrated using a series of graded ethanol solutions, stained with Weigert's iron hematoxylin for 10 min, rinsed with distilled water, stained with Biebrich scarlet-acid fuchsin solution for 5–10 min, washed with distilled water, and incubated with phosphotungstic-phosphomolybdic acid solution for 5–10 min. Next, the sections were counterstained with aniline blue for 5–10 min, washed with distilled water, dehydrated, mounted with a mounting medium, and covered with a cover slip.

Sirius red staining

The tissue sections were deparaffinized using xylene, rehydrated using a series of graded ethanol solutions, stained with Sirius red solution for 1 h, washed with distilled water, dehydrated, and cleared with xylene. Finally, the sections were mounted with a mounting medium and covered with a cover slip.

Transmission electron microscopy

One mouse from the control, treatment, and model groups was randomly selected for electron microscopy. Liver sections with a size of approximately 1 mm were treated with 2.5% glutaraldehyde (Scientific Phygene, Fuzhou, China) and rinsed thrice with phosphate-buffered saline (PBS) (PH = 7.4). The samples were fixed with 1% osmium tetroxide (Ted Pella, Redding, USA) for 2 h, rinsed thrice with PBS, and dehydrated using alcohol and acetone gradients as follows: 50% ethanol for 30 min, 70% ethanol for 30 h, 80% acetone for 30 min, and 90% acetone for 30 h. Next, the samples were washed thrice with 100% acetone and embedded in epoxy resin (Ted Pella, Redding, USA). Ultrathin sections (7 nm) were prepared using a microtome (LKB, Bromma, Sweden). The sections were stained with uranyl acetate (EMS, Hatfield, USA) for 30 min and 3% lead citrate (Ted Pella, Redding, USA) for 15 min. The target structures were observed

using a transmission electron microscope (JEM-1400, Japan Electron Optics Laboratory Co., Ltd. Tokyo, Japan).

High-throughput sequencing

Liver samples ($n = 3$ per group) from the control, treatment, and model groups stored in liquid nitrogen were randomly selected and sent to Huada Genomics (Wuhan, China, <http://www.genomics.cn>) for high-throughput sequencing. For specific sequencing steps, refer to the supplementary materials.

Differentially expressed mRNAs and miRNAs

The mRNA and miRNA expression levels were measured using fragments per kilobase of transcript per million mapped reads (FPKM) values. Sequencing data were subjected to quality control. Differential expression between groups was estimated using the Limma R package (version: 3.52.2) based on the generalized linear model. To comparatively analyze the expression levels between multiple groups, the control-model and model-treatment expression profiles were regressed using the Limma package in R software (version: 4.2.1, <https://posit.co/>). A loose threshold was set to avoid excessive filtering of differentially expressed mRNAs and miRNAs (log fold-change (FC) > 1 ; $p < 0.05$). Differentially expressed mRNAs and miRNAs were identified.

Enrichment analysis

The mRNA expression data were subjected to Gene Ontology (GO) and Kyoto Encyclopedia of Genes and Genomes (KEGG) pathway analyses to identify the enrichment of mRNAs in the GO terms biological process (BP), cellular component (CC), and molecular function (MF) and the KEGG signaling pathways. The clusterprofiler R package (version: 4.4.4) was used for enrichment analyses with the reference files based on the org.Mm.eg.db R package (version 3.1.0) for mouse enrichment analysis. Enrichment was considered significant at $p < 0.05$.

Immune infiltration analysis

CIBERSORT is a gene expression-based algorithm for the accurate detection of immune cell infiltration. The "CIBERSORT" R package (CIBERSORT script v1.03;

<http://cibersort.stanford.edu/>) developed by Newman *et al* [27] was used to successfully quantify 22 types of immune cells, including B cells, regulatory T cells, CD4+ T, CD8+ T, natural killer (NK) cells, mast cells, plasma cells, dendritic cells, neutrophils, eosinophils, and macrophages. The reference dataset of mouse immune cells was obtained from Chen *et al* [28]. The mRNA expression data of the control, model, and treatment groups were examined using the CIBERSORT algorithm with a perm of the deconvolution algorithm set to 1000.

Regulatory network of miRNA-mRNA

miRNA, a type of non-coding RNA, negatively regulates gene expression at the posttranscriptional level. To generate a preliminary regulatory network of miRNA-mRNA in the model and treatment groups, the upregulated miRNAs or mRNAs and downregulated mRNAs or miRNAs were selected and analyzed using Cytoscape software (version: 3.7.1, <https://cytoscape.org/>).

The Sun Yat-sen University research team developed the starBase database (<https://starbase.sysu.edu.cn/starbase2/>) to analyze interaction networks between long non-coding RNA, miRNA, competitive endogenous RNAs, RNA-binding proteins, and mRNAs using cross-linked immunoprecipitation sequencing (CLIP-seq) (high-throughput sequencing of RNA using CLIP, photoactivable ribonucleoside-enhanced CLIP, individual nucleotide resolution ultraviolet CLIP, and cross-linking ligation and sequencing of hybrids) data. To obtain the final miRNA-mRNA regulatory network, the initially selected network was intersected with the mouse interaction network obtained from the starBase database. Gene expression was verified using immunohistochemical analyses. This study aimed to establish the miRNA-mRNA-protein regulatory axis.

Immunohistochemistry

The paraffin sections (4 μ m) of mouse liver tissue were deparaffinized, rehydrated, and subjected to antigen retrieval in a buffer solution (pH 9.0). After blocking endogenous peroxidase activity and non-specific binding, the sections were incubated with anti-ZCCHC24 antibodies (1:100) at 4°C overnight. The sections were then washed and incubated with horseradish peroxidase-conjugated secondary antibodies at room

temperature for 1 h. Immunoreactive signals were developed using diaminobenzidine. The sections were counterstained with hematoxylin and observed under a microscope.

Statistical analyses

The average optical density (AOD) (integrated optical density/area) of positive reactions was calculated using Image-Pro Plus 6.0 software. Data were analyzed using the dplyr R package (v. 1.0.10) and visualized using the ggplot2 R package (v. 3.3.6). Categorical variables were analyzed using the Chi-squared test. Continuous variables were represented as mean \pm standard deviation. Continuous variables between two groups were compared using the Wilcoxon signed-rank test, while those between more than two groups were compared using the Kruskal-Wallis test. Differences were considered significant at $p < 0.05$.

RESULTS

Effect of GTPs on bodyweight and liver index

The flowchart of the study is shown in Figure 1. Mice in all groups survived during the experiment and did not exhibit aberrant behaviors in urination, defecation, food intake, or water consumption. As shown in Table 1 and Figure 2A, the bodyweight of mice in all groups increased with time and was not significantly different between the groups. The liver index (Table 3 and Figure 2B) values in the model group were significantly higher than those in the control group ($P = 0.016$) but were not significantly different between the treatment and model groups ($P = 0.51$). Additionally, the liver index values in the oil and GTP groups were not significantly different from those in the control group ($p > 0.05$).

Effect of GTPs on the serum levels of liver function markers and lipids

The serum levels of ALT, AST, TBIL, LDL, and TGs were analyzed. As shown in Figure 2C–2G, the serum levels of ALT, AST, TBIL, LDL, and TGs in the model group were significantly higher than those in the control group ($p < 0.001$). Compared with those in the model group, the serum levels of liver function markers and lipids were significantly downregulated in the treatment group ($p < 0.001$). The ALT, AST, and LDL

levels were not significantly different between the oil, control, and GTP groups ($p > 0.05$).

Effect of GTPs on hepatic histological characteristics

The results of liver HE staining are shown in Figure 3A. Mice in the model group exhibited significant hepatocyte ballooning degeneration, whereas those in the control, oil, and GTP groups did not exhibit hepatocyte ballooning degeneration.

Effect of GTPs on fat deposition

Oil red O staining was performed to further analyze the severity of fatty liver in different groups. As shown in Figure 3A, the model group exhibited the highest red color staining intensity in the liver, followed by the oil and control groups. Meanwhile, the liver of the treatment and GTP groups exhibited the lowest red staining intensity. Five positive staining sites in the liver of the model and treatment groups were selected to calculate the AOD. As shown in Figure 3B, the AOD in the model group ($7.270 \pm 1.120\%$) was significantly higher than that in the treatment ($0.185 \pm 0.061\%$) ($P = 0.037$) and control groups ($5.760 \pm 0.586\%$) ($p < 0.001$).

Effect of GTPs on polysaccharide accumulation

The results of PAS staining (Figure 3A) revealed the lack of red-stained areas in the liver of the model group, while the red-stained areas were upregulated in the control, oil, treatment, and GTP groups. Five positive staining sites in the liver of the model and treatment groups were selected to calculate the AOD. As shown in Figure 3C, the AOD in the model group ($0.431 \pm 0.083\%$) was significantly lower than that in the treatment ($1.170 \pm 0.099\%$) and control groups ($1.360 \pm 0.148\%$) ($p < 0.001$).

Effect of GTPs on collagen fibers

The results of Masson's trichrome staining (Figure 4A) revealed blue-stained fibrous tissue around the liver blood vessels in the model group. In contrast, the area of blue-stained fibrous tissue around the liver blood vessels in the treatment group was lower than that in the model group. Fibrous tissue was not detected around the liver blood vessels in the control, oil, and GTP groups. Five positive staining sites in the liver of the model and treatment groups were selected to calculate the AOD. As shown in Figure

4B, the AOD¹ in the model group ($0.337 \pm 0.113\%$) was significantly higher than that in the treatment group ($0.183 \pm 0.014\%$) ($P = 0.02$).

Effect of GTPs on collagen network

In Sirius red staining, type I collagen fibers exhibit strong orange-yellow or bright red colors under a polarized light microscope, whereas type III collagen fibers exhibit green color. In this study, the results of Sirius red staining (Figure 4A) revealed red-stained fibrous tissue around the blood vessels in the model group. The red-stained area around the blood vessels¹ in the treatment group was significantly lower than that in the model group. Red-stained fibrous tissue was not detected around the blood vessels of the control, oil, and GTP groups. Next, semiquantitative analysis was performed by selecting five positive staining sites in the liver of the model and treatment groups to calculate the AOD. The AOD³ in the model group ($1.240 \pm 0.125\%$) was significantly higher than that in the treatment group ($0.080 \pm 0.025\%$) ($P = 0.012$) (Figure 4C).

Effect of GTPs on the liver microstructures

The liver tissues of the control, model, and treatment groups were subjected to transmission electron microscopy. The number and size of lipid droplets in the liver were upregulated and the capillary bile ducts were significantly dilated in the model group (Figure 5A). Additionally, the mitochondria were slightly swollen, and the arrangement of the rough endoplasmic reticulum was disordered. In contrast, the number of lipid droplets in the liver was downregulated in the treatment group, while the capillary bile ducts exhibited physiological structure. Additionally, the swelling of the mitochondria was alleviated. The results of quantitative analysis of the diameter of small bile ducts (Figure 5B) were consistent with the observations in Figure 5A. However, these findings must be carefully interpreted. Additionally, further studies with large sample sizes are needed to confirm and generalize the results.

Effect of GTPs on mRNA and miRNA expression levels

The liver samples from the control, treatment, and model groups were subjected to high-throughput screening with 3 replicate samples for each group. The mRNA and miRNA expression data were subjected to quality control analysis. The differences in

the expression levels of miRNA and mRNA were minimal between the groups (Supplementary Figure 1A–1B). Next, principal component analysis was performed (Supplementary Figure 1C–1D). The mRNA and miRNA profiles of the control, model, and treatment groups exhibited distinct separation. Differential analysis revealed that compared with those in the model group, the number of upregulated mRNAs and miRNAs was 377 and 33, respectively, while the number of downregulated mRNAs and miRNAs was 583 and 7, respectively (Figure 6A–6B). The expression levels of the significantly upregulated and downregulated mRNAs and miRNAs are shown in Figure 6C–6D.

Enrichment analyses

⁴ Compared with those in the control group, the differentially expressed mRNAs in the model group were enriched in the following GO terms: BP term: fatty acid metabolic process, small molecule catabolic process, organic acid catabolic process, and carboxylic acid catabolic process (Figure 7A); CC term: mitochondrial protein-containing complex, mitochondrial inner membrane, ribosome subunit, and ribosome (Figure 7B); MF term: ribosome structure, electron transfer activity, oxidoreductase activity, and ubiquitin protein ligase binding (Figure 7C). Additionally, the differentially expressed mRNAs in the model group were enriched in the following KEGG pathways: fatty acid metabolism, ferroptosis, glutathione metabolism, and PPAR signaling pathway (Figure 7D). ¹² Compared with those in the model group, the differentially expressed mRNAs in the treatment group were enriched in the following GO terms: BP term: mitochondrial organization, proteasome protein catabolic process, and oxidative phosphorylation (Figure 8A); CC term: ribosome, ribosomal subunit, and large ribosomal subunit (Figure 8B); MF term: structural constituent of ribosome and molecular carrier activity (Figure 8C). Additionally, the differentially expressed mRNAs in the treatment group were enriched in the following KEGG pathways: mitochondrial autophagy, glutathione metabolism, oxidative phosphorylation, and drug metabolism (Figure 8D).

Immune infiltration analysis

The proportions and infiltration levels of immune cells in the control, model, and treatment groups are shown in Figure 9A–9B. Compared with those in the control group, the infiltration levels of monocytes and immature CD8⁺ T cells were significantly upregulated and the infiltration levels of immature CD4⁺ T cells were significantly downregulated ¹³ in the model group (Figure 9C). Meanwhile, compared with those in the model group, the infiltration levels of mast cells and active NK cells were significantly upregulated and the infiltration levels of M2 macrophages were significantly downregulated in the treatment group (Figure 9D). The correlation analysis results of various immune cells are shown in Figure 9E.

Construction of miRNA-mRNA-protein regulatory axis

Network diagrams (Figure 10A) of upregulated miRNAs and downregulated mRNAs, as well as that of downregulated miRNAs and upregulated mRNAs, in the treatment group were constructed. The data were matched with the miRNA-mRNA regulatory axis in the starBase database to obtain the mmu-miR-141-3p/Zcchc24 and mmu-miR-9-5p/Zbtb7a axes (Figure 10B). Correlation analysis (Figure 10C–10D) revealed that the correlation coefficients of mmu-miR-141-3p/Zcchc24 and mmu-miR-9-5p/Zbtb7a axes were −0.44 and −0.87, respectively. To validate these regulatory axes, the liver tissues of the control, model, and treatment groups were subjected to western blotting (Figure 10E). ¹ Compared with that in the model group, the average gray value of Zcchc24 protein was significantly downregulated in the treatment group (Figure 10F), indicating that protein expression changes were consistent with mRNA expression changes. However, Zbtb7a protein expression could not be validated using western blotting or immunohistochemical analysis due to low expression levels. These findings indicate that the mmu-miR-141-3p/Zcchc24 (mRNA)/Zcchc24 (protein) regulatory axis plays an important role in the protective effects of GTPs on DEHP-induced liver injury in mice.

DISCUSSION

This study aimed to investigate the therapeutic mechanism of GTPs in alleviating DEHP-induced liver dysfunction, blood lipid dysregulation, non-alcoholic fatty liver,

and liver fibrosis. Additionally, the role of a miRNA-mRNA-protein regulatory axis in the therapeutic effects of GTPs on DEHP-induced liver damage was elucidated (Figure 11). Thus, the findings of this study provide valuable insights into the potential therapeutic application of GTPs in DEHP-induced liver damage.

Histopathological analysis revealed that GTPs are effective in mitigating DEHP-induced non-alcoholic fatty liver disease and liver fibrosis in mice. This finding is consistent with that of previous studies, which reported the beneficial effects of GTP on liver health. For example, a meta-analysis [29] of 20 randomized controlled trials with 1536 participants revealed that green tea decreases the levels of total cholesterol and LDL. Zhao *et al* [30] reported that DEHP exposure significantly promotes inflammation, necrosis, and fibrosis in the liver and upregulates the expression of proteins associated with the development of liver inflammation and fibrosis. Kim *et al* [31] suggested that GTPs downregulate the expression of collagen content and type 1 collagen, and consequently alleviate liver fibrosis. Additionally, animal studies [32] have indicated that supplementation of green tea effectively prevented excessive accumulation of visceral and liver lipids, elevated blood glucose levels, and alleviated aberrant blood lipid levels, liver dysfunction, and hepatic steatosis in male C57BL/6 mice fed on a high-fat diet for six weeks as evidenced by the analysis of serum biochemical parameters, histological changes, lipid accumulation, inflammatory cytokines, and related indices. These findings further indicated the therapeutic potential of GTPs in liver-related conditions.

In this study, GTPs were found to ameliorate the DEHP-induced pathological damage to the liver microstructures, including the mitochondria, endoplasmic reticulum, and capillary bile ducts. These findings are consistent with those of previous studies, which reported the adverse effects of DEHP on liver microstructures. For instance, an animal study [33] indicated that DEHP induced mitochondrial and endoplasmic reticulum ultrastructural damages, characterized by increased fission and decreased fusion. Furthermore, Sun *et al* [34] suggested that DEHP promoted mitochondrial-associated endoplasmic reticulum membrane disruption, potentially through endoplasmic

reticulum unfolded protein response, to induce endoplasmic reticulum stress. Consistent with the results of this study, *in vitro* and *in vivo* experimental studies [35] have demonstrated that GTPs effectively alleviate acetaminophen-induced liver damage and mitochondrial dysfunction. These studies have shown that GTPs possess antioxidant capacity and can prevent liver mitochondrial damage. Specifically, GTPs have been found to enhance the membrane potential and activity of liver mitochondrial respiratory chain complexes, thereby protecting against mitochondrial dysfunction. However, further research is needed to fully elucidate the underlying mechanisms and to explore the clinical applications of GTPs in protecting liver microstructures from DEHP-induced damage.

In this study, we aimed to analyze the signaling pathways involved in DEHP-induced liver damage and the hepatoprotective effects of GTPs using high-throughput sequencing. Ferroptosis, a recently discovered non-apoptotic cell death process, is induced by intracellular iron-dependent lipid peroxidation damage. Consistent with the findings of this study, Yin *et al* [36] investigated the acute toxicity of DEHP exposure in marine medaka. They conducted transcriptome analysis on the liver of DEHP-exposed medaka and reported that females were more sensitive to the immune response than males under acute DEHP exposure conditions. Furthermore, they found that DEHP exposure promoted iron depletion, leading to iron overload, increased levels of MDA and lipid peroxidation, and decreased levels of glutathione. These findings suggest that DEHP can rapidly alter certain molecular regulatory patterns and induce cell death through ferroptosis. In addition, we discussed the PPAR signaling pathway, which is a classic pathway associated with hyperlipidemia, regulates lipid metabolism and blood lipid levels by mediating various biological functions, such as the synthesis and decomposition of cholesterol and the oxidation of fatty acids [37]. Previous studies [38, 39] have revealed that DEHP exposure dysregulates blood lipid levels and hepatic lipid metabolism in mice through the PPAR signaling pathway.

In this study, we utilized the CIBERSORT algorithm to identify specific alterations in immune cells. Consistent with the findings of this study, Yang *et al* [40] investigated the

effect of DEHP on the immune system of male C57Bl/6 mice and reported significant atrophy of the thymus and spleen with the proportions of immature CD4⁺ and CD8⁺ populations exhibiting the highest downregulation. Additionally, they found a downregulation in the number of T and B cells in the spleen. A previous study [41] reported that green tea extract promotes macrophage phagocytic activity at a dosage of 5 mg/kg bodyweight and enhances NK cell activity and T cell proliferation at a dosage of 40 mg/kg bodyweight, supporting the potential immunomodulatory effects of GTPs demonstrated in this study.

This study identified the mmu-miR-141-3p/*Zcchc24* (mRNA)/*Zcchc24* (protein) regulatory axis, which exerted its function through *Zcchc24*. *Zcchc24* is reported to be involved in cell differentiation in mice. Previous studies [42] have suggested that *Zcchc24* can serve as a key gene in liver cancer prediction models, which are used to predict patient survival time. Additionally, *Zcchc24* is one of the major selective splicing factors that plays a critical role in cell fate transition, development, and disease progression.

This study has several limitations. In this study, three samples per group were analyzed using high-throughput sequencing, which may have yielded biased results. Additionally, the levels of immune cells in each group were determined using the CIBERSORT algorithm. However, immune cell infiltration was not verified using alternate methods, such as single-cell sequencing. Furthermore, the importance of *Zcchc24* was not verified using knockdown or overexpression experiments.

CONCLUSION

In conclusion, this study demonstrated that GTPs can alleviate the DEHP-induced changes in liver function and lipid profiles, improve fatty liver disease and partial liver fibrosis, and regulate immune cell infiltration. Additionally, an important miRNA-mRNA-protein molecular regulation axis that plays a crucial role in the therapeutic effects of GTPs on DEHP-induced liver damage was validated.

ARTICLE HIGHLIGHTS

Research background

5

Di (2-ethylhexyl) phthalate (DEHP) is a widely used plasticizer that has been shown to cause liver injury. Previous studies have reported the therapeutic effects of green tea on organ damage caused by heavy metal exposure. However, there is limited research on the therapeutic effects of green tea polyphenols (GTPs) specifically on DEHP-induced liver damage

Research motivation

Despite the known therapeutic effects of green tea on heavy metal exposure-induced organ damage, there is a lack of studies investigating the specific therapeutic effects of green tea polyphenols (GTPs) on DEHP-induced liver damage.

Research objectives

The research objectives of this study were to evaluate the molecular mechanism underlying the therapeutic effects of green tea polyphenols (GTPs) on DEHP-induced liver damage.

Research methods

In this study, C57BL/6J mice were divided into different groups and treated with DEHP and GTPs. After 8 wk, various assessments were conducted, including examination of liver function, blood lipid profile, and liver histopathology. High-throughput sequencing was used to analyze differentially expressed miRNAs and mRNAs in the liver tissues. Functional enrichment analysis and immune infiltration prediction were performed, and the miRNA-mRNA regulatory axis was elucidated using the starBase database. Protein expression was evaluated using immunohistochemistry.

Research results

The results of this study showed that GTPs had beneficial effects on DEHP-induced liver damage in mice. GTPs alleviated liver dysfunction, blood lipid dysregulation, fatty liver disease, liver fibrosis, and mitochondrial and endoplasmic reticulum lesions. The infiltration of immune cells, such as macrophages, mast cells, and natural killer cells, varied between the model and treatment groups. Furthermore, the study identified specific miRNAs, mRNAs, and proteins that constituted a regulatory axis involved in mediating the therapeutic effects of GTPs on DEHP-induced liver damage

Research conclusions

The findings of this study indicate that GTPs have a therapeutic effect on DEHP-induced liver damage. GTPs were shown to alleviate liver dysfunction, blood lipid dysregulation, fatty liver disease, and partial liver fibrosis. Additionally, GTPs were found to regulate immune cell infiltration. The study also identified a significant miRNA-mRNA-protein regulatory axis involved in mediating the therapeutic effects of GTPs on DEHP-induced liver damage

Research perspectives

Further studies are needed to investigate the long-term effects of GTPs on DEHP-induced liver damage and to explore the potential mechanisms underlying the regulation of immune cell infiltration. Additionally, future research should focus on optimizing the dosage and administration of GTPs to maximize their therapeutic effects and minimize potential side effects. Furthermore, clinical trials are warranted to evaluate the efficacy and safety of GTPs as a potential therapeutic intervention for DEHP-induced liver damage in humans.

7

ACKNOWLEDGEMENTS

We thank Bullet Edits Limited for the linguistic editing and proofreading of the manuscript. We also thank Figdraw (www.figdraw.com) for the materials provided in the scientific research drawing.

7%

SIMILARITY INDEX

PRIMARY SOURCES

- 1 www.ncbi.nlm.nih.gov 90 words — 2%
Internet
- 2 Yuanyuan Huang, Chuancheng Wu, Youbin Ye, Jingwen Zeng et al. "The Increase of ROS Caused by the Interference of DEHP with JNK/p38/p53 Pathway as the Reason for Hepatotoxicity", International Journal of Environmental Research and Public Health, 2019 77 words — 1%
Crossref
- 3 Leyang Li, Su Chen, Hiroyuki Yokoyama, Hidetoshi Kaburagi et al. "Remodeling of Neuromuscular Junctions in Target Muscle Following Nerve Regeneration in Mice After Delayed Peripheral Nerve Repair", Neuroscience, 2023 31 words — 1%
Crossref
- 4 www.spandidos-publications.com 26 words — < 1%
Internet
- 5 www.science.gov 25 words — < 1%
Internet
- 6 Le T Vuong, Stephen R Dueker, Suzanne P Murphy. "Plasma β -carotene and retinol concentrations of children increase after a 30-d supplementation with the fruit *Momordica cochinchinensis* (gac)", The American Journal of Clinical Nutrition, 2002 18 words — < 1%

-
- 7 Yufei Zhou, Chunjiang Liu, Zhongzheng Zhang, Jian Chen, Di Zhao, Linnan Li, Mingyue Tong, Gang Zhang. "Identification and validation of diagnostic biomarkers of coronary artery disease progression in type 1 diabetes via integrated computational and bioinformatics strategies", Computers in Biology and Medicine, 2023
Crossref 18 words — < 1%
-
- 8 f6publishing.blob.core.windows.net
Internet 18 words — < 1%
-
- 9 Yuhao Sun, Zhenzhen Wang, Jing Dai, Ruyi Sha, Jianwei Mao, Yangchen Mao, Yanli Cui. "Improved Antioxidant Capacity of Akebia trifoliata Fruit Inoculated Fermentation by Plantilactobacillus plantarum, Mechanism of Anti-Oxidative Stress through Network Pharmacology, Molecular Docking and Experiment Validation by HepG2 Cells", Fermentation, 2023
Crossref 17 words — < 1%
-
- 10 Ema Onode, Takuya Uemura, Kiyohito Takamatsu, Takuya Yokoi et al. "Bioabsorbable nerve conduits three-dimensionally coated with human induced pluripotent stem cell-derived neural stem/progenitor cells promote peripheral nerve regeneration in rats", Scientific Reports, 2021
Crossref 16 words — < 1%
-
- 11 Q. Yang. "Effects of peroxisome proliferators on the thymus and spleen of mice", Clinical & Experimental Immunology, 11/2000
Crossref 16 words — < 1%
-
- 12 assets.researchsquare.com
Internet 15 words — < 1%
-

13	"Abstract", Hepatology International, 2008 Crossref	14 words — < 1%
14	www.tandfonline.com Internet	13 words — < 1%
15	link.springer.com Internet	12 words — < 1%

EXCLUDE QUOTES	ON	EXCLUDE SOURCES	< 12 WORDS
EXCLUDE BIBLIOGRAPHY	ON	EXCLUDE MATCHES	< 12 WORDS