

Accuracy of early detection of colorectal tumours using stool methylation markers: A meta-analysis

Hu Zhang, You-Qing Zhu, Jian Qi, Qi-Xian Wang, Sha-Sha Cai, Si-Ying Zhu, Xiao-Wen Zhu, Xin-Tao Wang

Hu Zhang, You-Qing Zhu, Jian Qi, Qi-Xian Wang, Sha-Sha Cai, Si-Ying Zhu, Xiao-Wen Zhu, Xin-Tao Wang, Department of Gastroenterology and Clinical Center of Intestinal and Colorectal Diseases of Hubei Province, Zhongnan Hospital of Wuhan University, Wuhan 430071, Hubei Province, China

Author contributions: Qi J and Zhu YQ designed this study and critically revised the article; Zhang H and Qi J were responsible for data acquisition and extracted the data; Zhang H drafted the manuscript, analyzed the data and interpreted the results; Wang QX, Cai SS, Zhu SY, Zhu XW and Wang XT were involved in editing the manuscript; all authors read and approved the final manuscript to be published.

Supported by The National Natural Science Foundation of China Youth Science Fund, No. 81101868; The Natural Science Foundation of Hubei Province of China, No. 2011CDB505

Correspondence to: Dr. Jian Qi, Department of Gastroenterology and Clinical Center of Intestinal and Colorectal Diseases of Hubei Province, Zhongnan Hospital of Wuhan University, 169 Wuchang Donghu Lu, Wuhan 430071, Hubei Province, China. qiqidelizi@aliyun.com

Telephone: +86-27-87336141 Fax: +86-27-87336141

Received: July 9, 2013 Revised: October 21, 2013

Accepted: November 2, 2013

Published online: November 26, 2013

Abstract

AIM: To evaluate the accuracy of methylation of genes in stool samples for diagnosing colorectal tumours.

METHODS: Electronic databases including PubMed, Web of Science, Chinese Journal Full Text Database and Wanfang Journals Full-text Database were searched to find relevant original articles about methylated genes used in diagnosing colorectal tumours. Quality assessment of diagnostic accuracy studies items were used to evaluate the quality of the included articles, and the Meta-disc 1.4 and SPSS 13.0 software programs were used for data analysis.

RESULTS: Thirty-four articles met the inclusion criteria, and 4151 patients were included. Pooled diagnostic

performances of SFRP2 methylation for colorectal cancer (CRC) provided the following results: the sensitivity was 79% (95%CI: 75%-82%), the specificity was 93% (95%CI: 90%-96%), the diagnostic odds ratio (DOR) was 47.57 (95%CI: 20.08-112.72), and the area under the curve was 0.9565. Additionally, the results of accuracy of SFRP2 methylation for detecting colorectal adenomas were as follows: the sensitivity was 43% (95%CI: 38%-49%), the specificity was 94% (95%CI: 91%-97%), the DOR was 11.06 (95%CI: 5.77-21.18), and the area under the curve was 0.9563.

CONCLUSION: Stool-based DNA testing may be useful for non-invasively diagnosing colorectal tumours, and SFRP2 methylation is a promising marker that has great potential in early CRC diagnosis.

© 2013 Baishideng Publishing Group Co., Limited. All rights reserved.

Key words: Colorectal carcinoma; Colorectal adenoma; Stool; Methylation; Meta-analysis

Core tip: The analysis of stool methylation markers as a non-invasive test is important for the early diagnosis of colorectal tumours. However, no consensus has been reached with regard to the role of stool methylation markers in colorectal tumour diagnosis. We performed a meta-analysis of 34 articles, and the pooled results showed that stool methylation markers could be used as a valuable diagnostic and predictive tool for colorectal tumours and SFRP2 methylation serves as a promising marker with great potential in early colorectal cancer diagnosis.

Zhang H, Zhu YQ, Qi J, Wang QX, Cai SS, Zhu SY, Zhu XW, Wang XT. Accuracy of early detection of colorectal tumours using stool methylation markers: A meta-analysis. *World J Meta-Anal* 2013; 1(3): 147-156 Available from: URL: <http://www.wjgnet.com/2308-3840/full/v1/i3/147.htm> DOI: <http://dx.doi.org/10.13105/wjma.v1.i3.147>

INTRODUCTION

Colorectal cancer (CRC) is the third most common malignancy and the second leading cause of cancer-related deaths in Western countries^[1,2]. The 5-year survival of stage I CRC has reached 90%^[3], but less than 10% of CRC cases have distant metastases^[4]. However, most cases of CRC are diagnosed at the middle or late stage because no typical symptoms for early-stage CRC exist^[5]. Therefore, the diagnosis of CRC at early stages has great importance for reducing CRC mortality.

Early diagnosis of CRC will help reduce mortality and the costs for surgery. The current colonoscopy screening test is of high efficacy, but the acceptability of this procedure in the general public is rather low. As an available non-invasive method, faecal testing has a unique advantage when compared to other screening modalities. Although faecal occult blood testing (FOBT) has been confirmed to reduce mortality due to CRC, the test has little or no impact on the incidence of CRC because of its low-level sensitivity for the detection of adenoma^[6], *i.e.*, a sensitivity of only 10%-20%^[7]. Compared to FOBT, the most important advantage of methylation markers in stool samples is their higher accuracy and sensitivity for the diagnosis of premalignant lesions of CRC^[8].

DNA methylation often occurs in the early stage of CRC, and many studies have been performed on the diagnosis of colorectal tumours by determining the methylation of genes in stool samples. However, the results of these studies are variable although inspiring. Thus, this meta-analysis was conducted to assess the accuracy of the detection of colorectal tumours using methylation markers in stool samples.

MATERIALS AND METHODS

Search strategy

A literature search was performed independently by two investigators (Zhang H and Qi J) using the following databases: PubMed, Web of Science, Chinese Journal Full Text Database and Wanfang Journals Full-text Database. All references that were cited in these studies and all published reviews were also searched. All English and Chinese references for analyses were published before January 2013. The following keywords were used in the search strategy: "colon/rectal/colorectal", "cancer/tumours", "stool," and "methylation". In this meta-analysis, 2 × 2 tables were constructed from each study for the true-positive, false-negative, true-negative and false-positive values.

Inclusion and exclusion criteria

Eligible studies were required to meet all of the following criteria: (1) the data were independent; (2) CRC was diagnosed by analyzing DNA methylation in stool sample; (3) patients were diagnosed with colorectal cancer or colorectal adenomas by pathology; and (4) the colonoscopy result of control individuals was normal.

Exclusion criteria for this meta-analysis were as fol-

lows: (1) studies on secondary CRC or primary CRC with distant metastases and (2) studies on CRC patients receiving chemotherapy or curative surgery.

Data extraction and quality assessment

The following data were extracted from each study: author, year of publication, country or region, sample size, the name of genes, the detection method of methylation and the study design. The data were independently extracted by two investigators (Zhang H and Qi J), and discrepancies were solved by a third investigator (Zhu YQ) and collective discussion. Quality Assessment of studies of Diagnostic Accuracy^[9] (QUADAS) was used to assess the quality of the primary studies with diagnostic accuracy, and quality scoring was appraised based on the empirical evidence, the experts' opinions and the formal consensus. Score of 1, 0 or -1 was given to the articles that were in compliance with the standards completely, unclear or out of standards, respectively, and the full score was 14.

Statistical analysis

All statistics were calculated and then combined using a random-effects model and 95%CI as effect measurements. The diagnostic odds ratio (DOR) reflects the relationship between the result of the diagnostic test and the disease. The summary receiver operation characteristic (SROC) curve displays the trade-off between sensitivity and specificity and represents a global summary of test performance. We used the *Q*-value, which is the intersection point of the SROC curve with a diagonal line from the left upper corner to the right lower corner of the receiver operation characteristic (ROC) space, which corresponds to the highest value of sensitivity and specificity for the test. The positive likelihood ratio (PLR) represents the value by which the odds of the disease increase when a test is positive, whereas the negative likelihood ratio (NLR) shows the value by which the odds of the disease decrease when a test is negative. Statistical heterogeneity was assessed using the Chi-square test, and alpha significance testing was performed at the two-tailed 0.05 level. The professional statistical software programs (Meta-DiSc 1.4 and SPSS 13.0) were used for analysis. Publication bias was assessed by Egger analysis.

RESULTS

The literature search retrieved 453 citations, of which 344 were excluded because they were duplicates. Of the 109 potentially eligible studies, 75 publications were excluded because they did not investigate colorectal tumour studies ($n = 6$), included no diagnostic value studies ($n = 15$), were reviews ($n = 26$) or had overlapping data ($n = 28$). Finally, 34 studies that focused on the target patient spectrum were included (Figure 1).

Study characteristics

Of the 34 studies, 7 were Chinese and 27 were English,

Table 1 The characteristics of the included studies in the meta-analysis and QUADAS scores

Ref.	Country/ region	Methylated genes	N	CRC		Adenoma		Normal		Blind design	Detection method	QUADAS score
				+	-	+	-	+	-			
Ahlquist <i>et al</i> ^[10]	Ireland	<i>Vimentin</i> /NDRG4/BMP3/TFPI2	98	26	4	18	4	5	41	Yes	QuARTS	11
Bosch <i>et al</i> ^[11] , 2011	Netherlands	<i>PHACTR3</i>	185	40	25	6	13	4	97	Unclear	qMSP	10
		<i>GATA4</i>	160	29	11	3	16	6	95			
		<i>OSMR</i>	185	25	40	4	15	7	94			
		<i>PHACTR3</i>	639	214	38	51	43	29	264			
Ahlquist <i>et al</i> ^[12]	Ireland	<i>PHACTR3</i>	639	214	38	51	43	29	264	Yes	QuARTS	11
Azuara <i>et al</i> ^[13]	Spain	<i>RARB2/P16/MGMT/APC</i>	98	25	13	20	20	0	20	Yes	MS-MCA	10
		<i>RARB2</i>	85	11	23	7	31	0	13			
		<i>P16</i>	77	9	21	6	28	0	13			
		<i>MGMT</i>	80	9	19	3	34	0	15			
		<i>APC</i>	77	9	19	9	25	0	15			
Tang <i>et al</i> ^[14]	China	<i>SRRP2</i>	262	142	27	29	34	2	28	Yes	MSP	9
Baek <i>et al</i> ^[15]	South Korea	<i>Vimentin/MGMT/MLH1</i>	149	45	15	31	21	5	32	Yes	MSP	9
		<i>MLH1</i>	149	18	42	6	46	0	37			
		<i>Vimentin</i>	149	23	37	8	44	0	37			
		<i>MGMT</i>	149	31	29	19	33	5	32			
Li <i>et al</i> ^[16] , 2009	United States	<i>Vimentin</i>	80	9	13	9	11	2	36	Unclear	Methy-BEAMing	5
Melotte <i>et al</i> ^[17] , 2009	Netherlands	<i>NDRG4</i>	150	42	33	nr	nr	3	72	Yes	qMSP	11
Ausch <i>et al</i> ^[18] , 2009	United States	<i>IGTA4</i>	37	nr	nr	7	2	6	22	Unclear	qMSP	4
Hellebrekers <i>et al</i> ^[19] , 2009	Netherlands	<i>GATA4</i>	150	44	31	nr	nr	9	66	Yes	qMSP	10
Mayor <i>et al</i> ^[20] , 2009	Spain	<i>EN1</i>	60	8	22	nr	nr	1	29	Unclear	MS-MCA	7
Kim <i>et al</i> ^[21] , 2009	United States	<i>OSMR/SFRP1</i>	42	12	8	6	11	0	5	Yes	qMSP	9
		<i>OSMR</i>	201	35	54	2	14	4	92			
		<i>SFRP1</i>	52	11	9	5	12	0	15			
		<i>SFRP2</i>	253	53	31	18	38	9	104			
Nagasaka <i>et al</i> ^[22] , 2009	Japan	<i>RASSF2</i>	253	38	46	7	49	6	107	Unclear	COBRA	10
		<i>SFRP2</i>	253	38	46	7	49	6	107			
Glöckner <i>et al</i> ^[23] , 2009	United States	<i>TFPI2</i>	129	44	14	7	19	2	43	Yes	qMSP	12
Wang <i>et al</i> ^[24] , 2008	China	<i>SFRP2</i>	133	60	9	21	13	2	28	Yes	MethyLight	8
Oberwalder <i>et al</i> ^[25] , 2008	Australia	<i>SFRP2</i>	19	nr	nr	6	7	0	6	Yes	MethyLight	9
Izkowitz <i>et al</i> ^[26] , 2008	United States	<i>Vimentin</i>	80	9	13	9	11	2	36	Yes	MSP	13
Huang <i>et al</i> ^[27] , 2007	China	<i>SFRP2/HPP1/MGMT</i>	97	50	2	15	6	1	23	Yes	MSP	8
		<i>SFRP2</i>	97	49	3	11	10	1	23			
		<i>HPP1</i>	97	37	15	12	9	0	24			
		<i>MGMT</i>	97	25	27	6	15	0	24			
Izkowitz <i>et al</i> ^[28] , 2007	United States	<i>Vimentin/HLTF</i>	162	31	9	nr	nr	19	103	Yes	MSP	13
		<i>HLTF</i>	162	15	25	nr	nr	9	113			
		<i>Vimentin</i>	162	29	11	nr	nr	16	106			
Abbaszadegan <i>et al</i> ^[29] , 2007	Hong Kong	<i>p16</i>	45	5	20	nr	nr	0	20	Unclear	MSP	8
Zhang <i>et al</i> ^[30] , 2007	Germany	<i>SFRP1</i>	44	16	4	7	0	2	15	Yes	MSP	9
Leung <i>et al</i> ^[31] , 2007	Hong Kong	<i>SFRP2/MGMT/MLH1/HLTF/ATM/APC</i>	75	16	4	18	7	3	27	Yes	MSP	13
		<i>SFRP2</i>	75	6	14	3	22	2	28			
		<i>MGMT</i>	75	4	16	3	22	0	30			
		<i>MLH1</i>	75	4	16	3	22	0	30			
		<i>HLTF</i>	75	5	15	5	20	1	29			
		<i>ATM</i>	75	5	15	5	20	0	30			
		<i>APC</i>	75	4	16	4	21	0	30			
		<i>MGMT/CDKN2A/MLH1</i>	48	nr	nr	16	13	7	12			
		<i>CDKN2A</i>	48	nr	nr	9	20	3	16			
<i>MGMT</i>	48	nr	nr	14	15	5	14					
<i>MLH1</i>	48	nr	nr	0	29	2	17					
Lenhard <i>et al</i> ^[33] , 2005	Germany	<i>HIC1</i>	71	11	15	4	9	0	32	Yes	MSP	11
Chen <i>et al</i> ^[34] , 2005	United States	<i>Vimentin</i>	263	43	51	6	44	8	111	Yes	MSP	11
Müller <i>et al</i> ^[35] , 2004	Australia	<i>SFRP2/SRRP5</i>	39	20	3	nr	nr	8	8	Unclear	MethyLight	5
		<i>SRRP2</i>	39	19	4	nr	nr	4	12			
		<i>SRRP5</i>	39	18	5	nr	nr	5	11			
		<i>SFRP2</i>	90	20	10	15	15	1	29			
Xu <i>et al</i> ^[36] , 2012	China	<i>SFRP2</i>	90	20	10	15	15	1	29	Unclear	MSP	5
Kang <i>et al</i> ^[37] , 2011	China	<i>MGMT/MAL/CDKN2A</i>	119	64	5	17	7	2	24	Unclear	MSP	7
		<i>MAL</i>	119	54	15	14	10	1	25			
		<i>CDKN2A</i>	119	36	33	10	14	0	26			
		<i>MGMT</i>	119	38	31	9	15	1	25			
Zhang <i>et al</i> ^[38] , 2011	China	<i>Vimentin/OSMR/TFPI2</i>	107	52	8	13	4	4	26	Unclear	MSP	9
		<i>Vimentin</i>	107	32	28	5	12	0	30			
		<i>OSMR</i>	107	41	19	7	10	0	30			
		<i>TFPI2</i>	107	45	15	11	6	4	26			

Fu <i>et al</i> ^[39] , 2010	China	<i>Vimentin</i>	22	5	9	nr	nr	0	8	Unclear	MSP	5
Ling <i>et al</i> ^[40] , 2009	China	<i>P16</i>	108	47	14	16	11	1	19	Unclear	MSP	7
Cheng <i>et al</i> ^[41] , 2007	China	<i>SFRP2</i>	97	49	3	11	10	1	23	Unclear	MSP	5
Zhao <i>et al</i> ^[42] , 2009	China	<i>NDRG4</i>	114	64	20	nr	nr	3	27	Unclear	MSP	6
Chang <i>et al</i> ^[43] , 2010	South Korea	<i>IGTA4/SFRP2/P16</i>	86	21	9	18	7	1	30	Yes	MSP	8
		<i>IGTA4</i>	86	11	19	4	21	0	31			
		<i>SFRP2</i>	86	18	12	11	14	0	31			
		<i>P16</i>	86	12	18	6	19	1	30			

+: Represents the number of individuals with the disease when the DNA methylation test was positive; -: Represents the number of individuals with the disease when the DNA methylation test was negative; nr: Not reported; N: Total number. CRC: Colorectal cancer.




Figure 1 Flowchart of the study selection.

and they included 4151 patients (Table 1). These studies were performed in 10 countries or regions (including China, the United States, the Netherlands, Spain, Japan, Germany, Iran, Hong Kong, Austria and South Korea). In these studies, 31 evaluated CRC, and 26 evaluated colorectal adenoma. Twenty-two studies focused on the methylation of a single gene, and the other 12 studies involved the methylation of multiple genes.

Genes evaluated in these studies are mainly involved in three types of regulation pathways: the Wnt pathway, the DNA damage repair pathway and other pathways. Four genes of the Wnt pathway were involved in 10 studies: secreted frizzled-related proteins (*SFRP1*, *SFRP2*, *SFRP5*) and adenomatous polyposis coli (*APC*). Two genes of the DNA damage repair pathway were involved in 7 studies: O-6-methylguanine-DNA methyltransferase (*MGMT*) and Mut L homologue 1 (*MLH1*). Twenty-seven studies involved 16 genes of other pathways: vimentin, oncostain M receptor- β (*OSMR*), phosphatase and actin regulator 3 (*PHACTR3*), cyclin-dependent kinase inhibitor 2A (*CDKN2A*), tissue factor pathway inhibitor 2 (*TFPI2*), hyperplastic polyposis protein gene (*HPP1*), GATA4, human lactoferrin (*HLTF*), *ATM*, ras association domain family 2 (*RASSF2*), *RARB2*, hypermethylated in cancer 1 (*HIC*), engrailed gene (*ENT1*), N-myc downstream-regulated gene family (*NDRG4*), *IGTA4* and T-cell differentiation protein (*MAL*).

Qualitative and quantitative methods were the two main types of methods used for methylation detection.

The qualitative method included methylation-specific PCR (MSP) and methylation-specific melting curve analysis (MS-MCA). The quantitative method included MethL-BEAMing; quantitative MSP (qMSP); MethyLight; combined bisulfite restriction analysis (COBRA); and quantitative, allele-specific, real-time target and signal amplification (QuARTS).

Colorectal carcinoma meta-analysis

The colorectal carcinoma results were pooled from 31 studies and are shown in Table 2. The meta-analysis showed that the sensitivity and specificity of gene methylation for the detection of colorectal carcinoma were 73% (95%CI: 71%-75%) and 92% (95%CI: 90%-93%), respectively. The PLR was 7.94 (95%CI: 6.08-10.36), the NLR was 0.31 (95%CI: 0.25-0.39), the DOR was 30.86 (95%CI: 22.33-42.66), and the symmetric area under the curve was 0.9286.

Heterogeneity was significant for the sensitivity ($P < 0.001$), specificity ($P = 0.0012$), PLR ($P = 0.0023$), NLR ($P < 0.001$), and DOR ($P = 0.0245$).

Of the involved regulation mechanisms, we found that DOR and AUC of the methylated genes belonging to the Wnt pathway were higher than the genes of the DNA damage repair pathway and other pathways. The sensitivity, specificity, DOR and AUC of different methylated genes in the three types of pathways were calculated (Table 2), and the results indicated that the accuracy of faecal SFRP2 methylation in the diagnosis of colorectal carcinoma was higher than that of other genes, with a sensitivity of 79% (95%CI: 75%-82%) (Figure 2A), a specificity of 93% (95%CI: 90%-96%) (Figure 2B), a DOR of 47.57 (95%CI: 20.08-112.72), and the area under the curve of 0.9565 (Figure 2C).

Colorectal adenoma meta-analysis

Pooled colorectal adenoma analysis (Table 3), including 26 studies, provided the following results: the sensitivity and specificity of gene methylation for colorectal adenoma diagnosis were 51% (95%CI: 47%-54%) and 92% (95%CI: 90%-93%), respectively. The PLR was 5.52 (95%CI: 4.23-7.19), the NLR was 0.52 (95%CI: 0.44-0.61), and the DOR and symmetric area under the curve were 12.61 (95%CI: 8.66-18.37) and 0.8830, respectively.

Heterogeneity was also significant regarding sensitivity ($P < 0.001$), specificity ($P = 0.0233$), PLR ($P = 0.1166$), NLR ($P < 0.001$), and DOR ($P = 0.0565$).

Table 2 Methylation of pooled genes for the diagnosis of colorectal cancer

Wnt pathway	DNA damage repair pathway	Other pathways	SE (95%CI)	SP (95%CI)	DOR (95%CI)	AUC
Wnt pathway	DNA damage repair pathway	Other pathways	73% (0.71-0.75)	92% (90%-93%)	30.86 (22.33-42.66)	0.929
Wnt pathway	-	-	74% (70%-77%)	93% (90%-95%)	33.92 (17.73-64.90)	0.932
-	DNA damage repair pathway	-	42% (36%-47%)	97% (94%-99%)	12.87 (5.98-27.72)	0.730
-	-	Other pathways	57% (55%-60%)	94% (93%-95%)	20.93 (15.56-28.15)	0.921
SFRP2	-	-	79% (75%-82%)	93% (90%-96%)	47.57 (20.08-112.72)	0.957
-	MGMT	-	47% (40%-53%)	95% (90%-98%)	11.67 (5.10-26.67)	0.709
-	MLH	-	28% (18%-39%)	100% (95%-100%)	23.68 (3.02-185.44)	0.500
-	-	Vimentin	48% (42%-54%)	93% (90%-95%)	14.95 (8.99-24.84)	0.862
-	-	OSMR	47% (40%-54%)	95% (91%-98%)	14.66 (5.06-42.47)	0.225
-	-	P16	50% (42%-58%)	98% (92%-100%)	24.39 (7.26-81.96)	0.975
SFRP2	MGMT	-	69% (66%-72%)	94% (91%-96%)	33.24 (16.76-65.93)	0.946
SFRP2	MLH	-	72% (68%-75%)	94% (92%-96%)	43.03 (20.15-91.87)	0.953
SFRP2	MLH	Vimentin	64% (61%-67%)	94% (92%-95%)	27.11 (16.48-44.61)	0.934
SFRP2	MLH	OSMR	65% (62%-69%)	95% (93%-96%)	33.10 (17.12-63.98)	0.951
SFRP2	MLH	P16	68% (64%-71%)	95% (93%-97%)	38.86 (20.11-67.54)	0.952

SE: Sensitivity; SP: Specificity; DOR: Diagnostic odds ratio; AUC: The area under the curve.




Figure 2 Forest plot of SFRP2 methylation in the diagnosis of colorectal cancer. A: The sensitivity of SFRP2 methylation in stool samples used for colorectal carcinoma diagnosis. The point estimates of specificity from each study are shown as red squares; B: The specificity of SFRP2 methylation in stool samples used for colorectal cancer diagnosis. The point estimates of specificity from each study are shown as blue squares; C: The summary receiver operating characteristic curves of SFRP2 methylation assays used for diagnosis of colorectal carcinoma. Red circles represent each study that was included in the meta-analysis. The size of each study is indicated by the size of the red circle. Summary receiver operating characteristic curves summarize the overall diagnostic accuracy. Error bars indicate the 95%CI, and df indicates the degrees of freedom.

The DOR and AUC of the methylated Wnt pathway genes were higher than those of the genes of the DNA

damage repair pathway and other pathways when grouping all of the genes by pathway for analysis. In these

Table 3 Methylation of pooled genes for the diagnosis of colorectal adenomas

Wnt pathway	DNA damage repair pathway	Other pathways	SE (95%CI)	SP (95%CI)	DOR (95%CI)	AUC
Wnt pathway	DNA damage repair pathway	Other pathways	51% (47%-54%)	92% (90%-93%)	12.61 (8.66-18.37)	0.883
Wnt pathway	-	-	40% (35%-46%)	95% (92%-97%)	10.81 (6.43-18.16)	0.932
-	DNA damage repair pathway	-	21% (17%-27%)	95% (91%-97%)	4.23 (2.01-8.88)	0.672
-	-	Other pathways	32% (28%-35%)	94% (93%-95%)	7.78 (5.48-11.05)	0.873
SFRP2	-	-	43% (38%-49%)	94% (91%-97%)	11.06 (5.77-21.18)	0.956
-	MGMT	-	29% (22%-36%)	93% (87%-96%)	4.42 (2.18-8.95)	0.614
-	MLH	-	8% (4%-16%)	98% (92%-100%)	2.35 (0.14-40.83)	-
-	-	Vimentin	23% (17%-31%)	95% (92%-98%)	8.30 (2.60-26.55)	0.898
-	-	OSMR	25% (14%-39%)	95% (91%-98%)	5.20 (1.44-18.82)	0.817
-	-	P16	33% (23%-44%)	97% (89%-100%)	13.27 (3.40-51.83)	0.97
SFRP2	MLH	-	34% (29%-39%)	95% (92%-97%)	9.62 (4.64-19.93)	0.947
SFRP2	MGMT	-	38% (33%-42%)	94% (91%-96%)	7.85 (4.79-12.87)	0.753
SFRP2	-	OSMR	41% (35%-46%)	95% (92%-96%)	9.25 (5.13-16.69)	0.948
SFRP2	-	Vimentin	36% (32%-41%)	95% (93%-96%)	9.88 (5.55-17.57)	0.946
SFRP2	-	P16	41% (36%-46%)	95% (92%-97%)	10.37 (6.21-17.31)	0.948
SFRP2	MGMT	Vimentin	34% (30%-38%)	94% (92%-96%)	7.81 (4.96-12.29)	0.804
SFRP2	MGMT	OSMR	36% (32%-41%)	94% (92%-96%)	7.25 (4.61-11.39)	0.775
SFRP2	MGMT	P16	37% (33%-41%)	94% (92%-96%)	7.92 (5.14-12.21)	0.772
SFRP2	MLH	Vimentin	31% (27%-35%)	95% (93%-97%)	8.99 (4.95-16.31)	0.944
SFRP2	MLH	OSMR	33% (29%-38%)	95% (93%-97%)	8.37 (4.50-15.59)	0.941
SFRP2	MLH	P16	34% (30%-38%)	95% (93%-97%)	9.98 (5.45-18.27)	0.947

SE: Sensitivity; SP: Specificity; DOR: Diagnostic odds ratio; AUC: The area under the curve.

regulation mechanisms, we also found that the sensitivity and specificity of the methylated genes in the Wnt pathway were higher than those in the DNA damage repair pathway and the other pathway. The sensitivity, specificity, DOR and AUC of the different methylated genes in the three types of pathways were calculated (Table 3), and the results indicated that the values of DOR and AUC of P16 and SFRP2 were higher than those of other genes, but the accuracy of faecal SFRP2 methylation for the diagnosis of colorectal adenoma was higher than that of P16 methylation according to sensitivity (Figure 3).

Meta-regression

In the meta-regression analysis, the difference in relative diagnostic odds ratio (RDOR) values between the higher and lower quality studies was not significant. We also noted that the differences between the blinded and non-blinded methods, qualitative and quantitative methods, single- and multiple-gene methylation did not reach statistical significance, indicating that these potential factors did not substantially affect the diagnostic accuracy, as shown in Table 4.

Publication bias

In our meta-analysis, publication bias was evaluated using the Egger test. The results showed no significant publication bias among the studies of SFRP2 methylation in faecal samples from CRC or adenoma patients (Figure 4).

DISCUSSION

It is widely accepted that DNA methylation in stool samples may be valuable for increasing the rate of CRC detection at earlier stage^[44]. In the present study, we fo-

cused on the detection performance of gene methylation in stool samples for patients with colorectal tumours. Our analysis suggests that the specificity of SFRP2 methylation is high (93% for CRC and 94% for colorectal adenoma) for the detection of colorectal tumours; however, it has moderate (79%) and low sensitivity (43%) for diagnosing CRC and adenoma, respectively. Compared to FOBT, with a sensitivity of 14% for colorectal tumour diagnosis^[45], the detection accuracy of faecal methylation biomarkers was higher as a CRC screening method.

The DOR is an indicator of test accuracy. The value of the DOR ranges from 0 to infinity, and higher values indicate better discriminatory test performance. In this meta-analysis, we found that the DOR of faecal SFRP2 methylation for colorectal carcinoma and adenoma were 47.57 and 11.06, respectively, which indicated a high level of overall accuracy for CRC and a low level for adenoma. The SROC curve represents an overall measure of the discriminatory power of a test. The area under the curve of 1 for any test indicates that the test is excellent. Our data showed that the area under the curve (AUC) of the SROC curve for faecal SFRP2 methylation for the diagnosis of colorectal carcinoma and adenoma were 0.9565 and 0.9563, respectively, which indicated that faecal SFRP2 methylation is an excellent diagnostic biomarker for colorectal tumours.

Because the DOR and SROC curve are not easy to use in clinical practice, the likelihood ratios are considered to be more clinically meaningful. For a high-quality diagnostic test, a PLR of > 10 or an NLR < 0.1 is typically required. However, our meta-analysis showed that neither PLR nor NLR alone was adequate to confirm or exclude the diagnosis of colorectal carcinoma or adenoma. The PLR value was 9.12 in the diagnosis analysis of




Figure 3 Forest plot of SFRP2 methylation in the diagnosis of colorectal adenomas. A: Plot and table of the sensitivity of SFRP2 for diagnosis of colorectal adenomas; B: Plot and table of the specificity of SFRP2 for diagnosis of colorectal adenomas; C: The symmetric summary receiver-operating characteristic of SFRP2 for diagnosis of colorectal adenomas.




Figure 4 Assessment of the publication bias in the faecal SFRP2 methylation for the diagnosis of colorectal cancer (A) and adenomas (B). No significant publication biases were found in any of these studies (all $P > 0.05$).

CRC, which suggested that patients with a positive faecal SFRP2 methylation assay had a nine-fold chance of being diagnosed with CRC than non-CRC. Therefore, a colonoscopy was necessary for patients with a positive faecal SFRP2 methylation assay to confirm the diagnosis of

CRC with high probability. On the other hand, an NLR of 0.24 in the diagnosis analysis of CRC suggested that if a faecal SFRP2 methylation assay result was negative, the probability rate of the individual having CRC was 24%. For the diagnosis of colorectal adenoma, a PLR of 5.99

Table 4 Weighted meta-regression on the diagnostic accuracy of the gene methylation assays

Covariate	Coefficient	SE	P value	RDOR	95%CI
QUADAS score ¹	-0.259	0.419	0.541	0.77	(0.33, 1.82)
Detection method ²	0.101	0.377	0.790	1.11	(0.51, 2.40)
Blinded design ³	-0.179	0.381	0.642	0.84	(0.38, 1.82)
Methylated genes ⁴	0.302	0.394	0.449	1.35	(0.60, 3.03)

¹Based on QUADAS score, studies were divided into those with higher quality (QUADAS score ≥ 10) and those with lower quality (QUADAS score < 10); ²Detection method, which was divided into qualitative and quantitative assay methods; ³Blinded design: the study was included with or without blinded design; ⁴Methylated genes, which were divided into single gene and combination genes. QUADAS: Quality Assessment for Studies of Diagnostic Accuracy was used to assess the quality of primary studies of diagnostic accuracy; SE: Standard error; RDOR: Relative diagnostic odds ratio.

suggested a moderate necessity to consider colonoscopy for patients with a positive faecal SFRP2 methylation assay to confirm the diagnosis of colorectal adenoma. Moreover, the NLR was 0.60 in the diagnosis analysis of colorectal adenoma. These data suggest that a negative faecal SFRP2 methylation assay result should not be used alone as a justification for denying or discontinuing the screening of colorectal adenomas.

An aberrant Wnt signalling pathway is an early event in 90% of colorectal carcinomas. SFRPs are secreted glycoproteins that antagonise Wnt signaling *via* different direct or indirect mechanisms. Thus, the role of SFRPs as a negative regulator of Wnt signaling may have important significance in tumourigenesis. These epigenetic events are involved in early steps of colon carcinogenesis, and changes in the status of DNA methylation are associated with early steps of the histologic progression of colon carcinoma. Our previous studies of CRC tissue showed that SFRP1 and SFRP2 were methylated in more than 80.6% of colorectal carcinomas^[46]. Therefore, faecal SFRP2 methylation could be expected to be a biomarker for the screening of colorectal tumours. Although it cannot be generally used as a screening tool for the financial limited, the analysis of methylation markers offers a variety of new opportunities for developing biomarkers for colorectal tumours at the molecular level.

Our meta-analysis had several limitations. First, none of the included studies were multicentre or large-blinded, randomized, controlled trials. Second, conference abstracts and non-English and non-Chinese language studies were excluded, which might have led to publication bias. Third, studies on DNA methylation with statistical significance tend to be published and cited. Finally, due to the absence of case-mix difference analysis, smaller trials may show larger treatment effects than larger studies (*e.g.*, patients with only localised *vs* metastatic disease).

To sum up, stool-based DNA methylation has been shown to be highly discriminatory in the detection of colorectal tumours. Our results demonstrate that SFRP2 methylation, as a non-invasive modality, shows promise for the accurate detection of CRC; however, a large num-

ber of studies are required to further confirm the role of faecal SFRP2 methylation for the early and accurate CRC diagnosis.

COMMENTS

Background

Colorectal cancer (CRC) is the third most common malignancy and the second leading cause of cancer-related deaths in Western countries. The diagnosis of CRC at early stages has great importance for reducing CRC mortality. Although significant advances have been achieved in diagnostic technologies, the current available modalities for diagnosing CRC remain suboptimal.

Research frontiers

DNA methylation often occurs during the early stages of colon tumours and has played an important role in oncology, especially in the early diagnosis of colorectal tumours. However, no consensus with regard to the role of stool methylation markers in colon tumours exists.

Innovations and breakthroughs

Stool methylation markers as an available non-invasive modality have high accuracy and sensitivity for the diagnosis of premalignant lesions of CRC. A few systematic reviews about the efficacy of stool methylation markers in colorectal tumour diagnosis exist. This article comprehensively assesses the accuracy of methylated genes in stool samples for diagnosing colorectal tumours.

Applications

Analysis of DNA methylation in stool samples may be used as a non-invasive test for the diagnosis of CRC, and SFRP2 methylation is a promising marker that has great potential in early CRC diagnosis.

Terminology

Diagnostic odds ratio (DOR) reflects the relationship between the result of the diagnostic test and the disease. The summary receiver operation characteristic (SROC) curve displays the trade-off between sensitivity and specificity and represents a global summary of test performance. We used the Q-value, the intersection point of the SROC curve with a diagonal line from the left upper corner to the right lower corner of the receiver operation characteristic (ROC) space, which corresponds to the highest value of sensitivity and specificity for the test. The positive likelihood ratio (PLR) represents the value by which the odds of the disease increase when a test is positive, whereas negative likelihood ratio (NLR) shows the value by which the odds of the disease decrease when a test is negative.

Peer review

This study reviewed 34 trials to evaluate the accuracy of stool methylation genes for diagnosing colorectal tumours. Based on these analyses, the authors conclude that stool SFRP2 methylation is a promising marker that has great potential in early CRC diagnosis. The analysis was carefully performed, the results were clearly presented and summarized, and valuable advice for early clinical diagnosis of colorectal tumours was provided.

REFERENCES

- 1 **Jemal A**, Siegel R, Ward E, Hao Y, Xu J, Thun MJ. Cancer statistics, 2009. *CA Cancer J Clin* 2009; **59**: 225-249 [PMID: 19474385 DOI: 10.3322/caac.20006]
- 2 **Schetter AJ**, Harris CC. Alterations of microRNAs contribute to colon carcinogenesis. *Semin Oncol* 2011; **38**: 734-742 [PMID: 22082759 DOI: 10.1053/j.seminoncol.2011.08.009]
- 3 **Markowitz SD**, Dawson DM, Willis J, Willson JK. Focus on colon cancer. *Cancer Cell* 2002; **1**: 233-236 [PMID: 12086859 DOI: 10.1016/S1535-6108(02)00053-3]
- 4 **Bosch LJ**, Carvalho B, Fijneman RJ, Jimenez CR, Pinedo HM, van Engeland M, Meijer GA. Molecular tests for colorectal cancer screening. *Clin Colorectal Cancer* 2011; **10**: 8-23 [PMID: 21609931 DOI: 10.3816/CCC.2011.n.002]
- 5 **Jemal A**, Siegel R, Ward E, Hao Y, Xu J, Murray T, Thun MJ. Cancer statistics, 2008. *CA Cancer J Clin* 2008; **58**: 71-96 [PMID: 18287387 DOI: 10.3322/CA.2007.0010]
- 6 **Heresbach D**, Manfredi S, D'halluin PN, Bretagne JF, Branger B. Review in depth and meta-analysis of controlled trials

- on colorectal cancer screening by faecal occult blood test. *Eur J Gastroenterol Hepatol* 2006; **18**: 427-433 [PMID: 16538116 DOI: 10.1097/00042737-200604000-00018]
- 7 **Tagore KS**, Levin TR, Lawson MJ. The evolution to stool DNA testing for colorectal cancer. *Aliment Pharmacol Ther* 2004; **19**: 1225-1233 [PMID: 15191503 DOI: 10.1111/j.1365-2036.2004.02005.x]
 - 8 **Berger BM**, Ahlquist DA. Stool DNA screening for colorectal neoplasia: biological and technical basis for high detection rates. *Pathology* 2012; **44**: 80-88 [PMID: 22198259 DOI: 10.1097/PAT.0b013e3283502fd]
 - 9 **Whiting P**, Rutjes AW, Reitsma JB, Bossuyt PM, Kleijnen J. The development of QUADAS: a tool for the quality assessment of studies of diagnostic accuracy included in systematic reviews. *BMC Med Res Methodol* 2003; **3**: 25 [PMID: 14606960 DOI: 10.1186/1471-2288-3-25]
 - 10 **Ahlquist DA**, Taylor WR, Mahoney DW, Zou H, Domanico M, Thibodeau SN, Boardman LA, Berger BM, Lidgard GP. The stool DNA test is more accurate than the plasma septin 9 test in detecting colorectal neoplasia. *Clin Gastroenterol Hepatol* 2012; **10**: 272-277.e1 [PMID: 22019796 DOI: 10.1016/j.cgh.2011.10.008]
 - 11 **Bosch LJ**, Oort FA, Neerincx M, Khalid-de Bakker CA, Terhaar sive Droste JS, Melotte V, Jonkers DM, Masclee AA, Mongera S, Grootecloes M, Louwagie J, van Criekinge W, Coupé VM, Mulder CJ, van Engeland M, Carvalho B, Meijer GA. DNA methylation of phosphatase and actin regulator 3 detects colorectal cancer in stool and complements FIT. *Cancer Prev Res (Phila)* 2012; **5**: 464-472 [PMID: 22135045 DOI: 10.1158/1940-6207.CAPR-11-0315]
 - 12 **Ahlquist DA**, Zou H, Domanico M, Mahoney DW, Yab TC, Taylor WR, Butz ML, Thibodeau SN, Rabeneck L, Paszat LF, Kinzler KW, Vogelstein B, Bjerregaard NC, Laurberg S, Sørensen HT, Berger BM, Lidgard GP. Next-generation stool DNA test accurately detects colorectal cancer and large adenomas. *Gastroenterology* 2012; **142**: 248-256; quiz e25-e26 [PMID: 22062357 DOI: 10.1053/j.gastro.2011.10.031]
 - 13 **Azuara D**, Rodriguez-Moranta F, de Oca J, Soriano-Izquierdo A, Mora J, Guardiola J, Biondo S, Blanco I, Peinado MA, Moreno V, Esteller M, Capellá G. Novel methylation panel for the early detection of colorectal tumors in stool DNA. *Clin Colorectal Cancer* 2010; **9**: 168-176 [PMID: 20643622 DOI: 10.3816/CCC.2010.n.023]
 - 14 **Tang D**, Liu J, Wang DR, Yu HF, Li YK, Zhang JQ. Diagnostic and prognostic value of the methylation status of secreted frizzled-related protein 2 in colorectal cancer. *Clin Invest Med* 2011; **34**: E88-E95 [PMID: 21463549]
 - 15 **Baek YH**, Chang E, Kim YJ, Kim BK, Sohn JH, Park DI. Stool methylation-specific polymerase chain reaction assay for the detection of colorectal neoplasia in Korean patients. *Dis Colon Rectum* 2009; **52**: 1452-1459; discussion 1452-1459 [PMID: 19617759 DOI: 10.1007/DCR.0b013e3181a79533]
 - 16 **Li M**, Chen WD, Papadopoulos N, Goodman SN, Bjerregaard NC, Laurberg S, Levin B, Juhl H, Arber N, Moinova H, Durkee K, Schmidt K, He Y, Diehl F, Velculescu VE, Zhou S, Diaz LA, Kinzler KW, Markowitz SD, Vogelstein B. Sensitive digital quantification of DNA methylation in clinical samples. *Nat Biotechnol* 2009; **27**: 858-863 [PMID: 19684580 DOI: 10.1038/nbt.1559]
 - 17 **Melotte V**, Lentjes MH, van den Bosch SM, Hellebrekers DM, de Hoon JP, Wouters KA, Daenen KL, Partouns-Hendriks IE, Stessels F, Louwagie J, Smits KM, Weijenberg MP, Sanduleanu S, Khalid-de Bakker CA, Oort FA, Meijer GA, Jonkers DM, Herman JG, de Bruïne AP, van Engeland M. N-Myc downstream-regulated gene 4 (NDRG4): a candidate tumor suppressor gene and potential biomarker for colorectal cancer. *J Natl Cancer Inst* 2009; **101**: 916-927 [PMID: 19535783 DOI: 10.1093/jnci/djp131]
 - 18 **Ausch C**, Kim YH, Tsuchiya KD, Dzieciatkowski S, Washington MK, Paraskeva C, Radich J, Grady WM. Comparative analysis of PCR-based biomarker assay methods for colorectal polyp detection from fecal DNA. *Clin Chem* 2009; **55**: 1559-1563 [PMID: 19541867 DOI: 10.1373/clinchem.2008.122937]
 - 19 **Hellebrekers DM**, Lentjes MH, van den Bosch SM, Melotte V, Wouters KA, Daenen KL, Smits KM, Akiyama Y, Yuasa Y, Sanduleanu S, Khalid-de Bakker CA, Jonkers D, Weijenberg MP, Louwagie J, van Criekinge W, Carvalho B, Meijer GA, Baylin SB, Herman JG, de Bruïne AP, van Engeland M. GATA4 and GATA5 are potential tumor suppressors and biomarkers in colorectal cancer. *Clin Cancer Res* 2009; **15**: 3990-3997 [PMID: 19509152 DOI: 10.1158/1078-0432.CCR-09-0055]
 - 20 **Mayor R**, Casadomé L, Azuara D, Moreno V, Clark SJ, Capellá G, Peinado MA. Long-range epigenetic silencing at 2q14.2 affects most human colorectal cancers and may have application as a non-invasive biomarker of disease. *Br J Cancer* 2009; **100**: 1534-1539 [PMID: 19384295 DOI: 10.1038/sj.bjc.6605045]
 - 21 **Kim MS**, Louwagie J, Carvalho B, Terhaar Sive Droste JS, Park HL, Chae YK, Yamashita K, Liu J, Ostrow KL, Ling S, Guerrero-Preston R, Demokan S, Yalniz Z, Dalay N, Meijer GA, Van Criekinge W, Sidransky D. Promoter DNA methylation of oncostatin m receptor-beta as a novel diagnostic and therapeutic marker in colon cancer. *PLoS One* 2009; **4**: e6555 [PMID: 19662090 DOI: 10.1371/journal.pone.0006555]
 - 22 **Nagasaka T**, Tanaka N, Cullings HM, Sun DS, Sasamoto H, Uchida T, Koi M, Nishida N, Naomoto Y, Boland CR, Matsubara N, Goel A. Analysis of fecal DNA methylation to detect gastrointestinal neoplasia. *J Natl Cancer Inst* 2009; **101**: 1244-1258 [PMID: 19700653 DOI: 10.1093/jnci/djp265]
 - 23 **Glöckner SC**, Dhir M, Yi JM, McGarvey KE, Van Neste L, Louwagie J, Chan TA, Kleeberger W, de Bruïne AP, Smits KM, Khalid-de Bakker CA, Jonkers DM, Stockbrügger RW, Meijer GA, Oort FA, Iacobuzio-Donahue C, Bierau K, Herman JG, Baylin SB, Van Engeland M, Schuebel KE, Ahuja N. Methylation of TFP12 in stool DNA: a potential novel biomarker for the detection of colorectal cancer. *Cancer Res* 2009; **69**: 4691-4699 [PMID: 19435926 DOI: 10.1158/0008-5472.CAN-08-0142]
 - 24 **Wang DR**, Tang D. Hypermethylated SFRP2 gene in fecal DNA is a high potential biomarker for colorectal cancer noninvasive screening. *World J Gastroenterol* 2008; **14**: 524-531 [PMID: 18203283 DOI: 10.3748/wjg.14.524]
 - 25 **Oberwalder M**, Zitt M, Wöntner C, Fiegl H, Goebel G, Zitt M, Köhle O, Mühlmann G, Ofner D, Margreiter R, Müller HM. SFRP2 methylation in fecal DNA--a marker for colorectal polyps. *Int J Colorectal Dis* 2008; **23**: 15-19 [PMID: 17639423 DOI: 10.1007/s00384-007-0355-2]
 - 26 **Itzkowitz S**, Brand R, Jandorf L, Durkee K, Millholland J, Rabeneck L, Schroy PC, Sontag S, Johnson D, Markowitz S, Paszat L, Berger BM. A simplified, noninvasive stool DNA test for colorectal cancer detection. *Am J Gastroenterol* 2008; **103**: 2862-2870 [PMID: 18759824 DOI: 10.1111/j.1572-0241.2008.02088.x]
 - 27 **Huang ZH**, Li LH, Yang F, Wang JF. Detection of aberrant methylation in fecal DNA as a molecular screening tool for colorectal cancer and precancerous lesions. *World J Gastroenterol* 2007; **13**: 950-954 [PMID: 17352030]
 - 28 **Itzkowitz SH**, Jandorf L, Brand R, Rabeneck L, Schroy PC, Sontag S, Johnson D, Skoletsky J, Durkee K, Markowitz S, Shuber A. Improved fecal DNA test for colorectal cancer screening. *Clin Gastroenterol Hepatol* 2007; **5**: 111-117 [PMID: 17161655 DOI: 10.1016/j.cgh.2006.10.006]
 - 29 **Abbaszadegan MR**, Tavasoli A, Velayati A, Sima HR, Vosooghnia H, Farzadnia M, Asadzadeh H, Gholamin M, Dadkhah E, Aarabi A. Stool-based DNA testing, a new noninvasive method for colorectal cancer screening, the first report from Iran. *World J Gastroenterol* 2007; **13**: 1528-1533 [PMID: 17461444]

- 30 **Zhang W**, Bauer M, Croner RS, Pelz JO, Lodygin D, Herme-king H, Stürzl M, Hohenberger W, Matzel KE. DNA stool test for colorectal cancer: hypermethylation of the secreted frizzled-related protein-1 gene. *Dis Colon Rectum* 2007; **50**: 1618-1626; discussion 1618-1626 [PMID: 17762966 DOI: 10.1007/s10350-007-0286-6]
- 31 **Leung WK**, To KF, Man EP, Chan MW, Hui AJ, Ng SS, Lau JY, Sung JJ. Detection of hypermethylated DNA or cyclooxygenase-2 messenger RNA in fecal samples of patients with colorectal cancer or polyps. *Am J Gastroenterol* 2007; **102**: 1070-1076 [PMID: 17378912 DOI: 10.1111/j.1572-0241.2007.01108.x]
- 32 **Petko Z**, Ghiassi M, Shuber A, Gorham J, Smalley W, Washington MK, Schultenover S, Gautam S, Markowitz SD, Grady WM. Aberrantly methylated CDKN2A, MGMT, and MLH1 in colon polyps and in fecal DNA from patients with colorectal polyps. *Clin Cancer Res* 2005; **11**: 1203-1209 [PMID: 15709190]
- 33 **Lenhard K**, Bommer GT, Asutay S, Schauer R, Brabletz T, Göke B, Lamerz R, Kolligs FT. Analysis of promoter methylation in stool: a novel method for the detection of colorectal cancer. *Clin Gastroenterol Hepatol* 2005; **3**: 142-149 [PMID: 15704048 DOI: 10.1016/S1542-3565(04)00624-X]
- 34 **Chen WD**, Han ZJ, Skoletsky J, Olson J, Sah J, Myeroff L, Platzer P, Lu S, Dawson D, Willis J, Pretlow TP, Lutterbaugh J, Kasturi L, Willson JK, Rao JS, Shuber A, Markowitz SD. Detection in fecal DNA of colon cancer-specific methylation of the nonexpressed vimentin gene. *J Natl Cancer Inst* 2005; **97**: 1124-1132 [PMID: 16077070 DOI: 10.1093/jnci/dji204]
- 35 **Müller HM**, Oberwalder M, Fiegl H, Morandell M, Goebel G, Zitt M, Mühlthaler M, Ofner D, Margreiter R, Widschwendter M. Methylation changes in faecal DNA: a marker for colorectal cancer screening? *Lancet* 2004; **363**: 1283-1285 [PMID: 15094274 DOI: 10.1016/S0140-6736(04)16002-9]
- 36 **Xu MH**, Cai KY, Tu Y. Stool DNA methylation analysis in the early diagnosis of colorectal cancer. *Chin J Clin Gastroenterol* 2012; **24**: 7-19 [DOI: 10.3870/lcxh.j.issn.1005-541X.2012.01.06]
- 37 **Kang YP**, Cao FA, Chang WJ, Lou Z, Wang Y, Wu LL, Fu CG, Cao GW. Stool DNA methylation in the screening early colorectal cancer. *J Gastrointest Surg* 2011; **14**: 52-561 [DOI: 10.3760/cma.j.issn.1671-0274.2011.01.017]
- 38 **Zhang JP**, Wang J, Gui YL, Zhu QQ, Xu ZW, Li JS. Stool DNA methylation in the diagnosis of colorectal cancer. *Chin Med J* 2011; **95**: 2482-2484 [DOI: 10.3760/cma.j.issn.0376-2491.2011.35.010]
- 39 **Fu L**, Sheng JQ, Meng XM, Meng MM, Jin P, Li AQ, Wu ZT, Li SR. Stool Vimentin methylation in the diagnosis of colorectal cancer. *J Gastroenterol Liver Dis* 2010; **19**: 601-603 [DOI: 10.3969/j.issn.1006-5709.2010.07.007]
- 40 **Ling ZA**, Chen LS, He CG. Stool P16 methylation in the diagnosis of colorectal cancer. *Colorectal Anus Surg* 2009; **15**: 144-148 [DOI: 10.3969/j.issn.1674-0491.2009.03.002]
- 41 **Cheng ZH**. Stool SFRP2 methylation in the diagnosis of colorectal cancer. *Shandong Yiyao Zazhi* 2007; **47**: 10-12 [DOI: 10.3969/j.issn.1002-266X.2007.06.004]
- 42 **Zhao HX**, Li QW, Dong WW, Duan XH, Zhu JH, Wang RL, Hao YX, Ye M, Xiao WH. Stool NDRG4 methylation in the diagnosis of colorectal cancer. *Zhongguo Yiyao Daobao* 2012; **9**: 29-31 [DOI: 10.3877/cma.j.issn.1674-0785.2012.10.012]
- 43 **Chang E**, Park DI, Kim YJ, Kim BK, Park JH, Kim HJ, Cho YK, Sohn CL, Jeon WK, Kim BI, Kim HD, Kim DH, Kim YH. Detection of colorectal neoplasm using promoter methylation of ITGA4, SFRP2, and p16 in stool samples: a preliminary report in Korean patients. *Hepatogastroenterology* 2010; **57**: 720-727 [PMID: 21033217]
- 44 **Kim MS**, Lee J, Sidransky D. DNA methylation markers in colorectal cancer. *Cancer Metastasis Rev* 2010; **29**: 181-206 [PMID: 20135198 DOI: 10.1007/s10555-010-9207-6]
- 45 **Imperiale TF**, Ransohoff DF, Itzkowitz SH, Turnbull BA, Ross ME. Fecal DNA versus fecal occult blood for colorectal cancer screening in an average-risk population. *N Engl J Med* 2004; **351**: 2704-2714 [PMID: 15616205 DOI: 10.1056/NEJ-Moa033403]
- 46 **Qi J**, Zhu YQ, Luo J, Tao WH. Hypermethylation and expression regulation of secreted frizzled-related protein genes in colorectal tumor. *World J Gastroenterol* 2006; **12**: 7113-7117 [PMID: 17131472]

P- Reviewers: Meshikhes AWN, Xia HHX

S- Editor: Cui XM **L- Editor:** Wang TQ **E- Editor:** Lu YJ





百世登

Baishideng®

Published by **Baishideng Publishing Group Co., Limited**

Flat C, 23/F., Lucky Plaza,

315-321 Lockhart Road, Wan Chai, Hong Kong, China

Fax: +852-65557188

Telephone: +852-31779906

E-mail: bpgoffice@wjgnet.com

<http://www.wjgnet.com>

