

# World Journal of *Clinical Cases*

*World J Clin Cases* 2023 July 26; 11(21): 4966-5192



**OPINION REVIEW**

- 4966 Flexner has fallen: Transitions in medical education system across time, a gradual return to pre-Flexnerian state (de-Flexnerization)  
*Alzerwi NAN*

**REVIEW**

- 4975 Manipulation of the intestinal microbiome-a slow journey to primetime  
*Kruger-Sharabi O, Malnick SDH, Fisher D*

**MINIREVIEWS**

- 4989 Menstrual cycle abnormalities in women with inflammatory bowel disease and effects of biological therapy on gynecological pathology  
*Malinauskiene V, Zuzo A, Liakina V, Kazenaite E, Stundiene I*
- 4996 Exhaled volatile organic compounds for diagnosis and monitoring of asthma  
*Savito L, Scarlata S, Bikov A, Carratù P, Carpagnano GE, Dragonieri S*

**ORIGINAL ARTICLE****Retrospective Cohort Study**

- 5014 Cohort analysis of pediatric intussusception score to diagnose intussusception  
*Rukwong P, Wangviwat N, Phewplung T, Sintusek P*

**Retrospective Study**

- 5023 Identification of functional genes regulating gastric cancer progression using integrated bioinformatics analysis  
*Yu K, Zhang D, Yao Q, Pan X, Wang G, Qian HY, Xiao Y, Chen Q, Mei K*
- 5035 Clinical outcomes and complications of single anastomosis duodenal-ileal bypass with sleeve gastrectomy: A 2-year follow-up study in Bogotá, Colombia  
*Ospina Jaramillo A, Riscanevo Bobadilla AC, Espinosa MO, Valencia A, Jiménez H, Montilla Velásquez MDP, Bastidas M*
- 5047 Image-based visualization of stents in mechanical thrombectomy for acute ischemic stroke: Preliminary findings from a series of cases  
*Yao QY, Fu ML, Zhao Q, Zheng XM, Tang K, Cao LM*
- 5056 Clinical effects of a traditional Chinese medicine nursing programme to intervene in gastric pain of the spleen and stomach with Qi deficiency  
*Yu YH, Yu YS, Cheng XH*

- 5063** Behavior analysis and formative assessments in online oral medicine education during the COVID-19 pandemic

*Ye JJ, Zhao YK, Teng ZS, Ye HW, Yuan Q, Nie X*

- 5073** Clinical study of acupuncture combined with surface anesthesia using proparacaine in geriatric cataract phacoemulsification

*Dong P, Wang LF, Zhang LX, Li F, Yin HS, Dou ZX, Huang XJ, Xu R, Zhang WL*

### Clinical Trials Study

- 5083** Safety evaluation of human umbilical cord-mesenchymal stem cells in type 2 diabetes mellitus treatment: A phase 2 clinical trial

*Lian XF, Lu DH, Liu HL, Liu YJ, Yang Y, Lin Y, Xie F, Huang CH, Wu HM, Long AM, Hui CJ, Shi Y, Chen Y, Gao YF, Zhang F*

### Observational Study

- 5097** Time usage analysis and satisfaction comparison by occupational area according to the sex of single-parent families

*Jang WH, Jang JS, Bang JH*

- 5108** Evaluation of bronchoscopic direct vision glottis anesthesia method in bronchoscopy

*Lang J, Guo ZZ, Xing SS, Sun J, Qiu B, Shu Y, Wang ZQ, Liu GX*

### CASE REPORT

- 5115** Type I Mirizzi syndrome treated by electrohydraulic lithotripsy under the direct view of SpyGlass: A case report

*Liang SN, Jia GF, Wu LY, Wang JZ, Fang Z, Wang SH*

- 5122** Primary ovarian angiosarcoma: Two case reports and review of literature

*Zhou Y, Sun YW, Liu XY, Shen DH*

- 5129** Acute myelomonocytic leukemia and T-lymphoblastic lymphoma as simultaneous bilineage hematologic malignancy treated with decitabine: A case report

*Jeon SY, Lee NR, Cha S, Yhim HY, Kwak JY, Jang KY, Kim N, Cho YG, Lee CH*

- 5136** Systematic review of fibroblastic rheumatism: A case report

*Guo H, Liang Q, Dong C, Zhang Q, Gu ZF*

- 5147** Improved super-elastic Ti-Ni alloy wire for treating adult skeletal class III with facial asymmetry: A case report

*Huang CY, Chen YH, Lin CC, Yu JH*

- 5160** Neuroendocrine carcinoma of the endometrium concomitant with Lynch syndrome: A case report

*Siu WYS, Hong MK, Ding DC*

- 5167** Myeloperoxidase-antineutrophil cytoplasmic antibody-associated vasculitis with headache and kidney involvement at presentation and with arthralgia at relapse: A case report

*Zhang X, Zhao GB, Li LK, Wang WD, Lin HL, Yang N*

- 5173** Rare combination of traumatic subarachnoid-pleural fistula and intracranial subdural hygromas: A case report  
*Chen PH, Li CR, Gan CW, Yang TH, Chang CS, Chan FH*
- 5179** *ADAMTS3* and *FLT4* gene mutations result in congenital lymphangiectasia in newborns: A case report  
*Liang ZW, Gao WL*
- 5187** Brucellosis presenting with pancytopenia and hearing loss: A case report  
*Chen X, Yang FB, Liang JY*

**ABOUT COVER**

Editorial Board Member of *World Journal of Clinical Cases*, Baharudin Ibrahim, BPharm, PhD, Associate Professor, Pharmacist, Department of Clinical Pharmacy and Pharmacy Practice, Faculty of Pharmacy, Universiti Malaya, Kuala Lumpur 50603, Malaysia. baharudin.ibrahim@um.edu.my

**AIMS AND SCOPE**

The primary aim of *World Journal of Clinical Cases* (*WJCC*, *World J Clin Cases*) is to provide scholars and readers from various fields of clinical medicine with a platform to publish high-quality clinical research articles and communicate their research findings online.

*WJCC* mainly publishes articles reporting research results and findings obtained in the field of clinical medicine and covering a wide range of topics, including case control studies, retrospective cohort studies, retrospective studies, clinical trials studies, observational studies, prospective studies, randomized controlled trials, randomized clinical trials, systematic reviews, meta-analysis, and case reports.

**INDEXING/ABSTRACTING**

The *WJCC* is now abstracted and indexed in Science Citation Index Expanded (SCIE, also known as SciSearch®), Journal Citation Reports/Science Edition, Current Contents®/Clinical Medicine, PubMed, PubMed Central, Reference Citation Analysis, China National Knowledge Infrastructure, China Science and Technology Journal Database, and Superstar Journals Database. The 2023 Edition of Journal Citation Reports® cites the 2022 impact factor (IF) for *WJCC* as 1.1; IF without journal self cites: 1.1; 5-year IF: 1.3; Journal Citation Indicator: 0.26; Ranking: 133 among 167 journals in medicine, general and internal; and Quartile category: Q4.

**RESPONSIBLE EDITORS FOR THIS ISSUE**

Production Editor: *Ying-Yi Yuan*; Production Department Director: *Xiang Li*; Editorial Office Director: *Jin-Lei Wang*.

**NAME OF JOURNAL**

*World Journal of Clinical Cases*

**ISSN**

ISSN 2307-8960 (online)

**LAUNCH DATE**

April 16, 2013

**FREQUENCY**

Thrice Monthly

**EDITORS-IN-CHIEF**

Bao-Gan Peng, Jerzy Tadeusz Chudek, George Kontogeorgos, Maurizio Serati, Ja Hyeon Ku

**EDITORIAL BOARD MEMBERS**

<https://www.wjgnet.com/2307-8960/editorialboard.htm>

**PUBLICATION DATE**

July 26, 2023

**COPYRIGHT**

© 2023 Baishideng Publishing Group Inc

**INSTRUCTIONS TO AUTHORS**

<https://www.wjgnet.com/bpg/gerinfo/204>

**GUIDELINES FOR ETHICS DOCUMENTS**

<https://www.wjgnet.com/bpg/GerInfo/287>

**GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH**

<https://www.wjgnet.com/bpg/gerinfo/240>

**PUBLICATION ETHICS**

<https://www.wjgnet.com/bpg/GerInfo/288>

**PUBLICATION MISCONDUCT**

<https://www.wjgnet.com/bpg/gerinfo/208>

**ARTICLE PROCESSING CHARGE**

<https://www.wjgnet.com/bpg/gerinfo/242>

**STEPS FOR SUBMITTING MANUSCRIPTS**

<https://www.wjgnet.com/bpg/GerInfo/239>

**ONLINE SUBMISSION**

<https://www.f6publishing.com>

## Observational Study

# Evaluation of bronchoscopic direct vision glottis anesthesia method in bronchoscopy

Jie Lang, Zhi-Zhen Guo, Shu-Shan Xing, Jian Sun, Bin Qiu, Yu Shu, Zhi-Qiang Wang, Gui-Xiang Liu

**Specialty type:** Medicine, general and internal**Provenance and peer review:** Unsolicited article; Externally peer reviewed.**Peer-review model:** Single blind**Peer-review report's scientific quality classification**Grade A (Excellent): 0  
Grade B (Very good): 0  
Grade C (Good): C  
Grade D (Fair): D  
Grade E (Poor): 0**P-Reviewer:** Taheri S, Iran;  
Tsuchiya M, Japan**Received:** April 27, 2023**Peer-review started:** April 27, 2023**First decision:** June 1, 2023**Revised:** June 6, 2023**Accepted:** July 3, 2023**Article in press:** July 3, 2023**Published online:** July 26, 2023**Jie Lang, Jian Sun, Bin Qiu, Yu Shu, Zhi-Qiang Wang, Gui-Xiang Liu**, Department of Thoracic Surgery, Tangshan People's Hospital, Tangshan 063000, Hebei Province, China**Zhi-Zhen Guo**, Department of Nephrology, Kailuan General Hospital, Tangshan 063000, Hebei Province, China**Shu-Shan Xing**, Department of Oncology, Tangshan Central Hospital, Tangshan 063000, Hebei Province, China**Corresponding author:** Shu-Shan Xing, MD, Deputy Director, Department of Oncology, Tangshan Central Hospital, No. 601 Changning Road, Tangshan 063000, Hebei Province, China. [xingshushan2022@163.com](mailto:xingshushan2022@163.com)

## Abstract

### BACKGROUND

Fibrobronchoscopy is a common adjunct tool that requires anesthesia and is widely used in the diagnosis and treatment of various respiratory diseases. However, current anesthesia methods, such as spray, nebulized inhalation, and cricothyroid membrane puncture, have their own advantages and disadvantages. Recently, studies have shown that bronchoscopic direct-view glottis anesthesia is a simple and inexpensive method that shortens the examination time and provides excellent anesthetic results.

### AIM

To evaluate the effectiveness of bronchoscopic direct vision glottis anesthesia for bronchoscopy.

### METHODS

The study included 100 patients who underwent bronchoscopy during thoracic surgery. A random number table method was used to divide the patients into control and observation groups (50 patients each). The control and observation groups were anesthetized using the nebulized inhalation and bronchoscopic direct vision glottis method, respectively. Hemodynamic indices [systolic blood pressure (SBP), diastolic blood pressure (DBP), heart rate (HR), and oxygen saturation (SpO<sub>2</sub>) before (T1), 5 min after anesthesia (T2), and at the end of the operation (T3)] serum stress hormone indices [norepinephrine (NE), epinephrine (E), adrenocorticotropic hormone (ACTH), and cortisol (Cor) before and after treatment] were compared between the 2 groups. Adverse effects were also

compared between the two groups.

## RESULTS

At T2 and T3, SBP, DBP, and HR were lower in the observation group than the control group, whereas SpO<sub>2</sub> was higher than the control group [(119.05 ± 8.01) mmHg *vs* (127.05 ± 7.83) mmHg, (119.35 ± 6.66) mmHg *vs* (128.39 ± 6.56) mmHg, (84.68 ± 6.04) mmHg *vs* (92.42 ± 5.57) mmHg, (84.53 ± 4.97) mmHg compared to (92.57 ± 6.02) mmHg, (74.25 ± 5.18) beats/min compared to (88.32 ± 5.72) beats/min, (74.38 ± 5.31) beats/min compared to (88.42 ± 5.69) beats/min, (97.36 ± 2.21)% *vs* (94.35 ± 2.16)%, (97.42 ± 2.36)% *vs* (94.38 ± 2.69%)], with statistically significant differences (all *P* < 0.05). After treatment, NE, E, ACTH, and Cor were significantly higher in both groups than before treatment, but were lower in the observation group than in the control group [(68.25 ± 8.87) ng/mL *vs* (93.35 ± 14.00) ng/mL, (53.59 ± 5.89) ng/mL *vs* (82.32 ± 10.70) ng/mL, (14.32 ± 1.58) pg/mL *vs* (20.35 ± 3.05) pg/mL, (227.35 ± 25.01) nmol/L *vs* (322.28 ± 45.12) nmol/L], with statistically significant differences (all *P* < 0.05). The incidence of adverse reactions was higher in the control group than in the observation group [12.00% (12/50) *vs* 6.00% (3/50)] (*P* < 0.05).

## CONCLUSION

The use of bronchoscopic direct vision glottis anesthesia method for bronchoscopy patients is beneficial for stabilizing hemodynamic indices during bronchoscopy and reducing the level of patient stress, with good safety and practicality.

**Key Words:** Direct bronchoscopy; Glottis anesthesia method; Bronchoscopy; Airway; Sedation

©The Author(s) 2023. Published by Baishideng Publishing Group Inc. All rights reserved.

**Core Tip:** We evaluated the use of bronchoscopic direct vision glottis anesthesia method for bronchoscopy patients, and found it was beneficial for stabilizing hemodynamic indices during bronchoscopy and reducing the patient stress level with good safety and practicality.

**Citation:** Lang J, Guo ZZ, Xing SS, Sun J, Qiu B, Shu Y, Wang ZQ, Liu GX. Evaluation of bronchoscopic direct vision glottis anesthesia method in bronchoscopy. *World J Clin Cases* 2023; 11(21): 5108-5114

**URL:** <https://www.wjgnet.com/2307-8960/full/v11/i21/5108.htm>

**DOI:** <https://dx.doi.org/10.12998/wjcc.v11.i21.5108>

## INTRODUCTION

Fibrobronchoscopy is a common adjunct tool in the clinical setting, and with technological advancements it has been used in the diagnosis and treatment of various respiratory diseases[1,2]. This examination method is invasive and requires the placement of a bronchoscope into the lower respiratory tract through the oral or nasal cavity. It enters the airway and bronchi through the larynx or even more distally to examine the tracheal and bronchial lesions and provide appropriate treatment according to the condition of the lesions. During bronchial manipulation, patients experience significant discomfort; therefore, anesthesia is necessary[2]. Previously, nebulized inhalation anesthesia was used, which, to some extent, relaxed and opened the patient's vocal cords, reduced airway resistance, and prevented bronchospasm; however, the overall anesthetic effect was poor[3]. Recent studies have shown that bronchoscopic direct vision vocal anesthesia-assisted bronchoscopy has better clinical performance[4]. Therefore, this study examined the effect of bronchoscopic direct vision vocal anesthesia on bronchoscopy in patients who underwent bronchoscopy.

## MATERIALS AND METHODS

### Subjects and methods

In total, 100 patients who underwent bronchoscopy at the Department of Thoracic Surgery of Tangshan People's Hospital between June 2017 and June 2019 were included in the study. The random number table method was used to divide the patients into control and observation groups (50 patients each). The inclusion criteria were: (1) Patients with respiratory system diseases that require bronchoscopy; (2) are older than 18 years and younger than 65 years, with no restriction on gender; (3) patients and/or family members voluntarily participate in this study, ensure compliance with all measures, and sign an informed consent form; and (4) are receiving bronchoscopy for the first time, have American Society of Anesthesiologists classification of grade I or II, and have been prescribed the anesthetic drugs used in this study. The exclusion criteria were: (1) Allergy to the drugs used in this study; (2) congenital malformations of the respiratory system; (3) mental disorders or cognitive abnormalities that prevent normal examination procedures; (4) women who are breast-

feeding or pregnant; and (5) long-term alcohol or drug use resulting in increased tolerance of the body to anesthetic drugs. The control group comprised of 32 males and 18 females, aged 42-65 years, with an average age of  $(52.25 \pm 8.13)$  years, an average body mass index (BMI) of  $(23.18 \pm 3.50)$  kg/m<sup>2</sup>, with six cases of pneumonia, 22 cases of lung cancer, 8 cases of chronic obstructive pulmonary disease, eight cases of bronchiectasis, and six cases of tuberculosis as the primary disease. The observation group comprised of 34 males and 16 females aged 41-64 years, with a mean age of  $(51.70 \pm 8.05)$  years, a mean BMI of  $(23.22 \pm 2.87)$  kg/m<sup>2</sup>, seven cases of pneumonia, 23 cases of lung cancer, five cases of chronic obstructive pulmonary disease, 10 cases of bronchiectasis, and five cases of tuberculosis as the primary cause. The study was performed after obtaining approval from the Medical Ethics Committee of Tangshan People's Hospital.

### Examination methods

Patients in the two groups routinely fasted for 8 h prior to the examination. Pulsed high-flow oxygen was used during the examination, and the patient vital signs were monitored. After completing the examination, all patients received supportive symptomatic treatment. An electronic bronchoscope (Olympus, BF-1T260, Japan) was used for the examination. The control group was anesthetized by the nebulized inhalation method using 15 mL of 2% lidocaine hydrochloride injection (Henan Province Shennong Pharmaceutical Co., Ltd., State Drug Quantifier H20043428, 5 mL:0.1 g, Lot No. 2017052258) for 10 min. Bronchoscopy was then performed, with the patients receiving oxygen during the operation. In the observation group, bronchoscopy was performed by direct vision bronchoscopy, and 1% lidocaine solution was injected directly into the vocal cords when the lens passed through the epiglottis for 2-3 mL. The patient was asked to breathe deeply to allow the anesthetic to enter the trachea through the vocal cords. The patient was allowed to breathe deeply for 1 minute after removing the bronchoscope. Total amount of lidocaine was controlled under 300mg per patient. The procedure was repeated and venous blood was collected from the patient before and after treatment for testing.

### Observation indices

Hemodynamic indices: Systolic blood pressure (SBP), diastolic blood pressure (DBP), heart rate (HR), and oxygen saturation (SpO<sub>2</sub>) before anesthesia (T1), 5 min after anesthesia (T2), and after the end of the operation (T3) in the two groups were measured. Serum stress hormone indicators: norepinephrine (NE), epinephrine (E), adrenocorticotrophic hormone (ACTH), and cortisol (Cor) were also measured. Serum was obtained after 5 mL of venous blood was drawn at 3000 r/min with a radius of 10 cm at high speed for 15 min, placed in a low temperature refrigerator at -20 °C and tested within 2 h by enzyme-linked immunosorbent assay (Wuhan Ruisheng Biotechnology Co., Ltd., China) using an automatic biochemistry instrument (Beckman Coulter, AU7900) following the kit and instrument's instructions. Adverse reaction indices: Adverse reactions that occurred during anesthesia in the two groups were recorded including limb twitching, abnormal breathing, choking, and poor memory.

### Statistical analysis

SPSS24.0 software was used for statistical analyses; the results of measurement data are expressed as (mean  $\pm$  SD), and the results of count data are expressed as number of cases (*n*) and percentage (%). Independent samples *t*-test was used to compare measurement data between groups, paired *t*-test was used to compare measurement data before and after treatment within groups, and  $\chi^2$  test was used to compare count data between groups. Statistical significance was set at *P* < 0.05.

## RESULTS

### Comparison of hemodynamic indexes between the two groups

At T1, the differences between SBP, DBP, HR, and SpO<sub>2</sub> in the two groups were not statistically significant (all *P* > 0.05); however, at T2 and T3, SBP, DBP, and HR were significantly higher and SpO<sub>2</sub> was significantly lower in the control group, whereas, none of the hemodynamic indices showed significant changes in the observation group. In T2 and T3, SBP, DBP, and HR in the observation group were lower than the control group, while SpO<sub>2</sub> was higher than the control group, and the differences were statistically significant (*P* < 0.05) (Table 1). Plasma concentrations were investigated and no significance was found between the case and control groups (case:  $2.25 \pm 0.12$ ; control:  $2.21 \pm 0.14$ ; *P* > 0.05).

### Comparison of serum stress hormone indexes between the two groups

Prior to treatment, there was no statistically significant difference between the two groups in NE, E, ACTH, and Cor levels (*P* > 0.05). Post treatment, the NE, E, ACTH, and Cor levels of the two groups were significantly higher, however, NE, E, ACTH, and Cor in the observation group were significantly lower than the control group (*P* < 0.05) (Table 2).

### Comparison of adverse reaction indicators between the two groups

The incidence of adverse reactions was higher in the control group than in the observation group (*P* < 0.05) (Table 3).

Table 1 Comparison of hemodynamic indices between two groups of patients undergoing bronchoscopy

Group	No.	SBP (mmHg)			DBP (mmHg)			HR (/min)			SpO <sub>2</sub> (%)		
		T1	T2	T3	T1	T2	T3	T1	T2	T3	T1	T2	T3
Case	50	118.02 ± 8.69	119.05 ± 8.01	119.35 ± 6.66	84.57 ± 5.38	84.68 ± 6.04	84.53 ± 4.97	73.58 ± 7.36	74.25 ± 5.18	74.38 ± 5.31	97.35 ± 2.35	97.36 ± 2.21	97.42 ± 2.36
Control	50	119.25 ± 7.95	127.05 ± 7.83 <sup>a</sup>	128.39 ± 6.56 <sup>a</sup>	84.35 ± 6.35	92.42 ± 5.57 <sup>a</sup>	92.57 ± 6.02 <sup>a</sup>	74.05 ± 6.57	88.32 ± 5.72 <sup>a</sup>	88.42 ± 5.69 <sup>a</sup>	97.41 ± 2.27	94.35 ± 2.16 <sup>a</sup>	94.38 ± 2.69 <sup>a</sup>
<i>t</i> value	-	-0.738	-5.05	-6.838	0.187	-6.661	-7.283	-0.337	-12.892	-12.756	-0.13	6.887	6.007
<i>P</i> value	-	0.462	< 0.001	< 0.001	0.852	< 0.001	< 0.001	0.737	< 0.001	< 0.001	0.897	< 0.001	< 0.001

<sup>a</sup>*P* < 0.05.

The control group received lidocaine nebulized inhalation anesthesia; the observation group received bronchoscopic direct vision glottis anesthesia. SBP: Systolic blood pressure; DBP: Diastolic blood pressure; HR: Heart rate; SPO<sub>2</sub>: Oxygen saturation.

## DISCUSSION

In the process of fiberoptic bronchoscopy, the bronchoscope will inevitably stimulate or pull on the pharynx and tracheal wall during surgery, and patients are prone to physiological reflexes such as wheezing, choking, and nausea, which not only seriously affect the operation of bronchoscopy but also cause anxiety, fear, and nervousness in patients[5]. Moreover, a significant proportion of patients are unable to cooperate with the completion of monitoring, making the examination meaningless[6].

In recent years, local anesthesia methods before bronchoscopy have included spray, nebulized inhalation, cricothyroid membrane puncture, tracheal drip, and nasal drip anesthesia methods; the first three are the most common[7,8]. Various local anesthesia methods have their own advantages and disadvantages; for example, for the cricothyroid puncture injection, endotracheal anesthesia effect is good. There is no anesthetic effect on the pear-shaped fossa, but it is an invasive operation, there is a risk of bleeding while increasing the patient's pain, and the patient compliance requirements are high[8]. The nebulized inhalation method is simple and easy, with high compliance; however, its anesthetic effect is impacted by many factors, such as awake patients, number of secretions in the airway, and individual differences, which have a greater impact on the amount of anesthetic required[9]. The use of intravenous anesthesia poses certain risks[10] requires the involvement of professional anesthesiologists, and the corresponding resuscitation measures need to be in a ready state. The bronchoscopic direct-view glottis anesthesia method has a good anesthetic effect on both the endotracheal and pear-shaped fossa[11], which can improve the patient's tolerance to bronchoscopy and reduce the degree of postoperative pain recall. Bronchoscopic direct-view glottis anesthesia is simple, inexpensive, shortens examination time, and provides excellent anesthetic results. This greatly reduces or directly eliminates all types of physiological and psychological abnormalities caused by the stimulation and pulling of tissues during the endoscopy and improves the patient's ability to tolerate and complete the examination.

The observation group in this study received bronchoscopic direct vision glottis anesthesia. Compared to the nebulized anesthesia method, the direct-view subglottic drip anesthesia method can be performed under direct vision and is easy to operate and promote. Clinical research has shown that the application of the direct-view subsonic drip method of anesthesia has a rapid onset of action, an ideal anesthetic effect, and reduces the time required for bronchoscopy. Compared with other anesthesia methods, this method requires fewer instruments and can reduce the economic burden

**Table 2 Comparison of serum stress hormone indicators between two groups of bronchoscopy patients**

Group	Case	NE (ng/mL)		E (ng/mL)		ACTH (pg/mL)		Cor (nmol/L)	
		Before	After	Before	After	Before	After	Before	After
Case	50	60.21 ± 8.43	68.25 ± 8.87 <sup>a</sup>	48.02 ± 6.72	53.59 ± 5.89 <sup>a</sup>	12.02 ± 1.80	14.32 ± 1.58 <sup>a</sup>	190.67 ± 24.79	227.35 ± 25.01 <sup>a</sup>
Control	50	59.62 ± 8.35	93.35 ± 14.00 <sup>a</sup>	47.55 ± 6.18	82.32 ± 10.70 <sup>a</sup>	11.82 ± 1.54	20.35 ± 3.05 <sup>a</sup>	189.25 ± 24.6	322.28 ± 45.12 <sup>a</sup>
<i>t</i> value	-	-0.352	10.709	-0.364	16.633	-0.597	12.413	-0.288	13.012
<i>P</i> value	-	0.726	< 0.001	0.717	< 0.001	0.552	< 0.001	0.774	< 0.001

<sup>a</sup>*P* < 0.05.

The control group received lidocaine nebulized inhalation anesthesia and the observation group received bronchoscopic direct vision glottis anesthesia. The values were compared before and after treatment. NE: Norepinephrine; E: Epinephrine; ACTH: Adrenocorticotropic hormone; Cor: Cortisol.

**Table 3 Comparison of adverse reaction indicators between two groups of patients undergoing bronchoscopy**

Group	No.	Body stirring	Abnormal breathing	Choking and coughing	Adverse memory	Total
Control	50	3 (5.26)	5 (10.00)	2 (4.00)	2 (4.00)	12 (24.00)
Case	50	2 (4.00)	1 (2.00)	0 (0.00)	0 (0.00)	3 (6.00)
$\chi^2$						6.353
<i>P</i> value						0.012

The control group received lidocaine-nebulized inhalation anesthesia, and the observation group received bronchoscopic direct vision vocal anesthesia.

on patients; therefore, it can be promoted in primary hospitals.

The study compared anesthesia between the two groups in three dimensions: hemodynamics, stress indicators, and adverse reactions. In the control group, only nebulized inhalation anesthesia was used, and the patients inevitably experienced choking and dyspnea during the operation, which led to dramatic changes in HR, SBP, DBP and SpO<sub>2</sub> indices at various time points. Patients in the observation group showed relatively stable hemodynamic indices owing to the precise application of anesthetic drugs; low anesthetic dosage, good local anesthesia effect, minimal response to the operation, and minimal response to tracheal stimulation, thus reflecting a smooth and stable operation process. The levels of serum stress indicators in the two groups were compared before and after treatment. The human adrenal glands are important organs in the stress response. Sympathetic activation promotes the synthesis and secretion of adrenaline and noradrenaline from the adrenal medulla, while activation of the hypothalamic-pituitary-adrenocortical axis promotes the secretion of cortisol from the adrenal cortex *via* ACTH secretion from the pituitary gland. During bronchoscopy, physiological and psychological symptoms caused by stimulation of the tracheal wall and pulling can directly activate the stress response, leading to the release of adrenomedullary hormones and corticosteroids into the circulation[12]. From these results, it is clear that the anesthesia protocol in the observation group effectively alleviated the physiological and psychological stress symptoms of patients during bronchoscopy and inhibited the process of activation of synergistic hormone release by the stress response.

In the comparison of adverse reactions, the observation group also had a clear advantage, as the degree of anesthesia with the bronchoscopic direct-view subglottic anesthesia method was significantly higher than that with the nebulized inhalation anesthesia method, and the patients had a lower risk of various adverse reactions. It should be noted that the use of the bronchoscopic direct visual glottis anesthesia method causes heavy choking symptoms in the early stage of anesthesia, so it is necessary to monitor the patient's breathing, blood pressure, HR, *etc.* at all times to ensure the safety of the patient.

## CONCLUSION

The use of bronchoscopic direct-view subglottic anesthesia for bronchoscopy patients with adequate anesthesia of the pear-shaped crypt and tracheal mucosa is conducive to stabilizing hemodynamic indices during the operation and reducing the level of patient stress, with good safety and practicality.

## ARTICLE HIGHLIGHTS

### Research background

Current anesthesia methods, such as spray, nebulized inhalation, and cricothyroid membrane puncture, exhibit diversity advantages and disadvantages.

### Research motivation

Evaluate the bronchoscopic direct vision glottis anesthesia in bronchoscopy.

### Research objectives

Investigate the effectiveness and side-effect of bronchoscopic direct vision glottis anesthesia in bronchoscopy.

### Research methods

Patients divided into case and control groups, relevant observation indicators and side-effect were investigated.

### Research results

The hemodynamic indices were more stable with lower stress levels in patients during the new anesthesia method.

### Research conclusions

Bronchoscopic direct vision glottis anesthesia exhibited good results.

### Research perspectives

Bronchoscopic direct vision glottis anesthesia method exhibits good clinical applicability and may be widely applicable in clinical practice in the future.

## FOOTNOTES

**Author contributions:** Lang J, Xing SS concept and designed the study; Guo ZZ, Sun J, and Qiu B were be responsible for materials and patients; Shu Y, Wang ZQ, and Liu GX collected and analyzed the data; all authors wrote and approved the final manuscript.

**Supported by** Hebei Science and Technology Program, No. 162777232 (to Xing SS).

**Institutional review board statement:** This study was approved by the ethics committee of Tangshan People's Hospital.

**Informed consent statement:** All study participants, or their legal guardian, provided informed written consent prior to study enrollment.

**Conflict-of-interest statement:** The authors declared no conflicts of interest.

**Data sharing statement:** All datasets are available from the corresponding author.

**STROBE statement:** The authors have read the STROBE Statement – checklist of items, and the manuscript was prepared and revised according to the STROBE Statement – checklist of items.

**Open-Access:** This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

**Country/Territory of origin:** China

**ORCID number:** Shu-Shan Xing [0009-0009-7711-0674](https://orcid.org/0009-0009-7711-0674).

**S-Editor:** Yan JP

**L-Editor:** A

**P-Editor:** Zhang XD

## REFERENCES

- 1 Wu B, Zhu W, Wang Q, Ren C, Wang L, Xie G. Efficacy and safety of ciprofol-remifentanil versus propofol-remifentanil during fiberoptic bronchoscopy: A prospective, randomized, double-blind, non-inferiority trial. *Front Pharmacol* 2022; **13**: 1091579 [PMID: [36618929](https://pubmed.ncbi.nlm.nih.gov/36618929/) DOI: [10.3389/fphar.2022.1091579](https://doi.org/10.3389/fphar.2022.1091579)]

- 2 **Wang L**, Wu Q, Wang M, Ming W, Sheng C, Zhang Y, Chen Y, Cao Y. The safety and efficacy of alfentanil combined with midazolam in fiberoptic bronchoscopy sedation: A randomized, double-blind, controlled trial. *Front Pharmacol* 2022; **13**: 1036840 [PMID: 36339547 DOI: 10.3389/fphar.2022.1036840]
- 3 **Lin AT**, Moore-Clingenpeel M, Karsies TJ. Comparison of two continuous nebulized albuterol doses in critically ill children with status asthmaticus. *J Asthma* 2020; **57**: 980-986 [PMID: 31119958 DOI: 10.1080/02770903.2019.1623249]
- 4 **Wang H**, Wei N, Tang Y, Wang Y, Luo G, Ren T, Xiong C, Li H, Wang M, Qian X. The Utility of Rapid On-Site Evaluation during Bronchoscopic Biopsy: A 2-Year Respiratory Endoscopy Central Experience. *Biomed Res Int* 2019; **2019**: 5049248 [PMID: 31886221 DOI: 10.1155/2019/5049248]
- 5 **Joseph L**, Goldberg S, Cohen S, Picard E. Ibuprofen does not prevent postbronchoscopy fever in children undergoing broncho-alveolar lavage. *Pediatr Pulmonol* 2020; **55**: 2737-2741 [PMID: 32725954 DOI: 10.1002/ppul.24992]
- 6 **Caron M**, Parrot A, Elabbadi A, Dupeyrat S, Turpin M, Baurty T, Rozencajg S, Blayau C, Fulgencio JP, Gibelin A, Blanchard PY, Rodriguez S, Daigné D, Allain MC, Fartoukh M, Pham T. Pain and dyspnea control during awake fiberoptic bronchoscopy in critically ill patients: safety and efficacy of remifentanil target-controlled infusion. *Ann Intensive Care* 2021; **11**: 48 [PMID: 33725225 DOI: 10.1186/s13613-021-00832-6]
- 7 **Lachkar S**, Gervereau D, Lanquetuit M, Thiberville L, Pradier HM, Roger M, Bota S, Guisier F, Salaün M. Hypnosis associated with 3D immersive virtual reality technology during bronchoscopy under local anesthesia. *J Thorac Dis* 2022; **14**: 3205-3210 [PMID: 36245621 DOI: 10.21037/jtd-22-461]
- 8 **Karewicz A**, Faber K, Karon K, Januszewska K, Ryl J, Korczynski P, Gorska K, Dabrowska M, Krenke R. Evaluation of patients' satisfaction with bronchoscopy procedure. *PLoS One* 2022; **17**: e0274377 [PMID: 36201528 DOI: 10.1371/journal.pone.0274377]
- 9 **Madan K**, Biswal SK, Mittal S, Hadda V, Mohan A, Khilnani GC, Pandey RM, Guleria R. 1% Versus 2% Lignocaine for Airway Anesthesia in Flexible Bronchoscopy Without Lignocaine Nebulization (LIFE): A Randomized Controlled Trial. *J Bronchology Interv Pulmonol* 2018; **25**: 103-110 [PMID: 29346249 DOI: 10.1097/LBR.0000000000000458]
- 10 **Joskova M**, Durdik P, Sutovska M, Grendar M, Koniar D, Hargas L, Banovcin P, Franova S. Negative impact of anesthesia with midazolam, sufentanil, and propofol used in pediatric flexible bronchoscopy on the tracheal ciliary beat frequency in guinea pigs. *J Pharmacol Sci* 2020; **142**: 165-171 [PMID: 32033880 DOI: 10.1016/j.jphs.2020.01.005]
- 11 **Pritchett MA**, Lau K, Skibo S, Phillips KA, Bhadra K. Anesthesia considerations to reduce motion and atelectasis during advanced guided bronchoscopy. *BMC Pulm Med* 2021; **21**: 240 [PMID: 34273966 DOI: 10.1186/s12890-021-01584-6]
- 12 **Zhou Y**, Wu W, Zhu Y, Lv X, Liu J. Inhibition of stress and spontaneous respiration: Efficacy and safety of monitored anesthesia care by target-controlled infusion remifentanil in combination with dexmedetomidine in fiberoptic bronchoscopy for patients with severe tracheal stenosis. *Front Med (Lausanne)* 2022; **9**: 972066 [PMID: 36388940 DOI: 10.3389/fmed.2022.972066]



Published by **Baishideng Publishing Group Inc**  
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA  
**Telephone:** +1-925-3991568  
**E-mail:** [bpgoffice@wjgnet.com](mailto:bpgoffice@wjgnet.com)  
**Help Desk:** <https://www.f6publishing.com/helpdesk>  
<https://www.wjgnet.com>

