

World Journal of *Gastrointestinal Oncology*

World J Gastrointest Oncol 2021 September 15; 13(9): 980-1212



REVIEW

- 980** Involvement of integrin-activating peptides derived from tenascin-C in colon cancer progression
Fujita M, Suzuki H, Fukai F
- 995** MicroRNA expression in inflammatory bowel disease-associated colorectal cancer
Grillo TG, Quaglio AEV, Beraldo RF, Lima TB, Baima JP, Di Stasi LC, Sasaki LY
- 1017** Association between intestinal neoplasms and celiac disease: A review
Wang M, Yu M, Kong WJ, Cui M, Gao F
- 1029** Real-time fluorescence image-guided gastrointestinal oncologic surgery: Towards a new era
Martínez-López E, Martínez-Pérez A, Navarro-Martínez S, Sebastián-Tomás JC, de'Angelis N, García-Granero E
- 1043** Neoadjuvant chemotherapy for colorectal liver metastases: A contemporary review of the literature
Guo M, Jin N, Pawlik T, Cloyd JM

MINIREVIEWS

- 1062** Review of incomplete macroscopic resections (R2) in rectal cancer: Treatment, prognosis and future perspectives
Pérez Lara FJ, Hebrero Jimenez ML, Moya Donoso FJ, Hernández Gonzalez JM, Pitarch Martinez M, Prieto-Puga Arjona T
- 1073** Potential utility of liquid biopsies in the management of patients with biliary tract cancers: A review
Shotton R, Lamarca A, Valle J, McNamara MG
- 1086** Conservative management of malignant gastric outlet obstruction syndrome-evidence based evaluation of endoscopic ultrasound-guided gastroentero-anastomosis
Cominardi A, Tamanini G, Brighi N, Fusaroli P, Lisotti A
- 1099** Overgrowth of *Lactobacillus* in gastric cancer
Li ZP, Liu JX, Lu LL, Wang LL, Xu L, Guo ZH, Dong QJ
- 1109** Evidence based tools to improve efficiency of currently administered oncotherapies for tumors of the hepatopancreatobiliary system
Herold Z, Szasz AM, Dank M
- 1121** Screening strategy for gastrointestinal and hepatopancreatobiliary cancers in cystic fibrosis
Hoskins B, Wasuwanich P, Scheimann AO, Karnsakul W
- 1132** Immune aspects of hepatocellular carcinoma: From immune markers for early detection to immunotherapy
Mattos ÁZ, Debes JD, Boonstra A, Vogel A, Mattos AA

- 1144 Characterization of metabolic landscape in hepatocellular carcinoma

Wu J, Xue R, Jiang RT, Meng QH

- 1157 Effect of oncometabolic surgery on gastric cancer: The remission of hypertension, type 2 diabetes mellitus, and beyond

Cheng YX, Peng D, Tao W, Zhang W

ORIGINAL ARTICLE

Basic Study

- 1164 Scoparone inhibits pancreatic cancer through PI3K/Akt signaling pathway

Li N, Yang F, Liu DY, Guo JT, Ge N, Sun SY

Retrospective Study

- 1184 Prognostic value of modified Lauren classification in gastric cancer

Ning FL, Zhang NN, Wang J, Jin YF, Quan HG, Pei JP, Zhao Y, Zeng XT, Abe M, Zhang CD

META-ANALYSIS

- 1196 Neoadjuvant chemotherapy without radiation as a potential alternative treatment for locally advanced rectal cancer: A meta-analysis

Wu P, Xu HM, Zhu Z

LETTER TO THE EDITOR

- 1210 Use of liquid biopsies in gastrointestinal cancers

Khachfe HH

ABOUT COVER

Editorial Board Member of *World Journal of Gastrointestinal Oncology*, Rossana Berardi, MD, PhD, Director, Full Professor, Medical Oncology, Università Politecnica delle Marche, Ancona 60126, Italy. r.berardi@staff.univpm.it

AIMS AND SCOPE

The primary aim of *World Journal of Gastrointestinal Oncology* (WJGO, *World J Gastrointest Oncol*) is to provide scholars and readers from various fields of gastrointestinal oncology with a platform to publish high-quality basic and clinical research articles and communicate their research findings online.

WJGO mainly publishes articles reporting research results and findings obtained in the field of gastrointestinal oncology and covering a wide range of topics including liver cell adenoma, gastric neoplasms, appendiceal neoplasms, biliary tract neoplasms, hepatocellular carcinoma, pancreatic carcinoma, cecal neoplasms, colonic neoplasms, colorectal neoplasms, duodenal neoplasms, esophageal neoplasms, gallbladder neoplasms, etc.

INDEXING/ABSTRACTING

The WJGO is now indexed in Science Citation Index Expanded (also known as SciSearch®), PubMed, PubMed Central, and Scopus. The 2021 edition of Journal Citation Reports® cites the 2020 impact factor (IF) for WJGO as 3.393; IF without journal self cites: 3.333; 5-year IF: 3.519; Journal Citation Indicator: 0.5; Ranking: 163 among 242 journals in oncology; Quartile category: Q3; Ranking: 60 among 92 journals in gastroenterology and hepatology; and Quartile category: Q3. The WJGO's CiteScore for 2020 is 3.3 and Scopus CiteScore rank 2020: Gastroenterology is 70/136.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: *Ying-Yi Yuan*, Production Department Director: *Xiang Li*, Editorial Office Director: *Ya-Juan Ma*.

NAME OF JOURNAL

World Journal of Gastrointestinal Oncology

ISSN

ISSN 1948-5204 (online)

LAUNCH DATE

February 15, 2009

FREQUENCY

Monthly

EDITORS-IN-CHIEF

Rosa M Jimenez Rodriguez, Pashtoon Kasi, Monjur Ahmed, Florin Burada

EDITORIAL BOARD MEMBERS

<https://www.wjgnet.com/1948-5204/editorialboard.htm>

PUBLICATION DATE

September 15, 2021

COPYRIGHT

© 2021 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjgnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>



Retrospective Study

Prognostic value of modified Lauren classification in gastric cancer

Fei-Long Ning, Nan-Nan Zhang, Jun Wang, Yi-Feng Jin, Hong-Guang Quan, Jun-Peng Pei, Yan Zhao, Xian-Tao Zeng, Masanobu Abe, Chun-Dong Zhang

ORCID number: Fei-Long Ning 0000-0002-8846-3709; Nan-Nan Zhang 0000-0003-0582-8934; Jun Wang 0000-0002-6754-5641; Yi-Feng Jin 0000-0002-3803-2142; Hong-Guang Quan 0000-0002-9677-630X; Jun-Peng Pei 0000-0002-1876-3204; Yan Zhao 0000-0002-7760-916X; Xian-Tao Zeng 0000-0003-1262-725X; Masanobu Abe 0000-0001-9472-5077; Chun-Dong Zhang 0000-0003-1804-1356.

Author contributions: Ning FL, Zhang NN, Wang J, and Zhang CD contributed to the concept and design of the study; Ning FL, Zhang NN, and Zhang CD contributed to the acquisition of the data; Ning FL, Zhang NN, Wang J, Zeng XT, and Zhang CD contributed to the analysis and interpretation of the data; Ning FL, Zhang NN, Wang J, Jin YF, Quan HG, Pei JP, Zhao Y, Zeng XT, Abe M, and Zhang CD contributed to the drafting of the manuscript; Ning FL, Zhang CD contributed to the critical revision of the manuscript for important intellectual content; Zhang CD contributed to obtaining the funding.

Supported by The China Scholarship Council, No. 201908050148.

Institutional review board statement: This study was

Fei-Long Ning, Jun Wang, Hong-Guang Quan, Department of General Surgery, Xuzhou Hospital of Traditional Chinese Medicine, Xuzhou 221000, Jiangsu Province, China

Nan-Nan Zhang, State Key Laboratory of Cancer Biology and National Clinical Research Center for Digestive Diseases, Xijing Hospital of Digestive Diseases, The Fourth Military Medical University, Xi'an 710000, Shanxi Province, China

Yi-Feng Jin, Department of General Surgery, Jiading Hospital of Traditional Chinese Medicine, Shanghai 201800, China

Jun-Peng Pei, Chun-Dong Zhang, Department of Gastrointestinal Surgery, The Fourth Affiliated Hospital of China Medical University, Shenyang 110032, Liaoning Province, China

Yan Zhao, Department of Stomach Surgery, Liaoning Cancer Hospital and Institute, Cancer Hospital of China Medical University, Shenyang 110042, Liaoning Province, China

Xian-Tao Zeng, Center for Evidence-Based and Translational Medicine, Zhongnan Hospital of Wuhan University, Wuhan 430071, Hubei Province, China

Xian-Tao Zeng, Department of Evidence-Based Medicine and Clinical Epidemiology, The Second Clinical College of Wuhan University, Wuhan 430071, Hubei Province, China

Masanobu Abe, Division for Health Service Promotion, The University of Tokyo, Tokyo 113-0033, Japan

Chun-Dong Zhang, Department of Gastrointestinal Surgery, Graduate School of Medicine, The University of Tokyo, Tokyo 113-0033, Japan

Corresponding author: Chun-Dong Zhang, MD, Lecturer, Department of Gastrointestinal Surgery, The Fourth Affiliated Hospital of China Medical University, No. 4 Chongshan East Road, Huanggu District, Shenyang 110032, Liaoning Province, China.

cdzhang@cmu.edu.cn

Abstract

BACKGROUND

It remains controversial as to which pathological classification is most valuable in predicting the overall survival (OS) of patients with gastric cancer (GC).

AIM

To assess the prognostic performances of three pathological classifications in GC

approved by the Institutional Review Board of The Fourth Affiliated Hospital of China Medical University (EC-2021-KS-047).

Informed consent statement:

Informed consent statement was waived by the Institutional Review Board of The Fourth Affiliated Hospital of China Medical University.

Conflict-of-interest statement:

All authors declare no potential conflicts of interest for this manuscript.

Data sharing statement:

The datasets used and/or analyzed during the current study are available from the SEER database.

Open-Access:

This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

Manuscript source: Unsolicited manuscript

Specialty type: Gastroenterology and hepatology

Country/Territory of origin: China

Peer-review report's scientific quality classification

Grade A (Excellent): 0
Grade B (Very good): B
Grade C (Good): 0
Grade D (Fair): 0
Grade E (Poor): 0

Received: April 12, 2021

Peer-review started: April 12, 2021

First decision: June 23, 2021

Revised: July 1, 2021

Accepted: July 27, 2021

Article in press: July 27, 2021

and develop a novel prognostic nomogram for individually predicting OS.

METHODS

Patients were identified from the Surveillance, Epidemiology, and End Results program. Univariate and multivariate analyses were performed to identify the independent prognostic factors. Model discrimination and model fitting were evaluated by receiver operating characteristic curves and Akaike information criteria. Decision curve analysis was performed to assess clinical usefulness. The independent prognostic factors identified by multivariate analysis were further applied to develop a novel prognostic nomogram.

RESULTS

A total of 2718 eligible GC patients were identified. The modified Lauren classification was identified as one of the independent prognostic factors for OS. It showed superior model discriminative ability and model-fitting performance over the other pathological classifications, and similar results were obtained in various patient settings. In addition, it showed superior net benefits over the Lauren classification and tumor differentiation grade in predicting 3- and 5-year OS. A novel prognostic nomogram incorporating the modified Lauren classification showed superior model discriminative ability, model-fitting performance, and net benefits over the American Joint Committee on Cancer 8th edition tumor-node-metastasis classification.

CONCLUSION

The modified Lauren classification shows superior net benefits over the Lauren classification and tumor differentiation grade in predicting OS. A novel prognostic nomogram incorporating the modified Lauren classification shows good model discriminative ability, model-fitting performance, and net benefits.

Key Words: Gastric cancer; Pathological classification; Prognostic model; Tumor-node-metastasis classification; Survival outcome

©The Author(s) 2021. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: In this study, we compared the prognostic performances among the modified Lauren classification, the Lauren classification, and tumor differentiation grade. The modified Lauren classification was identified as one of the independent prognostic factors for overall survival. It showed superior model discriminative ability, model-fitting performance, and net benefits over the other classifications. We further developed a novel prognostic nomogram of individually predicting overall survival by incorporating the modified Lauren classification.

Citation: Ning FL, Zhang NN, Wang J, Jin YF, Quan HG, Pei JP, Zhao Y, Zeng XT, Abe M, Zhang CD. Prognostic value of modified Lauren classification in gastric cancer. *World J Gastrointest Oncol* 2021; 13(9): 1184-1195

URL: <https://www.wjgnet.com/1948-5204/full/v13/i9/1184.htm>

DOI: <https://dx.doi.org/10.4251/wjgo.v13.i9.1184>

INTRODUCTION

Gastric cancer (GC) is the fifth most prevalent and the third leading cause of cancer death worldwide[1]. It is a complex, heterogeneous entity that encompasses tumors with varying histopathologies, molecular profiles, and behaviors; however, GC is considered as a single entity for the purpose of clinical management and treatment, without regard to its subtype[2,3]. To date, the gold standard for GC prognostication and treatment guidance is the anatomical American Joint Committee on Cancer (AJCC) tumor-node-metastasis (TNM) classification[4,5]. It has been widely applied in many clinical practices without reference to its histopathology because the value of the morphological features of GC in determining clinical outcomes is still limited[6]. In

Published online: September 15, 2021

P-Reviewer: Cho JH

S-Editor: Zhang H

L-Editor: Wang TQ

P-Editor: Li X



addition, many investigators are still trying to identify a more valuable classification with better prognostic value[3,7,8].

Due to the wide variations in the morphological features of GC, many histological classifications have been proposed, and they are currently in wide use[3,9-13]. One of these classifications is the tumor differentiation grade. GC can be classified as well differentiated, moderately differentiated, poorly differentiated, and undifferentiated, according to the degree of differentiation exhibited by the tumor[10]. The tumor differentiation grade has been identified as a prognostic risk factor for GC in some studies [14,15]. However, several recent studies have reported that the tumor differentiation grade is not significantly associated with the prognosis of GC patients[16-19]. Another classification is the Lauren classification[13]. Despite dating back to 1965, it remains one of the most commonly used pathological classifications in GC. This classification categorizes GC into intestinal, diffuse, and mixed types, according to its histology, and each type has a distinct pathology and prognosis[13,20-22]. However, several studies have reported that the Lauren classification is not significantly correlated with patient survival because anatomic and corresponding epidemiologic distinctions were not taken into account[23,24].

Recently, it has been proposed that the Lauren classification be modified to include both the Lauren classification and the anatomical location of GC, thus yielding at least three entirely distinct types, namely, the proximal non-diffuse type, distal non-diffuse type, and diffuse type[3]. Molecular biology analyses further showed that there were marked differences in the mRNA expression profiles of the three types. Recent studies performed in Asia also suggested that the modified Lauren classification could be a reliable prognostic factor for patients with GC[25,26].

However, it remains controversial as to which pathological classification is most valuable in predicting the overall survival (OS) in GC patients. Therefore, we aimed to assess the prognostic value of the tumor differentiation grade, Lauren classification, and modified Lauren classification in GC patients. We compared model discriminative ability, model-fitting performance, and net benefits to identify the optimal prognostic pathological classification for GC based on the updated Surveillance, Epidemiology, and End Results (SEER) program. We also developed a novel prognostic nomogram for individually predicting the 3- and 5-year OS by applying the optimal pathological classification.

MATERIALS AND METHODS

Data source

We included data of eligible primary operable GC patients from the SEER program (<https://seer.cancer.gov/>). Data were extracted with SEER*Stat 8.3.6 software (www.seer.cancer.gov/seerstat). The data-use agreement for the SEER program data file was approved. This study was approved by the Institutional Review Boards of The Fourth Affiliated Hospital of China Medical University (EC-2021-KS-047).

Inclusion and exclusion criteria

Patients were included if they met the inclusion criteria as follows: (1) Primary carcinoma of the stomach; (2) TNM classification available; (3) no distant metastases (M0 disease); (4) solitary cancer; (5) history of curable surgery; (6) no neoadjuvant radiochemotherapy; (7) postoperative survival longer than one month; (8) aged between 18 and 75 years; (9) histological information available; and (10) defined tumor sites. Patients were excluded if they met any of the exclusion criteria as follows: (1) Metastatic carcinoma of the stomach; (2) TNM classification unavailable; (3) distant metastases (M1); (4) multiple cancers; (5) no history of surgery; (6) preoperative radiotherapy or chemotherapy; (7) postoperative survival shorter than 1 mo; (8) aged < 18 or > 75 years; (9) histological information unavailable; and (10) undefined tumor sites.

Clinicopathologic features

The analyzed clinicopathologic features were gender, age, tumor size, depth of tumor invasion (pT stage), number of retrieved lymph nodes, number of positive lymph nodes (pN stage), tumor differentiation grade, and Lauren classification. Patients were uniformly reviewed and re-staged (pT or pN stage) according to the AJCC 8th edition TNM classification[4]. The last follow-up was in November 2016.

Statistical analysis

The OS was calculated from the time of diagnosis to the time of death from any reason. Kaplan–Meier survival curves with log-rank tests were applied to analyze the difference in the OS among the groups. Factors with *P* values less than 0.1 in univariate analysis were considered potential prognostic factors and included in the Cox proportional hazards regression model. Hazard ratios with 95% CIs were applied.

The model discriminative ability of different pathological classifications was assessed by areas under the receiver operating characteristic curves (AUC)[27]. The model-fitting performance was evaluated by Akaike information criteria (AIC). A higher AUC value indicated a better model discriminative ability, and a lower AIC value indicated a superior model-fitting performance. The differences in AUC values were assessed by DeLong test[28]. Decision curve analysis (DCA) was performed to assess clinical usefulness, and the net benefits of making a decision based on the models were calculated[29,30].

The modified Lauren classification is an adjusted categorization of the Lauren classification, and both classifications are considered highly relevant. The Cox proportional hazards regression model was employed by incorporating either the Lauren or modified Lauren classification. Finally, the independent prognostic factors identified by multivariate analysis were applied to the nomogram.

Statistical analyses were performed using SPSS 22.0 (SPSS, Inc., Chicago, IL, United States), MedCalc 15.2 (Ostend, Belgium), GraphPad Prism 7 (GraphPad Software Inc., San Diego, CA, United States), and R 3.5.6 (<http://www.R-project.org/>) software packages. All tests were two-sided, and *P*-values less than 0.05 were considered statistically significant.

RESULTS

Patient characteristics

A total of 2718 eligible patients with GC from the SEER program were included. The clinicopathological characteristics are summarized in Table 1. There were 1588 males (58.4%) and 1130 were females (41.6%). The median age of all patients was 61 years (range, 18–75 years), and the median follow-up period was 31 mo (range, 2–155 mo).

Prognostic factors of overall survival

Univariate analysis identified potential prognostic factors, namely, age, tumor size, number of retrieved lymph nodes, pT stage, pN stage, tumor differentiation grade, and the modified Lauren classification (log-rank tests, $P < 0.10$). These factors were further applied in multivariate analysis with the Cox proportional hazards regression model. The results indicated that the independent prognostic factors predicting OS were age, tumor size, number of retrieved lymph nodes, pT stage, pN stage, and the modified Lauren classification (Table 2). However, neither the tumor differentiation grade ($P = 0.115$) nor the Lauren classification ($P = 0.163$) was found to be an independent predictive factor for OS in further multivariate analysis (Supplementary Table 1).

Predictive performance evaluations of pathological classifications

We compared the model discriminative ability and model-fitting performance of the tumor differentiation grade, Lauren classification, and modified Lauren classification. The modified Lauren classification showed superior model discriminative ability (3-year OS, AUC, 0.679 *vs* 0.666, DeLong test, $P < 0.001$; 5-year OS, AUC, 0.702 *vs* 0.681, $P < 0.001$) and model-fitting performance (AIC, 25877 *vs* 25923) over the Lauren classification (Table 3, Supplementary Figure 1A and B). The modified Lauren classification also showed superior model discriminative ability (3-year OS, AUC, 0.679 *vs* 0.626, DeLong test, $P < 0.001$; 5-year OS, AUC, 0.702 *vs* 0.621, $P < 0.001$) and model-fitting performance (AIC, 25877 *vs* 25971) over the tumor differentiation grade (Table 3, Supplementary Figure 1A and B). In addition, the Lauren classification showed superior model discriminative ability (3-year OS, AUC, 0.666 *vs* 0.626, DeLong test, $P < 0.001$; 5-year OS, AUC, 0.681 *vs* 0.621, $P < 0.001$) and model-fitting performance (AIC, 25923 *vs* 25971) over the tumor differentiation grade (Table 3, Supplementary Figure 1A and B, Supplementary Figure 2).

The modified Lauren classification also showed superior model discriminative ability (higher AUC values) and model-fitting performance (lower AIC values) in patients that were stratified by gender (female, male), age (< 60 years, ≥ 60 years),

Table 1 Basic characteristics according to anatomical location using the modified Lauren classification

Variable	Distal non-diffuse type	Proximal non-diffuse type	Diffuse type
Gender (%)			
Male	416 (63.6)	324 (70.0)	848 (53.0)
Female	238 (36.4)	139 (30.0)	753 (47.0)
Age (%)			
< 60 yr	200 (30.6)	167 (36.1)	850 (53.1)
≥ 60 yr	454 (69.4)	296 (63.9)	751 (46.9)
Tumor size (%)			
< 4.0 cm	318 (48.6)	218 (47.1)	664 (41.5)
≥ 4.0 cm	310 (47.4)	221 (47.7)	771 (48.2)
Unknown	26 (4.0)	24 (5.2)	166 (10.4)
Retrieved lymph nodes (%)			
Adequate ($n \geq 16$)	326 (49.8)	261 (56.4)	831 (51.9)
Inadequate ($n < 16$)	328 (50.2)	202 (43.6)	770 (48.1)
AJCC 8 th pT stage (%)			
pT1	211 (32.3)	123 (26.6)	356 (22.2)
pT2	87 (13.3)	69 (14.9)	173 (10.8)
pT3	207 (31.7)	149 (32.2)	464 (29.0)
pT4a	101 (15.4)	89 (19.2)	480 (30.0)
pT4b	48 (7.3)	33 (7.1)	128 (8.0)
AJCC 8 th pN stage (%)			
pN0	302 (46.2)	197 (42.5)	532 (33.2)
pN1	117 (17.9)	79 (17.1)	260 (16.2)
pN2	115 (17.6)	90 (19.4)	302 (18.9)
pN3a	95 (14.5)	70 (15.1)	347 (21.7)
pN3b	25 (3.8)	27 (5.8)	160 (10.0)
Differentiation grade (%)			
Well differentiation	66 (10.1)	26 (5.6)	3 (0.2)
Moderate differentiation	269 (41.1)	170 (36.7)	44 (2.7)
Poor differentiation	311 (47.6)	259 (55.9)	1484 (92.7)
Undifferentiation	8 (1.2)	8 (1.7)	70 (4.4)

AJCC: American Joint Committee on Cancer; pN stage: Pathological N stage; pT stage: Pathological T stage.

tumor size (< 4.0 cm, ≥ 4.0 cm, unknown), number of retrieved lymph nodes (< 16, ≥ 16), pT stage (pT1, pT2–4), and pN stage (pN0, pN1–3). These results confirmed that the modified Lauren classification showed the best model discriminative ability and model-fitting performance among the three pathological classifications.

Clinical utility of pathological classifications

We conducted DCA to assess the clinical utility of the different pathological classifications. The results revealed that the modified Lauren classification had superior net benefits over the Lauren classification and tumor differentiation grade in predicting both 3- and 5-year OS (Supplementary Figure 1C and D). Specifically, the modified Lauren classification showed superior net benefits over the tumor differentiation grade between threshold probabilities of 50%–65% and 40%–80% in predicting 3- and 5-year OS, respectively (Supplementary Figure 1C and D). In addition, the modified Lauren

Table 2 Univariate and multivariable analyses of prognostic factors for overall survival

Variable	No. of patients (%)	Univariate analysis		Multivariate analysis	
		5-yr OS	P value	HR (95%CI)	P value
Gender (%)			0.111		
Male	1588 (58.4)	45.9%			
Female	1130 (41.6)	49.1%			
Age (%)			< 0.001		< 0.001
< 60 yr	1217 (44.8)	50.7%		1 (Ref)	–
≥ 60 yr	1501 (55.2)	44.4%		1.157 (1.360–1.692)	< 0.001
Tumor size (%)			< 0.001		0.001
≤ 4.0 cm	1200 (44.2)	63.9%		1 (Ref)	–
> 4.0 cm	1302 (47.9)	33.6%		1.179 (1.038–1.339)	0.011
Unknown	216 (7.9)	40.2%		1.457 (1.191–1.782)	< 0.001
Retrieved lymph nodes (%)			0.074		< 0.001
Adequate ($n \geq 16$)	1418 (52.2)	48.9%		1 (Ref)	–
Inadequate ($n < 16$)	1300 (47.8)	45.5%		1.550 (1.380–1.740)	< 0.001
AJCC 8 th pT stage (%)			< 0.001		< 0.001
pT1	690 (25.4)	80.9%		1 (Ref)	–
pT2	329 (12.1)	66.6%		1.535 (1.193–1.975)	0.001
pT3	820 (30.2)	38.5%		2.882 (2.334–3.558)	< 0.001
pT4a	670 (24.7)	23.4%		3.415 (2.740–4.256)	< 0.001
pT4b	209 (7.7)	18.6%		4.452 (3.458–5.732)	< 0.001
AJCC 8 th pN stage (%)			< 0.001		< 0.001
pN0	1031 (37.9)	71.6%		1 (Ref)	–
pN1	456 (16.8)	46.9%		1.467 (1.225–1.757)	< 0.001
pN2	507 (18.7)	37.5%		1.611 (1.353–1.919)	< 0.001
pN3a	512 (18.8)	24.8%		2.356 (1.976–2.809)	< 0.001
pN3b	212 (7.8)	9.2%		4.138 (3.306–5.181)	< 0.001
Differentiation grade (%)			0.011		0.135
Well differentiation	95 (3.5)	69.4%		1 (Ref)	–
Moderate differentiation	483 (17.8)	58.9%		0.974 (0.649–1.462)	0.898
Poor differentiation	2054 (75.5)	44.0%		1.123 (0.755–1.670)	0.566
Undifferentiation	86 (3.2)	35.6%		1.415 (0.876–2.285)	0.156
Modified Lauren classification (%)			< 0.001		0.013
Distal non-diffuse type	654 (24.1)	58.8%		1 (Ref)	–
Proximal non-diffuse type	463 (17.0)	48.3%		1.230 (1.033–1.466)	0.020
Diffuse type	1601 (58.9)	42.4%		1.246 (1.068–1.452)	0.005

AJCC: American Joint Committee on Cancer; HR: Hazard ratio; OS: Overall survival; pN stage: Pathological N stage; pT stage: Pathological T stage. Variables with *P* values less than 0.1 were included in the multivariate analysis.

classification also showed superior net benefits over the Lauren classification between threshold probabilities of 30%–45% and 40%–60% in predicting 3- and 5-year OS, respectively (Supplementary Figure 1C and D).

Table 3 Comparison of predictive performances between different pathological classifications and prognostic models

Pathological classification/prognostic model	AUC (95%CI)		AIC
	3-yr overall survival	5-yr overall survival	
Differentiation grade	0.626 (0.608–0.644)	0.621 (0.602–0.639)	25971
Lauren classification	0.666 (0.647–0.683)	0.681 (0.663–0.699)	25923
Modified Lauren classification	0.679 (0.661–0.696)	0.702 (0.685–0.719)	25877
DeLong tests for AUCs			
Differentiation grade <i>vs</i> Lauren	$P < 0.001$	$P < 0.001$	–
Lauren <i>vs</i> modified Lauren	$P < 0.001$	$P < 0.001$	–
Modified Lauren <i>vs</i> differentiation grade	$P < 0.001$	$P < 0.001$	–
Novel prognostic model	0.803 (0.786–0.819)	0.804 (0.787–0.820)	20010
Age, tumor size, retrieved lymph nodes, pT stage, pN stage, modified Lauren classification			
Control model	0.776 (0.759–0.793)	0.776 (0.759–0.793)	20144
AJCC 8 th pTNM stage (pT stage, pN stage)			

AIC: Akaike's Information Criterion; AJCC: American Joint Committee on Cancer; AUC: Area under curve; pN stage: Pathological N stage; pT stage: Pathological T stage. A higher area under the curve indicated better model discrimination and a lower Akaike's Information Criterion indicates superior model-fitting; Differentiation grade, well *vs* moderate *vs* poor *vs* undifferentiation; Lauren classification, intestinal type *vs* diffuse type *vs* mixed type; Modified Lauren classification, distal non-diffuse *vs* proximal non-diffuse *vs* diffuse type.

Novel prognostic nomogram model vs AJCC 8th edition TNM classification

We further developed a novel prognostic model of age, tumor size, number of retrieved lymph nodes, pT stage, pN stage, and the modified Lauren classification by multivariate analysis using the Cox proportional hazards regression model. A novel nomogram for individually predicting 3- and 5-year OS was established by applying significant prognostic factors, including age, tumor size, number of retrieved lymph nodes, pT stage, pN stage, and the modified Lauren classification (Supplementary Figure 3A).

This novel prognostic model showed superior model discriminative ability (3-year OS, AUC, 0.803 *vs* 0.776, DeLong test; 5-year OS, AUC, 0.804 *vs* 0.776) and model-fitting performance (AIC, 20010 *vs* 20144) over the AJCC 8th edition TNM classification (pT stage, pN stage) (Table 3, Supplementary Figure 2C and D, Supplementary Figure 3B).

We further conducted DCA to assess the clinical utility of the novel prognostic model and the AJCC 8th edition TNM classification. The novel prognostic model showed superior net benefits over the AJCC 8th edition TNM classification between threshold probabilities of 40%–90% and 50%–95% in predicting 3- and 5-year OS, respectively (Supplementary Figure 1E and F).

DISCUSSION

Several pathological classifications of GC are currently in use due to the various morphological characteristics of GC[3,9-13]. However, it remains controversial as to which classification is best. Therefore, we performed a systematic analysis of the three most well-known pathological classifications and compared prognostic predictive performance with clinical use. In addition to the commonly used Lauren classification and tumor differentiation grade, we also compared a new classification, the modified Lauren classification. In our study, pN and pT stages were the most important prognostic factors for survival, thus validating the quality of the participants.

Tumor differentiation grades are commonly used for GC, and the four types of GC are defined as well differentiated, moderately differentiated, poorly differentiated, and undifferentiated[31]. It has been widely accepted that poorly differentiated tumors usually spread more extensively than well differentiated tumors by the time of surgery, and patients with more differentiated tumors have obvious survival advantages after curative resection[14,15]. However, recent studies have reported that

the tumor differentiation grade is not significantly associated with the prognosis of patients with GC[16-19]. In the current study, the tumor differentiation grade was significantly associated with the prognosis in log-rank tests; however, it was not an independent prognostic factor for OS. This discrepancy may be due to the mixture of differentiated and undifferentiated GC histologies[18,32]. In addition, it suggests that some well-differentiated types of GC can change to poorly differentiated types with tumor progression[33,34]. Therefore, further studies are needed to understand the significance of the tumor differentiation grade of GC.

The Lauren classification of GC is one of the most widely applied histological grading systems in predicting survival[21]. It has been reported that Lauren-classified tumor subtypes can respond differently to chemotherapy, thus yielding different survival outcomes[20]. However, the Lauren classification has also been demonstrated to have inadequate prognostic discriminative performance, and therefore, its prognostic accuracy remains controversial[23,24]. Specific pathogenetic and morphologic features of intestinal and diffuse types may underlie their different behaviors[22]. Population-based studies have reported the different epidemiological features of Lauren-classified subtypes and cancer of the cardia[35,36]. Epidemiologically, the intestinal type of GC, particularly that of the antrum, is often strongly associated with chronic inflammation as a consequence of chronic infection with *Helicobacter pylori*[37,38]. Anatomically, proximal GC can be classified as a third type of GC for which inflammation of a different type may be the driving force for carcinogenesis[39]. Furthermore, the anatomical location of GC is clinically relevant, and proximal third GC is associated with a worse prognosis than middle or distal third GC [40,41].

Therefore, a location-modified Lauren classification has been proposed. It defines the subtypes of GC by incorporating epidemiological and histopathological data together with the anatomical location[3]. Several studies have revealed that the modified Lauren classification has better discriminative ability and monotonicity than the Lauren classification[25,26]. The results of the current study demonstrated that the modified Lauren classification showed superior model discriminative ability, model-fitting performance, and net benefits compared with other classifications. Similar findings were also obtained in populations stratified by gender, age, tumor size, number of retrieved lymph nodes, pT stage, and pN stage. Decision curve analysis confirmed its clinical usefulness over other classifications.

It remains unclear why the modified Lauren classification showed a significantly better prognostic performance. A previous study has reported that the Kirsten rat sarcoma viral oncogene homolog pathway was downregulated in proximal non-diffuse GC compared with diffuse GC[42]. In addition, genomic analysis has confirmed that the modified Lauren classification can achieve a clear molecular distinction[3]. Moreover, HER2 amplification or overexpression is not uniform across different GC subtypes; it is most prevalent in proximal GC (HER2 positivity rate, approximately 30%) and least prevalent in diffuse GC (HER2 positivity rate, approximately 5%)[43]. Furthermore, whole-genome sequencing of diffuse GC uncovered mutations in *RHOA*, a gene encoding a well-studied small GTPase, in 15%–25% of diffuse tumors but not in non-diffuse tumors[44].

Nomograms are visualization tools for individually predicting survival[45,46] with improved predictive accuracy and comprehensive outcomes for many types of cancers [47]. Therefore, we developed a novel prognostic nomogram of age, tumor size, number of retrieved lymph nodes, pT stage, pN stage, and the modified Lauren classification. This novel prognostic model achieved superior model discriminative ability, model-fitting performance, and net benefits over the AJCC 8th edition TNM classification. These findings support the consideration of more factors spanning different aspects of the disease as the most promising approach to improve the clinical management of GC. However, the findings of the current study still need to be interpreted with caution because specific intervention factors of the surgical procedures, chemo-radiotherapeutic regimens, and drug doses were not applied in the current study.

CONCLUSION

In summary, the modified Lauren classification provides superior model discriminative ability, model-fitting performance, and net benefits over the tumor differentiation grade and Lauren classification. It also shows good applicability in various clinical settings. The novel prognostic nomogram incorporating the modified Lauren

classification shows good model discriminative ability, model-fitting performance, and net benefits. However, the findings of the current study require further validation.

ARTICLE HIGHLIGHTS

Research background

It remains controversial as to which pathological classification is most valuable in predicting overall survival (OS) in patients with gastric cancer (GC).

Research motivation

Recently, it has been proposed that the Lauren classification be modified to include both the Lauren classification and the anatomical location of GC, thus yielding at least three entirely distinct types, namely, the proximal non-diffuse type, distal non-diffuse type, and diffuse type.

Research objectives

To assess the prognostic performances of three pathological classifications in GC and develop a novel prognostic nomogram for individually predicting OS.

Research methods

We retrospectively reviewed and analyzed the data identified from the Surveillance, Epidemiology, and End Results program.

Research results

A total of 2718 eligible GC patients were identified. The modified Lauren classification was identified as one of the independent prognostic factors for OS. It showed superior model discriminative ability and model-fitting performance over the other pathological classifications, and similar results were obtained in various patient settings. In addition, it showed superior net benefits over the Lauren classification and tumor differentiation grade in predicting 3- and 5-year OS. A novel prognostic nomogram incorporating the modified Lauren classification showed superior model discriminative ability, model-fitting performance, and net benefits over the American Joint Committee on Cancer 8th edition tumor-node-metastasis classification.

Research conclusions

The modified Lauren classification shows superior net benefits over the Lauren classification and tumor differentiation grade in predicting OS. A novel prognostic nomogram incorporating the modified Lauren classification shows good model discriminative ability, model-fitting performance, and net benefits.

Research perspectives

A large prospective study is needed to validate our findings.

ACKNOWLEDGEMENTS

We acknowledge the efforts of Surveillance, Epidemiology, and End Results (SEER) program tumor registries for creating the SEER database (<https://seer.cancer.gov/>).

REFERENCES

- 1 Siegel RL, Miller KD, Jemal A. Cancer statistics, 2019. *CA Cancer J Clin* 2019; **69**: 7-34 [PMID: 30620402 DOI: 10.3322/caac.21551]
- 2 Shah MA, Kelsen DP. Gastric cancer: a primer on the epidemiology and biology of the disease and an overview of the medical management of advanced disease. *J Natl Compr Canc Netw* 2010; **8**: 437-447 [PMID: 20410336 DOI: 10.6004/jnccn.2010.0033]
- 3 Shah MA, Khanin R, Tang L, Janjigian YY, Klimstra DS, Gerdes H, Kelsen DP. Molecular classification of gastric cancer: a new paradigm. *Clin Cancer Res* 2011; **17**: 2693-2701 [PMID: 21430069 DOI: 10.1158/1078-0432.CCR-10-2203]
- 4 Amin MB, Edge SB, Greene F, Byrd DR, Brookland RK, Washington MK, Gershenwald JE, Compton CC, Hess KR, Sullivan DC, Jessup JM, Brierley JD, Gaspar LE, Schilsky RL, Balch CM,

- Winchester DP, Asare EA, Madera M, Gress DM, Meyer LR. AJCC cancer staging manual. 8th ed. New York: Springer, 2017
- 5 **Japanese Gastric Cancer Association.** Japanese gastric cancer treatment guidelines 2018 (5th edition). *Gastric Cancer* 2021; **24**: 1-21 [PMID: [32060757](#) DOI: [10.1007/s10120-020-01042-y](#)]
- 6 **Dai W**, Mo S, Xiang W, Han L, Li Q, Wang R, Xu Y, Cai G. The Critical Role of Tumor Size in Predicting Prognosis for T1 Colon Cancer. *Oncologist* 2020; **25**: 244-251 [PMID: [32162825](#) DOI: [10.1634/theoncologist.2019-0469](#)]
- 7 **Turner ES**, Turner JR. Expanding the Lauren classification: a new gastric cancer subtype? *Gastroenterology* 2013; **145**: 505-508 [PMID: [23891604](#) DOI: [10.1053/j.gastro.2013.07.019](#)]
- 8 **Songun I**, van de Velde CJ, Arends JW, Blok P, Grond AJ, Offerhaus GJ, Hermans J, van Krieken JH. Classification of gastric carcinoma using the Goseki system provides prognostic information additional to TNM staging. *Cancer* 1999; **85**: 2114-2118 [PMID: [10326687](#) DOI: [10.1002/\(sici\)1097-0142\(19990515\)85:10<2114::aid-cncr3>3.0.co;2-u](#)]
- 9 **Japanese Gastric Cancer Association.** Japanese classification of gastric carcinoma: 3rd English edition. *Gastric Cancer* 2011; **14**: 101-112 [PMID: [21573743](#) DOI: [10.1007/s10120-011-0041-5](#)]
- 10 **Hirota T**, Itabashi M, Suzuki K, Yoshida S. Clinicopathologic study of minute and small early gastric cancer. Histogenesis of gastric cancer. *Pathol Annu* 1980; **15**: 1-19 [PMID: [6256703](#)]
- 11 **Goseki N**, Takizawa T, Koike M. Differences in the mode of the extension of gastric cancer classified by histological type: new histological classification of gastric carcinoma. *Gut* 1992; **33**: 606-612 [PMID: [1377153](#) DOI: [10.1136/gut.33.5.606](#)]
- 12 **Jass JR**, Sobin LH, Watanabe H. The World Health Organization's histologic classification of gastrointestinal tumors. A commentary on the second edition. *Cancer* 1990; **66**: 2162-2167 [PMID: [2171747](#) DOI: [10.1002/1097-0142\(19901115\)66:10<2162::aid-cncr2820661020>3.0.co;2-n](#)]
- 13 **Lauren P.** The two histological main types of gastric carcinoma: diffuse and so-called intestinal-type carcinoma. An attempt at a histo-clinical classification. *Acta Pathol Microbiol Scand* 1965; **64**: 31-49 [PMID: [14320675](#) DOI: [10.1111/apm.1965.64.1.31](#)]
- 14 **Chu MP**, Hecht JR, Slamon D, Wainberg ZA, Bang YJ, Hoff PM, Sobrero A, Qin S, Afenjar K, Houe V, King K, Koski S, Mulder K, Hiller JP, Scarfe A, Spratlin J, Huang YJ, Khan-Wasti S, Chua N, Sawyer MB. Association of Proton Pump Inhibitors and Capecitabine Efficacy in Advanced Gastroesophageal Cancer: Secondary Analysis of the TRIO-013/LOGiC Randomized Clinical Trial. *JAMA Oncol* 2017; **3**: 767-773 [PMID: [27737436](#) DOI: [10.1001/jamaoncol.2016.3358](#)]
- 15 **Van Cutsem E**, Sagaert X, Topal B, Haustermans K, Prenen H. Gastric cancer. *Lancet* 2016; **388**: 2654-2664 [PMID: [27156933](#) DOI: [10.1016/S0140-6736\(16\)30354-3](#)]
- 16 **Bonnot PE**, Piessen G, Kepenekian V, Decullier E, Pocard M, Meunier B, Bereder JM, Abboud K, Marchal F, Quenet F, Goere D, Msika S, Arvieux C, Pirro N, Wernert R, Rat P, Gagnière J, Lefevre JH, Courvoisier T, Kianmanesh R, Vaudoyer D, Rivoire M, Meeus P, Passot G, Glehen O; FREGAT and BIG-RENAPE Networks. Cyto-reductive Surgery With or Without Hyperthermic Intraperitoneal Chemotherapy for Gastric Cancer With Peritoneal Metastases (CYTO-CHIP study): A Propensity Score Analysis. *J Clin Oncol* 2019; **37**: 2028-2040 [PMID: [31084544](#) DOI: [10.1200/JCO.18.01688](#)]
- 17 **Zhang CD**, Shen MY, Zhang JK, Ning FL, Zhou BS, Dai DQ. Prognostic significance of distal subtotal gastrectomy with standard D2 and extended D2 lymphadenectomy for locally advanced gastric cancer. *Sci Rep* 2015; **5**: 17273 [PMID: [26602830](#) DOI: [10.1038/srep17273](#)]
- 18 **Feng F**, Liu J, Wang F, Zheng G, Wang Q, Liu S, Xu G, Guo M, Lian X, Zhang H. Prognostic value of differentiation status in gastric cancer. *BMC Cancer* 2018; **18**: 865 [PMID: [30176846](#) DOI: [10.1186/s12885-018-4780-0](#)]
- 19 **Jiang Y**, Li T, Liang X, Hu Y, Huang L, Liao Z, Zhao L, Han Z, Zhu S, Wang M, Xu Y, Qi X, Liu H, Yang Y, Yu J, Liu W, Cai S, Li G. Association of Adjuvant Chemotherapy With Survival in Patients With Stage II or III Gastric Cancer. *JAMA Surg* 2017; **152**: e171087 [PMID: [28538950](#) DOI: [10.1001/jamasurg.2017.1087](#)]
- 20 **Jiménez Fonseca P**, Carmona-Bayonas A, Hernández R, Custodio A, Cano JM, Lacalle A, Echavarría I, Macías I, Mangas M, Visa L, Buxo E, Álvarez Manceño F, Viudez A, Pericay C, Azkarate A, Ramchandani A, López C, Martínez de Castro E, Fernández Montes A, Longo F, Sánchez Bayona R, Limón ML, Díaz-Serrano A, Martín Carnicero A, Arias D, Cerdà P, Rivera F, Vieitez JM, Sánchez Cánovas M, Garrido M, Gallego J. Lauren subtypes of advanced gastric cancer influence survival and response to chemotherapy: real-world data from the AGAMENON National Cancer Registry. *Br J Cancer* 2017; **117**: 775-782 [PMID: [28765618](#) DOI: [10.1038/bjc.2017.245](#)]
- 21 **Lee JH**, Chang KK, Yoon C, Tang LH, Strong VE, Yoon SS. Lauren Histologic Type Is the Most Important Factor Associated With Pattern of Recurrence Following Resection of Gastric Adenocarcinoma. *Ann Surg* 2018; **267**: 105-113 [PMID: [27759618](#) DOI: [10.1097/SLA.0000000000002040](#)]
- 22 **Pernot S**, Terme M, Radosevic-Robin N, Castan F, Badoual C, Marcheteau E, Penault-Llorca F, Bouche O, Bennouna J, Francois E, Ghiringhelli F, De La Fouchardiere C, Samalin E, Baptiste Bachet J, Borg C, Boige V, Voron T, Stanbury T, Tartour E, Gourgou S, Malka D, Taieb J. Infiltrating and peripheral immune cell analysis in advanced gastric cancer according to the Lauren classification and its prognostic significance. *Gastric Cancer* 2020; **23**: 73-81 [PMID: [31267360](#) DOI: [10.1007/s10120-019-00983-3](#)]
- 23 **Huang SC**, Ng KF, Yeh TS, Cheng CT, Lin JS, Liu YJ, Chuang HC, Chen TC. Subtraction of Epstein-Barr virus and microsatellite instability genotypes from the Lauren histotypes: Combined molecular and histologic subtyping with clinicopathological and prognostic significance validated in a

- cohort of 1,248 cases. *Int J Cancer* 2019; **145**: 3218-3230 [PMID: [30771224](#) DOI: [10.1002/ijc.32215](#)]
- 24 **de Aguiar VG**, Segatelli V, Macedo ALV, Goldenberg A, Gansl RC, Maluf FC, Usón Junior PLS. Signet ring cell component, not the Lauren subtype, predicts poor survival: an analysis of 198 cases of gastric cancer. *Future Oncol* 2019; **15**: 401-408 [PMID: [30620220](#) DOI: [10.2217/fon-2018-0354](#)]
- 25 **Choi JK**, Park YS, Jung DH, Son SY, Ahn SH, Park DJ, Kim HH. Clinical Relevance of the Tumor Location-Modified Lauren Classification System of Gastric Cancer. *J Gastric Cancer* 2015; **15**: 183-190 [PMID: [26468416](#) DOI: [10.5230/jgc.2015.15.3.183](#)]
- 26 **Zhao LY**, Wang JJ, Zhao YL, Chen XZ, Yang K, Chen XL, Zhang WH, Liu K, Song XH, Zheng JB, Zhou ZG, Yu PW, Li Y, Hu JK. Superiority of Tumor Location-Modified Lauren Classification System for Gastric Cancer: A Multi-Institutional Validation Analysis. *Ann Surg Oncol* 2018; **25**: 3257-3263 [PMID: [30051368](#) DOI: [10.1245/s10434-018-6654-8](#)]
- 27 **Zweig MH**, Campbell G. Receiver-operating characteristic (ROC) plots: a fundamental evaluation tool in clinical medicine. *Clin Chem* 1993; **39**: 561-577 [PMID: [8472349](#)]
- 28 **DeLong ER**, DeLong DM, Clarke-Pearson DL. Comparing the areas under two or more correlated receiver operating characteristic curves: a nonparametric approach. *Biometrics* 1988; **44**: 837-845 [PMID: [3203132](#)]
- 29 **Fitzgerald M**, Saville BR, Lewis RJ. Decision curve analysis. *JAMA* 2015; **313**: 409-410 [PMID: [25626037](#) DOI: [10.1001/jama.2015.37](#)]
- 30 **Vickers AJ**, Elkin EB. Decision curve analysis: a novel method for evaluating prediction models. *Med Decis Making* 2006; **26**: 565-574 [PMID: [17099194](#) DOI: [10.1177/0272989X06295361](#)]
- 31 **Camargo MC**, Kim WH, Chiaravalli AM, Kim KM, Corvalan AH, Matsuo K, Yu J, Sung JJ, Herrera-Goepfert R, Meneses-Gonzalez F, Kijima Y, Natsugoe S, Liao LM, Lissowska J, Kim S, Hu N, Gonzalez CA, Yatabe Y, Koriyama C, Hewitt SM, Akiba S, Gullet ML, Taylor PR, Rabkin CS. Improved survival of gastric cancer with tumour Epstein-Barr virus positivity: an international pooled analysis. *Gut* 2014; **63**: 236-243 [PMID: [23580779](#) DOI: [10.1136/gutjnl-2013-304531](#)]
- 32 **Horiuchi Y**, Fujisaki J, Yamamoto N, Ishizuka N, Omae M, Ishiyama A, Yoshio T, Hirasawa T, Yamamoto Y, Nagahama M, Takahashi H, Tsuchida T. Mixed poorly differentiated adenocarcinoma in undifferentiated-type early gastric cancer predicts endoscopic noncurative resection. *Gastric Cancer* 2018; **21**: 689-695 [PMID: [29236187](#) DOI: [10.1007/s10120-017-0788-4](#)]
- 33 **Tanaka K**, Shimura T, Kitajima T, Kondo S, Ide S, Okugawa Y, Saigusa S, Toiyama Y, Inoue Y, Araki T, Uchida K, Mohri Y, Kusunoki M. Tropomyosin-related receptor kinase B at the invasive front and tumour cell dedifferentiation in gastric cancer. *Br J Cancer* 2014; **110**: 2923-2934 [PMID: [24853179](#) DOI: [10.1038/bjc.2014.228](#)]
- 34 **Friedmann-Morvinski D**, Verma IM. Dedifferentiation and reprogramming: origins of cancer stem cells. *EMBO Rep* 2014; **15**: 244-253 [PMID: [24531722](#) DOI: [10.1002/embr.201338254](#)]
- 35 **Kaneko S**, Yoshimura T. Time trend analysis of gastric cancer incidence in Japan by histological types, 1975-1989. *Br J Cancer* 2001; **84**: 400-405 [PMID: [11161407](#) DOI: [10.1054/bjoc.2000.1602](#)]
- 36 **Dassen AE**, Lemmens VE, van de Poll-Franse LV, Creemers GJ, Breninkmeijer SJ, Lips DJ, Vd Wurff AA, Bosscha K, Coebergh JW. Trends in incidence, treatment and survival of gastric adenocarcinoma between 1990 and 2007: a population-based study in the Netherlands. *Eur J Cancer* 2010; **46**: 1101-1110 [PMID: [20219351](#) DOI: [10.1016/j.ejca.2010.02.013](#)]
- 37 **Choi IJ**, Kim CG, Lee JY, Kim YI, Kook MC, Park B, Joo J. Family History of Gastric Cancer and *Helicobacter pylori* Treatment. *N Engl J Med* 2020; **382**: 427-436 [PMID: [31995688](#) DOI: [10.1056/NEJMoa1909666](#)]
- 38 **Ajani JA**, Lee J, Sano T, Janjigian YY, Fan D, Song S. Gastric adenocarcinoma. *Nat Rev Dis Primers* 2017; **3**: 17036 [PMID: [28569272](#) DOI: [10.1038/nrdp.2017.36](#)]
- 39 **Shoji Y**, Nunobe S, Ida S, Kumagai K, Ohashi M, Sano T, Hiki N. Surgical outcomes and risk assessment for anastomotic complications after laparoscopic proximal gastrectomy with double-flap technique for upper-third gastric cancer. *Gastric Cancer* 2019; **22**: 1036-1043 [PMID: [30838469](#) DOI: [10.1007/s10120-019-00940-0](#)]
- 40 **Rosa F**, Quero G, Fiorillo C, Bissolati M, Cipollari C, Rausei S, Chiari D, Ruspi L, de Manzoni G, Costamagna G, Doglietto GB, Alfieri S. Total vs proximal gastrectomy for adenocarcinoma of the upper third of the stomach: a propensity-score-matched analysis of a multicenter western experience (On behalf of the Italian Research Group for Gastric Cancer-GIRCG). *Gastric Cancer* 2018; **21**: 845-852 [PMID: [29423892](#) DOI: [10.1007/s10120-018-0804-3](#)]
- 41 **Kajiyama Y**, Tsurumaru M, Udagawa H, Tsutsumi K, Kinoshita Y, Ueno M, Akiyama H. Prognostic factors in adenocarcinoma of the gastric cardia: pathologic stage analysis and multivariate regression analysis. *J Clin Oncol* 1997; **15**: 2015-2021 [PMID: [9164213](#) DOI: [10.1200/JCO.1997.15.5.2015](#)]
- 42 **Hiyama T**, Haruma K, Kitadai Y, Masuda H, Miyamoto M, Tanaka S, Yoshihara M, Shimamoto F, Chayama K. K-ras mutation in helicobacter pylori-associated chronic gastritis in patients with and without gastric cancer. *Int J Cancer* 2002; **97**: 562-566 [PMID: [11807778](#) DOI: [10.1002/ijc.1644](#)]
- 43 **Bang YJ**, Van Cutsem E, Feyereislova A, Chung HC, Shen L, Sawaki A, Lordick F, Ohtsu A, Omuro Y, Satoh T, Aprile G, Kulikov E, Hill J, Lehle M, Rüschhoff J, Kang YK, ToGA Trial Investigators. Trastuzumab in combination with chemotherapy versus chemotherapy alone for treatment of HER2-positive advanced gastric or gastro-oesophageal junction cancer (ToGA): a phase 3, open-label, randomised controlled trial. *Lancet* 2010; **376**: 687-697 [PMID: [20728210](#) DOI: [10.1016/S0140-6736\(10\)61121-X](#)]
- 44 **Kakiuchi M**, Nishizawa T, Ueda H, Gotoh K, Tanaka A, Hayashi A, Yamamoto S, Tatsuno K, Katoh H, Watanabe Y, Ichimura T, Ushiku T, Funahashi S, Tateishi K, Wada I, Shimizu N, Nomura S,

- Koike K, Seto Y, Fukayama M, Aburatani H, Ishikawa S. Recurrent gain-of-function mutations of RHOA in diffuse-type gastric carcinoma. *Nat Genet* 2014; **46**: 583-587 [PMID: [24816255](#) DOI: [10.1038/ng.2984](#)]
- 45 **Randall RL**, Cable MG. Nominal nomograms and marginal margins: what is the law of the line? *Lancet Oncol* 2016; **17**: 554-556 [PMID: [27301026](#) DOI: [10.1016/S1470-2045\(16\)00072-3](#)]
- 46 **Balachandran VP**, Gonen M, Smith JJ, DeMatteo RP. Nomograms in oncology: more than meets the eye. *Lancet Oncol* 2015; **16**: e173-e180 [PMID: [25846097](#) DOI: [10.1016/S1470-2045\(14\)71116-7](#)]
- 47 **Carmona-Bayonas A**, Jiménez-Fonseca P, Lamarca Á, Barriuso J, Castaño Á, Benavent M, Alonso V, Riesco-Martínez MDC, Alonso-Gordoa T, Custodio A, Sánchez Cánovas M, Hernando Cubero J, López C, Lacasta A, Fernández Montes A, Marazuela M, Crespo G, Escudero P, Díaz JÁ, Feliciangeli E, Gallego J, Llanos M, Segura Á, Vilardell F, Percovich JC, Grande E, Capdevila J, Valle JW, García-Carbonero R. Prediction of Progression-Free Survival in Patients With Advanced, Well-Differentiated, Neuroendocrine Tumors Being Treated With a Somatostatin Analog: The GETNE-TRASGU Study. *J Clin Oncol* 2019; **37**: 2571-2580 [PMID: [31390276](#) DOI: [10.1200/JCO.19.00980](#)]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

Telephone: +1-925-3991568

E-mail: bpgoffice@wjgnet.com

Help Desk: <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

