

Contrast-enhanced colour-Doppler sonography versus pH ...

<https://www.researchgate.net/publication/5373762...>

Oral US contrast agent administration has only been described in one study involving the diagnosis of gastroesophageal reflux [28]. The procedures, uses and safety of IV and intravesical contrast ...

Sonographic assessments of gastrointestinal and biliary ...

https://www.researchgate.net/publication/26275391_Sonographic_assessments_of...

The aim of this study was to evaluate the feasibility of using swallow contrast-enhanced ultrasound (swallow-CEUS) to diagnose ZD. Ten consecutive patients with ZD (7 men and 3 women, aged 67 ...

2017 Scientific Program - 2017 - Journal of Ultrasound in ...

<https://onlinelibrary.wiley.com/doi/full/10.1002/jum.14288>

Contrast-enhanced ultrasound (lymphosonography) was performed 0, 0.25, 0.5, 1, 2, 4, 6, and 24 hours post-Sonazoid administration to identify SLNs using Cadence Pulse Sequencing on an S3000 scanner (Siemens Healthineers, Mountain View, CA) with a 12L4 probe.

Radiographic and Endoscopic Diagnosis and Treatment of ...

https://www.researchgate.net/publication/51614124_Radiographic_and_Endoscopic...

Both intravenous and oral contrast should be administered if possible, given the increased accuracy of delayed contrast-enhanced CT images [19, 20]. Imaging 40-60 min after ingestion has been ...

Gastrointestinal Ultrasound In Neonates, Infants And ...

<https://www.researchgate.net/publication/261989061...>

The study of the gastrointestinal tract by imaging, particularly using ultrasound, is a required instrument for diagnosis of acute and chronic gastrointestinal pathologies in pediatric age.

Clinical Performance of PET/CT in Evaluation of Cancer ...

<https://jnm.snmjournals.org/content/44/8/1200> ▾

This study assessed the clinical performance of a combined PET/CT system using 18F-FDG in oncologic patients. Methods: 18F-FDG PET/CT was used to evaluate 204 patients with 586 suspicious lesions. All patients had available follow-up data, enabling assessment of the clinical significance of hybrid PET/CT findings. Differences in interpretation between PET, CT, and fused PET/CT data were ...

国内版

国际版

Microsoft Bing

Contrast-enhanced ultrasound using SonoVue mixe

624 Results

Any time

Open links in new tab

ALL

IMAGES

VIDEOS

Contrast-enhanced colour-Doppler sonography versus pH...

https://www.researchgate.net/publication/5373762...

Oral US contrast agent administration has only been described in one study involving the diagnosis of gastroesophageal reflux [28]. The procedures, uses and safety of IV and intravesical contrast ...

Sonographic assessments of gastrointestinal and biliary ...

https://www.researchgate.net/publication/26275391...

The aim of this study was to evaluate the feasibility of using swallow contrast-enhanced ultrasound (swallow-CEUS) to diagnose ZD. Ten consecutive patients with ZD (7 men and 3 women, aged 67 ...

2017 Scientific Program - 2017 - Journal of Ultrasound in ...

https://onlinelibrary.wiley.com/doi/full/10.1002/jum.14288

Contrast-enhanced ultrasound (lymphosonography) was performed 0, 0.25, 0.5, 1, 2, 4, 6, and 24 hours post-Sonazoid administration to identify SLNs using Cadence Pulse Sequencing on an S3000 scanner (Siemens Healthineers, Mountain View, CA) with a 12L4 probe.

Radiographic and Endoscopic Diagnosis and Treatment of ...

https://www.researchgate.net/publication/51614124

Search Tools

Turn off Hover Translation (关闭翻译)

激活 Windows

转到“设置”以激活 Windows。

Chat with Bing

Sign in



61671-Manuscript-File-revision.doc

Quotes Included
Bibliography Included1%
SIMILAR

Name of Journal: *World Journal of Clinical Cases*

Manuscript NO: 61671

Manuscript Type: CASE REPORT

Contrast-enhanced ultrasound using SonoVue mixed with oral gastrointestinal contrast agent to evaluate esophageal hiatal hernia: Three-cases report

Wang J *et al.* Diagnosing EHH by CEUS

Jingyu Wang, Yan Luo, Wenying Wang, Shicheng Zheng, Lian He, Chunyan Xie, Li Peng

Match Overview

- 1 Crossref 13 words
Jiabing Zhang, Nan Li, Feihong Dong, Jian An, Jue Zhang, Yunkun Luo. "Ultrasound diffraction attenuation microscopy in ... <1%
- 2 Crossref 12 words
Georg R. Linke. "Efficacy of Laparoscopic Mesh-Augmented Hiataloplasty in GERD and Symptomatic Hiatal Hernia. Stur ... <1%



PAGE: 1 OF 11



Text-Only Report

国内版 国际版

Contrast-enhanced ultrasound using SonoVue mixed with oral gastri



ALL IMAGES VIDEOS

2,590 Results Any time ▾

Contrast-enhanced colour-Doppler sonography versus pH ...

<https://www.researchgate.net/publication/5373762...>

The aim of this study was to evaluate the accuracy of contrast-enhanced colour-Doppler ultrasound (CDUS) in the diagnosis of gastro-oesophageal reflux in children.

Sonographic assessments of gastrointestinal and biliary ...

https://www.researchgate.net/publication/26275391_Sonographic_assessments_of...

The aim of this study was to evaluate the feasibility of using swallow contrast-enhanced ultrasound (swallow-CEUS) to diagnose ZD. Ten consecutive patients with ZD (7 men and 3 women, aged 67 ...

Radiographic and Endoscopic Diagnosis and Treatment of ...

https://www.researchgate.net/publication/51614124_Radiographic_and_Endoscopic...

Both intravenous and oral contrast should be administered if possible, given the increased accuracy of delayed contrast-enhanced CT images [19, 20]. Imaging 40-60 min after ingestion has been ...

Sonographic diagnosis features of Zenker diverticulum ...

<https://www.researchgate.net/publication/44798451...>

The aim of this study was to evaluate the feasibility of using swallow contrast-enhanced ultrasound (swallow-CEUS) to diagnose ZD. Ten consecutive patients with ZD (7 men and 3 women, aged 67 ...

Oral Contrast Echocardiography for the Diagnosis of Left ...

<https://www.researchgate.net/publication/23481546...>

Using two- (2DTTE) and three-dimensional transthoracic echocardiography (3DTTE) and an oral contrast agent (a carbonated beverage), a mass-like lesion behind the left ventricular posterior wall in ...

ESPR 2017 | SpringerLink

<https://link.springer.com/article/10.1007/s00247-017-3820-2> ▾

Apr 26, 2017 · Others US technologies are used in children to better approach the disease activity. It includes, hydrosonography using specific oral agents (Mannitol, Sorbitol, PolyEthylene Glycol, etc...), contrast-enhanced ultrasound and dynamic contrast-enhanced ultrasound (nowadays, contrast agent is off-label in children) (34) and elastography (35).

Oral presentations | SpringerLink