

Yersinia type III effectors perturb host innate immune responses

Pha K *et al.* *Yersinia* effectors perturb host immune responses

Khavong Pha, Lorena Navarro

Khavong Pha, Lorena Navarro, Department of ¹⁶Microbiology and Molecular Genetics, University of California, Davis, Davis, CA 95616, United States

Author contributions: Pha K and Navarro L solely contributed to this paper.

Supported by the ASM Robert D. Watkins Graduate Fellowship and UC Davis

Match Overview

- | | | |
|---|---|-----|
| 1 | CrossCheck 26 words
Grishin, Andrey M., Ksenia A. Beyrakhova, and Miroslaw Cygler. "Structural insight into effector proteins of gram..." | <1% |
| 2 | CrossCheck 15 words
Goure, Julien, Petr Broz, Olivier Attree, Guy R. Cornelis, and Ina Attree. "Protective Anti - V Antibodies Inhibit Pse ..." | <1% |
| 3 | Internet 14 words
crawled on 10-Mar-2010
www.rbej.com | <1% |
| 4 | CrossCheck 14 words
Hacker, Jörg, and James B. Kaper. "Pathogenicity Islands and the Evolution of Microbes", Annual Review of Micro | <1% |
| 5 | Internet 14 words
crawled on 04-Nov-2009
www.dtic.mil | <1% |
| 6 | CrossCheck 13 words
Hiroko Tsutsui. "The TLR4/TRIF-Mediated Activation of NLRP3 Inflammasome Underlies Endotoxin-Induced Lh ..." | <1% |

Modulation of innate immune responses by Yersinia type III ...

www.researchgate.net/publication/247770428_Modulation_of_innate_immune_responses

Publication » .Modulation of innate immune responses by Yersinia type III secretion system translocators and ... Yersinia species translocates seven effectors into infected host cells. example of a perturbation that is sensed as a pattern of

Modulation of innate immune responses by Yersinia type III ...

www.ncbi.nlm.nih.gov/pmc/articles/PMC3788085/

29 Jul 2013 ... Modulation of innate immune responses by Yersinia type III secretion ... type III secretion systems (T3SSs) that translocate effectors into host cells. ... to the cytosol is an example of a perturbation that is sensed as a pattern of....

Type III Secretion Systems and Disease

www.ncbi.nlm.nih.gov/pmc/articles/PMC2176049/

Type III secretion systems (T3SSs) are complex bacterial structures that This manipulation can result in both the induction and inhibition of host innate immune signaling. ... The T3SS effectors of Salmonella, Yersinia, and Shigella target these ... T3SSs are involved in the perturbation of host cellular immune responses.

Yersinia type III secretion

jcb.rupress.org/content/158/3/401.full

5 Aug 2002 ... The injected Yops perturb the dynamics of the cytoskeleton, disrupting ... The Yop virulon is an archetype of the type III secretion systems (TTSSs)* Translocation of the Yop effectors into the host cell requires the Thus, YopH not only contributes to Yersinia's evasion of the innate immune response by....

Innate Immune Recognition of Yersinia pseudotuberculosis Type III ...

journals.plos.org/plospathogens/article?id=10.1371/journal.ppat.1000686

4 Dec 2009 ... A common result of this "injection" is perturbation of normal host processes, ... T3SS effector Yops presumably travel through the type III needle and ... TLR activation generates strong innate immune responses, many of which....

Inflammasome Activation in Response to the Yersinia Type III ... - mBio

mbio.asm.org/content/6/1/e02095-14.full

17 Feb 2015 ... Type III secretion systems (T3SS) translocate effector proteins into ... The innate immune system plays a crucial role in host defense against pathogens. or YopD alters the

学术搜索

找到约 834 条结果 (用时0.06秒)

文章

我的图书馆

时间不限

2015以来

2014以来

2011以来

自定义范围...

按相关性排序

按日期排序

搜索所有网页

中文网页

简体中文网页

☒ 包括专利☒ 包含引用☐ 创建快讯小提示: 只搜索**中文(简体)**结果, 可在 [学术搜索设置](#) 指定搜索语言[Subterfuge and manipulation: type III effector proteins of phytopathogenic bacteria](#)SR Grant, EJ Fisher, JH Chang, BM Mole... - Annu. Rev. ..., 2006 - [annualreviews.org](#)... Table 2 lists **type III effector** proteins from plant pathogens with known or predicted enzymatic function ... Several **types** of proteases have been identified ... P. syringae homologs to the **Yersinia** protease YopT include HopN1 (HopPtoN), HopAR1 (AvrPphB), and HopC1 (HopPtoC) (25 ...

被引用次数: 351 相关文章 所有 9 个版本 引用 保存

[unc.edu](#) 中的 [\[PDF\]](#)[Yersinia type III secretion send in the effectors](#)GR Cornelis - The Journal of cell biology, 2002 - [jcb.rupress.org](#)... not least, YopP/YopJ induces apoptosis of macrophages but not of other cell **types** (Denecker et ... integrate in the bacterial genome in such a way that they received a **type III** secretion signal ... Yops not only leads to a better understanding of the mechanism of **Yersinia** infection but ...

被引用次数: 380 相关文章 所有 13 个版本 引用 保存

[rupress.org](#) 中的 [\[HTML\]](#)[\[HTML\] Arabidopsis RIN4 is a target of the type III virulence effector AvrRpt2 and modulates RPS2-mediated resistance](#) [sciencedirect.com](#) 中的 [\[HTML\]](#)

D Mackey, Y Belkadir, JM Alonso, JR Ecker, JL Dangl - Cell, 2003 - Elsevier

... avrRpm1 and avrRpt2 was inoculated into leaves of Col-0, rps2, rpm1/rps2, or 3 independent transgenic ... We proposed that the targeting of RIN4 by the **type III effectors** is monitored by RPM1, leading to rapid ... (B) AvrRpt2 is delivered into the plant cell via **type III** secretion and ...

被引用次数: 642 相关文章 所有 15 个版本 引用 保存

[Type III secretion systems and disease](#)

B Coburn, I Sekirov, BB Finlay - Clinical microbiology reviews, 2007 - Am Soc Microbiol

... This bacterial **type III** secretion system (T3SS) is of special interest to those studying **host**-pathogen interactions because, by utilizing ... which is also known to be involved in **type 2** secretion and in the assembly of **type IV** bacterial ... **Yersinia** pestis is the causative agent of plague.[asm.org](#) 中的 [\[HTML\]](#)

[网页](#)[图片](#)[视频](#)[新闻](#)[更多 ▾](#)[搜索工具](#)

找到约 50,600 条结果 (用时 0.71 秒)

Google 学术: Yersinia type III effectors perturb host innate immune responses

Subterfuge and manipulation: type III effector proteins ... - Grant - 被引用次数: 352

Yersinia type III secretion send in the effectors - Cornelis - 被引用次数: 382

... RIN4 is a target of the type III virulence effector ... - Mackey - 被引用次数: 644

"Modulation of innate immune responses by Yersinia type III ...

www.researchgate.net/.../247770428_Modulation_of_innate_i... - 翻译此页

Yersinia enterocolitica and Yersinia pseudotuberculosis induce macrophage apoptosis and subvert host innate immunity by injecting effectors through the T3SS ...

Modulation of innate immune responses by Yersinia type III ...

www.ncbi.nlm.nih.gov > ... > Literature > PubMed Central (PMC) - 翻译此页

2013年7月29日 - Modulation of innate immune responses by Yersinia type III secretion ...

type III secretion systems (T3SSs) that translocate effectors into host cells. ... to the cytosol is an example of a perturbation that is sensed as a pattern of ...

Innate Immune Recognition of Yersinia pseudotuberculosis ...

www.ncbi.nlm.nih.gov > ... > PubMed Central (PMC) ▾ 翻译此页

作者: V Auerbuch - 2009 - 被引用次数: 46 - 相关文章

2009年12月4日 - Innate Immune Recognition of Yersinia pseudotuberculosis Type III

Secretion A common result of this "injection" is perturbation of normal host processes, to the ... Targeting of effector Yops into the host cell cytosol requires the ... TLR activation generates strong innate immune responses, many of which ...

Yersinia type III secretion - The Journal of Cell Biology

jcb.rupress.org > 2002 Archive > 5 August ▾ 翻译此页