

World Journal of *Clinical Cases*

World J Clin Cases 2022 July 6; 10(19): 6341-6758



MINIREVIEWS

- 6341** Review of clinical characteristics, immune responses and regulatory mechanisms of hepatitis E-associated liver failure
Chen C, Zhang SY, Chen L
- 6349** Current guidelines for *Helicobacter pylori* treatment in East Asia 2022: Differences among China, Japan, and South Korea
Cho JH, Jin SY
- 6360** Review of epidermal growth factor receptor-tyrosine kinase inhibitors administration to non-small-cell lung cancer patients undergoing hemodialysis
Lan CC, Hsieh PC, Huang CY, Yang MC, Su WL, Wu CW, Wu YK

ORIGINAL ARTICLE**Case Control Study**

- 6370** Pregnancy-related psychopathology: A comparison between pre-COVID-19 and COVID-19-related social restriction periods
Chieffo D, Avallone C, Serio A, Kotzalidis GD, Balocchi M, De Luca I, Hirsch D, Gonzalez del Castillo A, Lanzotti P, Marano G, Rinaldi L, Lanzone A, Mercuri E, Mazza M, Sani G
- 6385** Intestinal mucosal barrier in functional constipation: Dose it change?
Wang JK, Wei W, Zhao DY, Wang HF, Zhang YL, Lei JP, Yao SK

Retrospective Cohort Study

- 6399** Identification of risk factors for surgical site infection after type II and type III tibial pilon fracture surgery
Hu H, Zhang J, Xie XG, Dai YK, Huang X

Retrospective Study

- 6406** Total knee arthroplasty in Ranawat II valgus deformity with enlarged femoral valgus cut angle: A new technique to achieve balanced gap
Lv SJ, Wang XJ, Huang JF, Mao Q, He BJ, Tong PJ
- 6417** Preliminary evidence in treatment of eosinophilic gastroenteritis in children: A case series
Chen Y, Sun M
- 6428** Self-made wire loop snare successfully treats gastric persimmon stone under endoscopy
Xu W, Liu XB, Li SB, Deng WP, Tong Q
- 6437** Neoadjuvant transcatheter arterial chemoembolization and systemic chemotherapy for the treatment of undifferentiated embryonal sarcoma of the liver in children
He M, Cai JB, Lai C, Mao JQ, Xiong JN, Guan ZH, Li LJ, Shu Q, Ying MD, Wang JH

- 6446 Effect of cold snare polypectomy for small colorectal polyps

Meng QQ, Rao M, Gao PJ

- 6456 Field evaluation of COVID-19 rapid antigen test: Are rapid antigen tests less reliable among the elderly?

Tabain I, Cucevic D, Skreb N, Mrzljak A, Ferencak I, Hruskar Z, Mistic A, Kuzle J, Skoda AM, Jankovic H, Vilibic-Cavlek T

Observational Study

- 6464 Tracheobronchial intubation using flexible bronchoscopy in children with Pierre Robin sequence: Nursing considerations for complications

Ye YL, Zhang CF, Xu LZ, Fan HF, Peng JZ, Lu G, Hu XY

- 6472 Family relationship of nurses in COVID-19 pandemic: A qualitative study

Çelik MY, Kiliç M

META-ANALYSIS

- 6483 Diagnostic accuracy of ≥ 16 -slice spiral computed tomography for local staging of colon cancer: A systematic review and meta-analysis

Liu D, Sun LM, Liang JH, Song L, Liu XP

CASE REPORT

- 6496 Delayed-onset endophthalmitis associated with *Achromobacter* species developed in acute form several months after cataract surgery: Three case reports

Kim TH, Lee SJ, Nam KY

- 6501 Sustained dialysis with misplaced peritoneal dialysis catheter outside peritoneum: A case report

Shen QQ, Behera TR, Chen LL, Attia D, Han F

- 6507 Arteriovenous thrombotic events in a patient with advanced lung cancer following bevacizumab plus chemotherapy: A case report

Kong Y, Xu XC, Hong L

- 6514 Endoscopic ultrasound radiofrequency ablation of pancreatic insulinoma in elderly patients: Three case reports

Rossi G, Petrone MC, Capurso G, Partelli S, Falconi M, Arcidiacono PG

- 6520 Acute choroidal involvement in lupus nephritis: A case report and review of literature

Yao Y, Wang HX, Liu LW, Ding YL, Sheng JE, Deng XH, Liu B

- 6529 Triple A syndrome-related achalasia treated by per-oral endoscopic myotomy: Three case reports

Liu FC, Feng YL, Yang AM, Guo T

- 6536 Choroidal thickening with serous retinal detachment in BRAF/MEK inhibitor-induced uveitis: A case report

Kiraly P, Groznic AL, Valentinčić NV, Mekjavić PJ, Urbančić M, Ocvirk J, Mesti T

- 6543 Esophageal granular cell tumor: A case report

Chen YL, Zhou J, Yu HL

- 6548** Hem-o-lok clip migration to the common bile duct after laparoscopic common bile duct exploration: A case report
Liu DR, Wu JH, Shi JT, Zhu HB, Li C
- 6555** Chidamide and sintilimab combination in diffuse large B-cell lymphoma progressing after chimeric antigen receptor T therapy
Hao YY, Chen PP, Yuan XG, Zhao AQ, Liang Y, Liu H, Qian WB
- 6563** Relapsing polychondritis with isolated tracheobronchial involvement complicated with Sjogren's syndrome: A case report
Chen JY, Li XY, Zong C
- 6571** Acute methanol poisoning with bilateral diffuse cerebral hemorrhage: A case report
Li J, Feng ZJ, Liu L, Ma YJ
- 6580** Immunoabsorption therapy for Klinefelter syndrome with antiphospholipid syndrome in a patient: A case report
Song Y, Xiao YZ, Wang C, Du R
- 6587** Roxadustat for treatment of anemia in a cancer patient with end-stage renal disease: A case report
Zhou QQ, Li J, Liu B, Wang CL
- 6595** Imaging-based diagnosis for extraskeletal Ewing sarcoma in pediatrics: A case report
Chen ZH, Guo HQ, Chen JJ, Zhang Y, Zhao L
- 6602** Unusual course of congenital complete heart block in an adult: A case report
Su LN, Wu MY, Cui YX, Lee CY, Song JX, Chen H
- 6609** Penile metastasis from rectal carcinoma: A case report
Sun JJ, Zhang SY, Tian JJ, Jin BY
- 6617** Isolated cryptococcal osteomyelitis of the ulna in an immunocompetent patient: A case report
Ma JL, Liao L, Wan T, Yang FC
- 6626** Magnetic resonance imaging features of intrahepatic extramedullary hematopoiesis: Three case reports
Luo M, Chen JW, Xie CM
- 6636** Giant retroperitoneal liposarcoma treated with radical conservative surgery: A case report and review of literature
Lieto E, Cardella F, Erario S, Del Sorbo G, Reginelli A, Galizia G, Urraro F, Panarese I, Auricchio A
- 6647** Transplanted kidney loss during colorectal cancer chemotherapy: A case report
Pośpiech M, Kolonko A, Nieszporek T, Kozak S, Kozaczka A, Karkoszka H, Winder M, Chudek J
- 6656** Massive gastrointestinal bleeding after endoscopic rubber band ligation of internal hemorrhoids: A case report
Jiang YD, Liu Y, Wu JD, Li GP, Liu J, Hou XH, Song J

- 6664** Mills' syndrome is a unique entity of upper motor neuron disease with N-shaped progression: Three case reports
Zhang ZY, Ouyang ZY, Zhao GH, Fang JJ
- 6672** Entire process of electrocardiogram recording of Wellens syndrome: A case report
Tang N, Li YH, Kang L, Li R, Chu QM
- 6679** Retroperitoneal tumor finally diagnosed as a bronchogenic cyst: A case report and review of literature
Gong YY, Qian X, Liang B, Jiang MD, Liu J, Tao X, Luo J, Liu HJ, Feng YG
- 6688** Successful treatment of Morbihan disease with total glucosides of paeony: A case report
Zhou LF, Lu R
- 6695** Ant sting-induced whole-body pustules in an inebriated male: A case report
Chen SQ, Yang T, Lan LF, Chen XM, Huang DB, Zeng ZL, Ye XY, Wan CL, Li LN
- 6702** Plastic surgery for giant metastatic endometrioid adenocarcinoma in the abdominal wall: A case report and review of literature
Wang JY, Wang ZQ, Liang SC, Li GX, Shi JL, Wang JL
- 6710** Delayed-release oral mesalamine tablet mimicking a small jejunal gastrointestinal stromal tumor: A case report
Frosio F, Rausa E, Marra P, Boutron-Ruault MC, Lucianetti A
- 6716** Concurrent alcoholic cirrhosis and malignant peritoneal mesothelioma in a patient: A case report
Liu L, Zhu XY, Zong WJ, Chu CL, Zhu JY, Shen XJ
- 6722** Two smoking-related lesions in the same pulmonary lobe of squamous cell carcinoma and pulmonary Langerhans cell histiocytosis: A case report
Gencer A, Ozcibik G, Karakas FG, Sarbay I, Batur S, Borekci S, Turna A
- 6728** Proprotein convertase subtilisin/kexin type 9 inhibitor non responses in an adult with a history of coronary revascularization: A case report
Yang L, Xiao YY, Shao L, Ouyang CS, Hu Y, Li B, Lei LF, Wang H
- 6736** Multimodal imaging study of lipemia retinalis with diabetic retinopathy: A case report
Zhang SJ, Yan ZY, Yuan LF, Wang YH, Wang LF
- 6744** Primary squamous cell carcinoma of the liver: A case report
Kang LM, Yu DP, Zheng Y, Zhou YH
- 6750** Tumor-to-tumor metastasis of clear cell renal cell carcinoma to contralateral synchronous pheochromocytoma: A case report
Wen HY, Hou J, Zeng H, Zhou Q, Chen N

ABOUT COVER

Editorial Board Member of *World Journal of Clinical Cases*, Abdulqadir Jeprel Naswhan, MSc, RN, Director, Research Scientist, Senior Lecturer, Senior Researcher, Nursing for Education and Practice Development, Hamad Medical Corporation, Doha 576214, Qatar. anashwan@hamad.qa

AIMS AND SCOPE

The primary aim of *World Journal of Clinical Cases* (*WJCC*, *World J Clin Cases*) is to provide scholars and readers from various fields of clinical medicine with a platform to publish high-quality clinical research articles and communicate their research findings online.

WJCC mainly publishes articles reporting research results and findings obtained in the field of clinical medicine and covering a wide range of topics, including case control studies, retrospective cohort studies, retrospective studies, clinical trials studies, observational studies, prospective studies, randomized controlled trials, randomized clinical trials, systematic reviews, meta-analysis, and case reports.

INDEXING/ABSTRACTING

The *WJCC* is now indexed in Science Citation Index Expanded (also known as SciSearch®), Journal Citation Reports/Science Edition, Scopus, PubMed, and PubMed Central. The 2021 Edition of Journal Citation Reports® cites the 2020 impact factor (IF) for *WJCC* as 1.337; IF without journal self cites: 1.301; 5-year IF: 1.742; Journal Citation Indicator: 0.33; Ranking: 119 among 169 journals in medicine, general and internal; and Quartile category: Q3. The *WJCC*'s CiteScore for 2020 is 0.8 and Scopus CiteScore rank 2020: General Medicine is 493/793.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: *Xu Guo*; Production Department Director: *Xiang Li*; Editorial Office Director: *Jin-Lei Wang*.

NAME OF JOURNAL

World Journal of Clinical Cases

ISSN

ISSN 2307-8960 (online)

LAUNCH DATE

April 16, 2013

FREQUENCY

Thrice Monthly

EDITORS-IN-CHIEF

Bao-Gan Peng, Jerzy Tadeusz Chudek, George Kontogeorgos, Maurizio Serati, Ja Hyeon Ku

EDITORIAL BOARD MEMBERS

<https://www.wjgnet.com/2307-8960/editorialboard.htm>

PUBLICATION DATE

July 6, 2022

COPYRIGHT

© 2022 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjgnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>

Retrospective Study

Total knee arthroplasty in Ranawat II valgus deformity with enlarged femoral valgus cut angle: A new technique to achieve balanced gap

Shuai-Jie Lv, Xiao-Jian Wang, Jie-Feng Huang, Qiang Mao, Bang-Jian He, Pei-Jian Tong

Specialty type: Orthopedics**Provenance and peer review:**

Unsolicited article; Externally peer reviewed.

Peer-review model: Single blind**Peer-review report's scientific quality classification**

Grade A (Excellent): 0

Grade B (Very good): 0

Grade C (Good): C

Grade D (Fair): 0

Grade E (Poor): 0

P-Reviewer: Abulsoud MI, Egypt**Received:** July 23, 2021**Peer-review started:** July 23, 2021**First decision:** September 2, 2021**Revised:** September 10, 2021**Accepted:** April 21, 2022**Article in press:** April 21, 2022**Published online:** July 6, 2022

Shuai-Jie Lv, Jie-Feng Huang, Qiang Mao, Bang-Jian He, Pei-Jian Tong, Department of Orthopedics and Traumatology, The First Affiliated Hospital and First Clinical College of Zhejiang Chinese Medical University, Hangzhou 310006, Zhejiang Province, China

Xiao-Jian Wang, The First Clinical Medical School, Zhejiang Chinese Medical University, Hangzhou 310053, Zhejiang Province, China

Corresponding author: Pei-Jian Tong, MD, Chief Doctor, Department of Orthopedics and Traumatology, The First Affiliated Hospital and First Clinical College of Zhejiang Chinese Medical University, No. 54 Youdian Road, Shangcheng District, Hangzhou 310006, Zhejiang Province, China. 351653161@qq.com

Abstract

BACKGROUND

Nearly 10% of patients undergoing primary total knee arthroplasty (TKA) have valgus deformity (VD) of the knee. For severe VD of the knee, a more lateral structural release is needed to achieve balance between medial and lateral space and neutral femorotibial mechanical axis (FTMA), which is challenging and technical.

AIM

To introduce a new surgical technique of resection, soft tissue release, and FTMA for Ranawat type-II VD with a 5-year follow-up.

METHODS

A retrospective study was conducted on patients who underwent TKA from December 2011 to December 2014. Hip-knee-ankle (HKA), range of motion (ROM), Oxford knee score (OKS), and knee society score (KSS) were used to assess the joint activity of patients in the new theory TKA group (NT-TKA) and were compared with those of the conventional TKA group (C-TKA).

RESULTS

A total of 103 people (103 knees) were included in this study, including 42 patients with an average follow-up period of 83 mo in the C-TKA group and 61 patients with an average follow-up period of 76 mo in the NT-TKA group. Six patients had constrained prosthesis, one had common peroneal nerve injury, and two had joint instability in the C-TKA group, but none of these occurred in the NT-TKA group. There were significant statistical differences in constrained

prosthesis usage and complications between the groups ($P = 0.002$ and $P = 0.034$, respectively). The KSS at 1 mo post-operation for the C-TKA and NT-TKA groups were 11.2 ± 3.8 and 13.3 ± 2.9 , respectively, with a significant difference ($P = 0.007$). However, the data of HKA, ROM, OKS KSS, and prosthesis survival rate were insignificant ($P > 0.05$) in both the preoperative and follow-up periods.

CONCLUSION

Adopting 5° - 7° valgus cut angle for VD and sacrificing 2° neutral FTMA for severe VD which cannot be completely corrected during TKA can reduce the need for soft tissue release, maintain early joint stability, reduce the use of constrained prostheses, and minimize postoperative complications.

Key Words: Valgus, knee; Arthroplasty; Osteotomy; Mechanical axis; Bone and soft-tissue balance

©The Author(s) 2022. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: This study aimed to provide a new theory of surgical resection, soft tissue release, and femorotibial mechanical axis (FTMA) for Ranawat type-II valgus deformity (VD) of the knee using bone and soft tissue co-balance theory to reduce the complexity of the surgery and to improve postoperative outcomes. Bone and soft tissue balance is equally important in a total knee arthroplasty procedure. Adopting 5° - 7° valgus cut angle for VD and sacrificing 2° neutral FTMA for severe VD during surgery can reduce the need for soft tissue release, maintain joint stability, and abrogate the use of constrained prostheses while minimizing postoperative complications.

Citation: Lv SJ, Wang XJ, Huang JF, Mao Q, He BJ, Tong PJ. Total knee arthroplasty in Ranawat II valgus deformity with enlarged femoral valgus cut angle: A new technique to achieve balanced gap. *World J Clin Cases* 2022; 10(19): 6406-6416

URL: <https://www.wjgnet.com/2307-8960/full/v10/i19/6406.htm>

DOI: <https://dx.doi.org/10.12998/wjcc.v10.i19.6406>

INTRODUCTION

Nearly 10% of patients undergoing primary total knee arthroplasty (TKA) have valgus deformity (VD) of the knee. These deformities are commonly accompanied by the lateral femur and tibia bone defect, lateral soft tissue tightening, and medial soft-tissue laxity. Ranawat *et al*[1] pioneered the classification of VD of the knee into three types: I, II, and III. TKA is indicated for type I and can be corrected with relative ease. Type II is more challenging and technical, and type III usually requires a constrained prosthesis.

To improve clinical outcomes, several studies have described the surgical approach, soft tissue balance, and resection. Rossi *et al*[2] proposed that a valgus cut angle (VCA) of 3° is a favorable approach. However, the femoral condyle requires greater medial than lateral resection to restore the femorotibial mechanical axis (FTMA), which may result in a hypoplastic lateral condyle[3]. Lateral soft tissue releases are required to balance a VD of the knee. Moreover, there are concerns that excessive soft tissue release could result in joint instability, and therefore, reducing the destruction of lateral soft tissue to protect joint stability is of great significance during TKA. Neutral FTMA, as one of the gold standards for TKA, is universally accepted to improve both postoperative knee function and prosthesis survival rate. However, some scholars[4,5] proposed that the alignment position of the FTMA may have little effect on TKA and the prosthesis survival rate. For severe VD of the knee, more soft tissue destruction and even restrictive prosthesis may be required to achieve neutral FTMA. Therefore, the balance between osteotomy, soft tissue release, and FTMA is very critical and valuable.

This study aimed to provide a new theory of surgical resection, soft tissue release, and FTMA for Ranawat type-II VD of the knee using the theory of bone and soft tissue co-balance to reduce the complexity of the surgery and to improve postoperative outcomes.

MATERIALS AND METHODS

Patients

This retrospectively study reviewed the data of patients who underwent conventional TKA or new technique TKA at the Department of Orthopedics and Traumatology at the First Affiliated Hospital of Zhejiang Chinese Medical University from January 2011 to December 2014. Patients with Ranawat type-II VD caused by either osteoarthritis, rheumatoid arthritis, or hemophilic arthritis were included in this study. There was no age limit for enrolled patients. A basic understanding of the score sheet and consent for physical examination were also required. Ranawat type-II was defined as the femorotibial axis (FTA) $> 10^\circ$ on anteroposterior radiographs upon standing. Patients with combined joint immobilization, infection, old fracture, or a previous resection were excluded from the study. This study was approved by the hospital ethics and review committee of the First Affiliated Hospital of Zhejiang Chinese Medical University (2019-K-207-01). All patients provided informed consent for this study.

Surgical procedure

All patients were routinely given prophylactic antibiotics within 30 min before surgery. The supine position and tourniquet were used after a general anesthetic. The incision site was at the mid-level knee *via* a medial patella approach. For the distal femur resection in the new theory TKA group (NT-TKA), VCA was increased by 2° based on the traditional 3° , and the medial femoral condyle determined the resection thickness. A neutral resection was used routinely in the proximal tibia (Figure 1). Next, the surgeon extended the knee joint, evaluated the extension gap through a spacer block, and then performed joint space balance according to the straightening priority. Osteophytes that affect soft tissue balance were resected in a cuff-like manner, releasing soft tissue attachment point. Pie-crusting release was then performed on the lateral collateral ligament (LCL), the iliotibial band (ITB), and the posterolateral capsule (PLC) with a maximum of eight stitches through a 20 mL syringe needle. If the joint space was difficult to balance, the VCA was expanded to 7° with continuous release of soft tissues until the knee joint extension gap was in a rectangular shape (Figures 1 and 2A). External rotation resection and a "4 in 1" guide resection were performed at the 90° position of knee flexion. The external rotation resection angle of the anterior femoral condyle can be appropriately increased, usually 3° - 6° , depending on the severity of the knee valgus. If the knee flexion was tense, the cuff-like release of the anterior soft tissue was performed (Figure 2B). For patients with an unsuccessful correction of the neutral FTMA, a 2° valgus FTMA was sacrificed to reduce the risk of ligament failure, joint instability, and consumption of constrained prostheses.

In the conventional TKA group (C-TKA), the distal femur cut bone angle was set to 3° , and the soft tissue was released according to the above steps until the joint space was balanced and the FTMA returned to 0° . Once the resection and the soft tissue balance were done, the posterior stabilized components were inserted. If the knee was unstable, legacy constrained condylar knee (LCKK) components were installed. If there was a femoral or tibial bone defect, remedial measures such as bone cement, small bone blocks, screws, and bone grafting were taken according to the degree of the defect. The patella was not resurfaced, but osteophyte cleaning and denervation were achieved. Finally, the surgeon stitched and bandaged the incision site. All knee prostheses were purchased from Zimmer (Zimmer, NexGen, LPS/LCKK, Warsaw, United States). All procedures were performed by the same orthopedic surgeon with more than 30 years of clinical experience.

Postoperative management

Postoperatively, prophylactic low molecular weight heparin (0.4 mL/4100 IU, 1 vial a day) was initiated and maintained for 5 wk for prevention of deep venous thrombosis or VTE. Cefuroxime sodium (1.5 g, twice a day) was given and maintained for 1 wk to prevent infection. In addition, all patients received physiotherapy and rehabilitation training under the guidance of a professional rehabilitation specialist. At day 1 post-surgery, patients were allowed to walk with full weight-bearing under the protection of crutches until the patient was confident walking independently.


Postoperative effect evaluation

Hip-knee-ankle (HKA), range of motion (ROM), Oxford knee score (OKS), and knee society score (KSS) were used to evaluate the joint function of patients before and after surgery. Any adverse events were recorded during the postoperative follow-up period.

Data collection and follow-up

The sample sizing was calculated by KSS. The data of patients enrolled in this study were accessed *via* an electronic medical record system by two investigators who received standardized training. Main biodata included age, gender, body mass index, diagnosis, FTMA, and operation time.

All patients were discharged from the hospital 2 wk after surgery and were followed by telephone and outpatient clinic visits at 1, 3, and 6 mo for the first year, and then an annual telephone or outpatient follow-up for 3 years. Physical examination and routine radiographs (X-ray) of the surgical site were performed at each follow-up visit by the attending physician. For patients living in far-flung areas, the



DOI: 10.12998/wjcc.v10.i19.6406 Copyright ©The Author(s) 2022.

Figure 1 Process of total knee arthroplasty. A: Preoperative X-ray of the knee joint in anterior and lateral position; B: The distal femoral valgus osteotomy angle was 7° (red arrow); C: The thickness of the medial femoral condyle osteotomy was more than that of the lateral condyle; D: Tibial plateau osteotomy was taken as the medial tibial plateau reference; E: The osteophyte of the lateral tibial plateau was removed, and the lateral plateau bone defect (black circle) was visible; F: Release of the lateral collateral ligament; G: The thickness of distal femoral osteotomy was increased because the extension space was smaller than the flexion space; H: Postoperative X-ray of the knee joint in anterior and lateral position.

X-ray results from the local hospital were accepted.


Statistical analysis

Data were analyzed by another researcher who also received standardized training of SPSS for Mac (Version 21; IBM Corp, Armonk, NY, United States). Normally distributed and non-normally distributed continuous variables are presented as the mean \pm SD and median (range), respectively. Categorical variables are presented as counts (percentages). Normally distributed and non-normally distributed continuous variables were analyzed by independent-sample *t*-test and Mann-Whitney *U* test. Categorical variables were analyzed by the chi-square test. Statistical significance was defined at $P < 0.05$.

RESULTS

Patient characteristics

A total of 109 patients (110 knees) with knee valgus who underwent primary TKA were included in this study from January 2011 to December 2014. Six patients dropped out from the study, including two patients who died of a cardiovascular event, three patients lost to follow-up, and 1 bedridden patient with cerebral hemorrhage. Finally, 103 patients (103 knees), including 69 females and 34 males with a mean age of 63 years (range, 38 to 79 years), were included in this study with a mean follow-up period



DOI: 10.12998/wjcc.v10.i19.6406 Copyright ©The Author(s) 2022.

Figure 2 The new osteotomy technique, soft tissue balancing technique, and follow-up results of joint function. A: Schematic diagram of the distal femoral valgus osteotomy. Preoperative lower limb mechanical axis (MA) (a1, blackline), 3° (a2), 5° (a3), and 7° (a4) valgus osteotomy in distal femur osteotomy (white line: new MA; green line: anatomical axis of femur; yellow line: bone cutting plane) are shown, and 5° and 7° valgus cut angle can correct more osseous MA and increase the relative length of the lateral collateral ligament compared with 3° (a5); B: A 65-year-old man with 18° of valgus knee deformity in the new theory total knee arthroplasty (NT-TKA) group. The surgical procedure included osteophytes removal (b1), soft tissue release with a 20 mL syringe needle (b2), osteotomy (b3), straightening gap balance (b4), soft tissue cuff-like release (b5), and buckling gap balance (b6); C and D: Comparison of Oxford knee score (OKS), knee society score (KSS), and range of motion (ROM) between pre- and post-operation in conventional total knee arthroplasty (C-TKA) group (C) and NT-TKA group (D), respectively. ^bP < 0.01, compared with pre-OKS, KSS, and ROM in NT-TKA and pre-KSS clinical stability in C-TKA. ^aP < 0.05, compared with pre-OKS in C-TKA. C-TKA: Conventional total knee arthroplasty; NT-TKA: New theory total knee arthroplasty; OKS: Oxford knee score; KSS: Knee society score; ROM: Range of motion.

of 79 mo (range, 60 to 104 mo). Of the patients included, 42 (42 knees; 15 males and 27 females) with an average follow-up of 83 mo (range, 63-104 mo) were assigned to the C-TKA group, and 61 (61 knees; 19 males and 42 females) with an average follow-up period of 76 mo (range, 60-97 mo) were assigned to the NT-TKA group. There were no statistically significant differences between the two groups with regard to age, gender, body mass index, diagnosis, FTA, and length of follow-up (Table 1).

Comparison of postoperative outcomes between different groups

The KSS data in clinical stability at 1 mo post-operation for the C-TKA and NT-TKA groups were 11.2 ± 3.8 and 13.3 ± 2.9 , respectively, with a significant difference ($P < 0.01$); however, there was no significant difference in joint stability at the last follow-up (Table 2). The HKA, ROM, OKS KSS, and prosthesis survival rate data had no significant differences in both the preoperative and follow-up periods. In the C-TKA and NT-TKA groups, the last follow-up patient data showed improved HKA, ROM, OKS, and KSS compared to preoperative data (Figure 2C and D).

There were six patients with constrained prostheses, one with common peroneal nerve injury, and two with joint instability in the C-TKA group, but no patients developed such complications in the NT-TKA group. There were significant differences in the constrained prosthesis usage and complications between the groups ($P < 0.01$ and $P < 0.05$, respectively) (Table 2). At the last follow-up, the prostheses were all stable, and no patients underwent joint revision in either group.

DISCUSSION

This study proposed to apply the theory of synergistic balance between bone and soft tissue in TKA surgery, which reduced the need for soft tissue loosening, operation time, and constrained prosthesis while maintaining early stability of the joint.

TKA for VD requires the surgeon to be equipped with experience in dealing with different bone and soft-tissue conditions, positioning of the resection, soft-tissue release, and possible complications. For patients with severe VD, sacrificing 2° of neutral FTMA can abrogate excessive release of the lateral soft tissue and reduce the incidence of joint instability while maintaining joint function. To our knowledge, this is the first study to follow patients for more than 5 years in exchange for soft tissue stabilization at the expense of FTMA. In our opinion, TKA should be pursued with the common balance of bone and soft tissue since both are critical and closely related to preoperative joint function.

TKA for Ranawat type-II VD is challenging due to intra- and extra-articular bony abnormalities. Improper limb and component alignments occur due to distortion of the bony canal and variations in femoral anatomy. Therefore, a conventional system to guide alignment may be inappropriate[1,6]. The VCA can be determined in two ways during distal femur resection: Fixed VCA and individualized angle. Koskinen *et al*[7] suggested using a 3° VCA of the distal femur instead of a 5°-7° to achieve optimal correction of the underlying deformity for the knee valgus. However, Huang *et al*[6] found conventional 3° VCA in patients with Ranawat type-II VD knee might be unreasonable because of the coronal femoral bowing deformities. To restore the FTMA, distal femoral resection should be performed with an overcorrection of 2°[8]. Tucker *et al*[9] believed that a VCA of 5° valgus or more should be applied if the trapezoidal extension gap was unbalanced. However, fixed VCA might result in unexpected results. Some authors[10,11] advocated for an individualized angle, defined by the anatomical and mechanical axes of the femur. However, a hypoplastic lateral femoral condyle in the valgus knee might lead to the offsetting of the central knee joint line; hence, the actual anatomical axis of the femur is not achieved. In this study, the conventional 3° VCA was increased to 5°-7°, which corrected more osseous FTMA and increased the relative length of the lateral soft tissue. Moreover, 7° VCA is relatively safe in the knee valgus, like ordinary TKA, and does not damage the collateral ligament.

Balancing the soft tissue in Ranawat type-II valgus knee is another challenge and might contribute to complications such as peroneal nerve palsy and joint instability. Several surgical techniques have been developed to release the lateral supporting structures in TKA, including cruciform release, Z-plasty, and pie-crust technology. From our experience, the first step is to remove osteophytes, which affect soft tissue tension. We recommend to use the pie-crusting technique with a needle, not a blade, which can reduce the risk of joint instability.

Effective restoration of the FTMA mainly depends on the precise resection and soft tissue release in TKA. Poor prosthesis position will increase the failure rate of TKA[12]. A 8-12-year follow-up study of 115 TKA patients by Jeffrey *et al*[13] found that the rate of joint prosthesis loosening was about 3% with FTMA in a neutral position (within $\pm 3^\circ$) and 24% beyond this range ($> 3^\circ$) ($P < 0.05$). Studies have shown that a neutral MA can protect the soft tissue, improve joint function, and prolong prosthesis survival when adapted as a gold standard. However, Stucinskas *et al*[5] conducted a 1-year follow-up of 91 TKA patients and found that patients with neutral FTMA had no significant results in ROM, KSS, and muscle strength compared with patients with non-neutral FTMA. Bellemans *et al*[14] also held the same view that restoration of neutral FTMA in all patients may not be befitting and would be unnecessary. Although the mean coronal alignment was almost constant, most patients did not achieve

Table 1 Baseline characteristics of 103 cases in conventional total knee arthroplasty group and new theory total knee arthroplasty group

Characteristic	C-TKA group (n = 42 case)	NT-TKA group (n = 61 case)	P value
Knee number (n)	42	61	
Age (yr)	62.9 (39-77)	63.5 (38-79)	0.083
Gender, n (%)			0.628
Male	15 (35.71)	19 (31.15)	
Female	27 (64.29)	42 (68.85)	
BMI (kg/m ²)	24.0 ± 2.2	25.1 ± 2.8	0.091
Diagnosis, n (%)			0.770
Primary osteoarthritis	30 (71.43)	44 (72.13)	
Rheumatoid arthritis	11 (26.19)	14 (22.58)	
Hemophilic arthritis	1 (2.38)	3 (4.92)	
FTA (°)	26.8 ± 5.7	26.3 ± 5.2	0.784
Operation time	105.6 ± 17.7	99.2 ± 11.0	0.031
Follow-up (mo)	83.4 ± 10.1	75.6 ± 10.0	0.907

Continuous data are shown as the mean ± SD or median (range), and categorical data are presented as n (%). BMI: Body mass index; FTA: Femorotibial axis; C-TKA: Conventional total knee arthroplasty; NT-TKA: New theory total knee arthroplasty.

neutral FTMA[14]. Hirschmann *et al*[4] measured the HKA angle of 308 non-osteoarthritic knees of 160 patients and found that constitutional varus and valgus probabilities were 32.4% and 26.6%, respectively. Similarly, Bellemans *et al*[14] found an incidence of 48.2% and 4.8%, respectively. Lately, the concept of anatomic restoration or kinematically aligned (KA) has gained traction[15]. This theory does not necessarily restore neutral FTMA but rather the natural alignment of the knee. KA TKA reduces the need for soft tissue release and naturally restores the FTMA more to the pre-arthritis state of the knee. Laende *et al*[16] conducted a randomized controlled trial with a 2-year follow-up and found no significant differences in the prosthesis fixation and OKS scores between KA TKA and FTMA TKA. A meta-analysis by Hetaimish *et al*[17] showed that even if KA TKA did not deliberately pursue the neutral MA, there was no significant difference in the incidence of postoperative complications, hemoglobin level, or hospital stay compared with FTMA TKA. However, there was a benefit trend towards shorter operation time ($P = 0.01$) and better overall knee function ($P < 0.01$) in patients with FTMA TKA. Besides, more patients who undergo FTMA TKA might have an outlier varus and valgus alignment when compared to FTMA due to high intraoperative forces in the medial and lateral compartments of the knee[18]. Based on our theory, which was similar to KA, patients with poor preoperative lateral soft tissue status and $VD > 20^\circ$ would be sacrificed a 2° valgus.

The constrained prosthesis was considered a better option when the above strategy did not achieve joint stability. The constrained prosthesis can simplify operation procedures, avoid excessive soft tissue release, and shorten operation time but not without its associated prohibitive cost, more bone loss, and knee pain. Meanwhile, the rate of constrained prosthesis revision was twice as high at 10 years and three times higher at 20 years postoperatively[19]. Therefore, several studies recommend minimizing the use of constrained prostheses under the premise of stability[20]. The balance between bone and soft tissue should be considered during any TKA procedure. If joint space balance cannot be achieved by soft tissue release, the VCA should be increased appropriately. If the joint gap is still unbalanced, a 2° FTMA is sacrificed for the joint stability and unconstrained prosthesis (Table 2). Statistical results show that the probability of peroneal nerve injury is 0.3%-9.5%, which manifests as sagging feet, spanning gait, inability to supinate the foot, muscle atrophy, and loss of cutaneous sensation in the innervated area. In the current study, there were 1 of 42 patients with peroneal nerve injury caused by traction, which recovered fully following VD correction in the C-TKA group but no patient in the NT-TKA cohort developed this injury.

Limitations of this study include possible biases like patient performance bias and lack of independent review of radiographs. In addition, this was a single surgeon case series using the new concept of resection and soft tissue release. Also, our study is a short-term study, and the theory of bone and soft tissue co-balance requires longer follow-up to ascertain its impact on joint function and prosthetic survival. The application of finite element analysis can provide simulated solutions for such clinical conundrums. Future research should focus on the influence of resection on soft tissue release and the critical value of ligament release by finite element analysis, along with mechanical changes in

Table 2 Comparison of surgery-related results between conventional total knee arthroplasty group and new theory total knee arthroplasty group

Parameter	C-TKA group (n = 42 case)	NT-TKA group (n = 61 case)	P value
HKA (°)			
Pre-	17.2 ± 5.9	16.0 ± 4.5	0.076
Post-	0.5 ± 1.0	0.8 ± 1.1	0.225
P value	< 0.001	<0.001	-
ROM (°)			
Pre-	73.0 ± 18.7	75.0 ± 20.1	0.369
Post-	93.5 ± 13.6	102.9 ± 11.8	0.781
P value	0.074	< 0.001	-
OKS			
Pre-	42.0 ± 4.5	42.8 ± 4.5	0.654
Post-	17.4 ± 3.3	17.8 ± 3.6	0.854
P value	0.038	0.173	-
KSS			
Pre-clinical	14.9 ± 12.1	13.9 ± 9.7	0.077
Post-clinical	78.9 ± 6.4	81.3 ± 5.0	0.655
P value	0.010	< 0.001	-
Pre-functional	34.2 ± 19.4	25.7 ± 20.9	0.530
Post-functional	69.9 ± 17.1	60.0 ± 12.1	0.172
P value	0.117	< 0.001	-
KSS clinical stability			
Preoperative	9.3 ± 4.4	10.0 ± 4.0	0.378
1 mo (post)	11.2 ± 3.8	13.3 ± 2.9	0.007
Last follow-up	14.3 ± 1.8	14.5 ± 1.5	0.132
Complications	3	0	0.034
Instability	2	0	
Nerve injury	1	0	
Prosthesis			0.002
Posterior stabilized	36	61	
Constrained prosthesis	6	0	
Survival rate	42/42	61/61	

Continuous data are shown as the mean ± SD. HKA: Hip-knee-ankle; ROM: Range of motion; OKS: Oxford knee score; KSS: Knee society score; C-TKA: Conventional total knee arthroplasty; NT-TKA: New theory total knee arthroplasty.

the medial and lateral prosthesis planes post-VD correction.

CONCLUSION

Bone and soft tissue balance is equally important in a TKA procedure. Adopting 5°-7° VCA for VD and sacrificing 2° neutral FTMA for severe VD of the knee during TKA can reduce the need for soft tissue release, maintain joint stability, and abrogate the use of constrained prostheses while minimizing postoperative complications.

ARTICLE HIGHLIGHTS

Research background

Total knee arthroplasty (TKA) for Ranawat Type II valgus deformity (VD) of the knee is challenging and technical due to resection and soft tissue balance.

Research motivation

Unreasonable osteotomy and soft tissue release in pursuit of neutral femorotibial mechanical axis (FTMA) will increase joint instability, limited prosthesis use, and complications.

Research objectives

This study aimed to provide a new technique of surgical resection, soft tissue release, and FTMA for the VD of the knee during TKA.

Research methods

Sixty-one patients had a valgus cut angle (VCA) of 5°-7° in the new theory TKA group (NT-TKA). We chose to sacrifice the 2° FTMA to reduce the risk of survival failure, joint installation, and consumption of constrained procedures if the FTMA still cannot return to the neutral position after the maximum soft tissue release. Forty-two patients in the conventional TKA group (C-TKA) were treated with a VCA of 3°-5° and pursuit of neutral FTMA.

Research results

The constrained prosthesis usage and complications in NT-TKA were lower than those in C-TKA ($P = 0.002$ and $P = 0.034$, respectively). The KSS at 1 mo post-operation for NT-TKA was higher than that in C-TKA ($P = 0.007$).

Research conclusions

Adopting 5°-7° VCA for VD and sacrificing 2° neutral FTMA for severe VD which cannot be completely corrected during TKA can reduce the need for soft tissue release, maintain early joint stability, reduce the use of constrained prostheses, and minimize postoperative complications.

Research perspectives

This series of surgical technique for VD of the knee during TKA have certain clinical guiding significance.

ACKNOWLEDGEMENTS

We would like to thank Tong PJ and Lv SJ for their help in research ideas and design; He BJ and Lv SJ for their role in research implementation and manuscript writing; Huang JF for manuscript editing and writing; Wang XJ, Mao Q, Lv SJ, and He BJ for their immense support in data collection and manuscript submission.

FOOTNOTES

Author contributions: Tong PJ and Lv SJ conceived and coordinated the study, designed, performed, and analyzed the experiments, and wrote the paper; He BJ and Lv SJ carried out the data collection and analysis, and revised the paper; all authors reviewed the results and approved the final version of the manuscript.

Supported by the Project of Excellent Young Talents of Traditional Chinese Medicine of Zhejiang Province, No. 2019ZQ016; and the Zhejiang Medical and Health Science and Technology Young Talents Program, No. 2019RC059.

Institutional review board statement: This study was approved by the hospital ethics and review committee of the First Affiliated Hospital of Zhejiang Chinese Medical University (No. 2019-K-207-01).

Informed consent statement: All patients provided informed consent for this study.

Conflict-of-interest statement: We have no financial relationships to disclose.

Data sharing statement: No additional data are available.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-

NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

Country/Territory of origin: China

ORCID number: Shuai-Jie Lv 0000-0003-1755-6665; Xiao-Jian Wang 0000-0002-2459-7388; Jie-Feng Huang 0000-0002-1982-6641; Qiang Mao 0000-0002-9382-6527; Bang-Jian He 0000-0002-7853-813X; Pei-Jian Tong 0000-0003-0948-8674.

S-Editor: Yan JP

L-Editor: Wang TQ

P-Editor: Yan JP

REFERENCES

- Ranawat AS, Ranawat CS, Elkus M, Rasquinha VJ, Rossi R, Babhulkar S. Total knee arthroplasty for severe valgus deformity. *J Bone Joint Surg Am* 2005; **87** Suppl 1: 271-284 [PMID: 16140800 DOI: 10.2106/JBJS.E.00308]
- Rossi R, Rosso F, Cottino U, Dettoni F, Bonasia DE, Bruzzone M. Total knee arthroplasty in the valgus knee. *Int Orthop* 2014; **38**: 273-283 [PMID: 24366186 DOI: 10.1007/s00264-013-2227-4]
- McAuley JP, Collier MB, Hamilton WG, Tabaraee E, Engh GA. Posterior cruciate-retaining total knee arthroplasty for valgus osteoarthritis. *Clin Orthop Relat Res* 2008; **466**: 2644-2649 [PMID: 18712454 DOI: 10.1007/s11999-008-0436-8]
- Moser LB, Hess S, Amsler F, Behrend H, Hirschmann MT. Native non-osteoarthritic knees have a highly variable coronal alignment: a systematic review. *Knee Surg Sports Traumatol Arthrosc* 2019; **27**: 1359-1367 [PMID: 30848310 DOI: 10.1007/s00167-019-05417-2]
- Stucinskas J, Robertsson O, Sirka A, Lebedev A, Wingstrand H, Tarasevicius S. Moderate varus/valgus malalignment after total knee arthroplasty has little effect on knee function or muscle strength. *Acta Orthop* 2015; **86**: 728-733 [PMID: 26058850 DOI: 10.3109/17453674.2015.1059689]
- Huang TW, Chuang PY, Lee CY, Lin SJ, Huang KC, Shen SH, Tsai YH, Lee MS, Hsu RW. Total knee arthroplasty in patients with Ranawat type-II valgus arthritic knee with a marked coronal femoral bowing deformity: comparison between computer-assisted surgery and intra-articular resection. *J Orthop Surg Res* 2016; **11**: 88 [PMID: 27488841 DOI: 10.1186/s13018-016-0422-x]
- Koskinen E, Remes V, Paavolainen P, Harilainen A, Sandelin J, Tallroth K, Kettunen J, Ylinen P. Results of total knee replacement with a cruciate-retaining model for severe valgus deformity--a study of 48 patients followed for an average of 9 years. *Knee* 2011; **18**: 145-150 [PMID: 20471274 DOI: 10.1016/j.knee.2010.04.001]
- Lombardi AV Jr, Dodds KL, Berend KR, Mallory TH, Adams JB. An algorithmic approach to total knee arthroplasty in the valgus knee. *J Bone Joint Surg Am* 2004; **86-A** Suppl 2: 62-71 [PMID: 15691110 DOI: 10.2106/00004623-200412002-00010]
- Tucker A, O'Brien S, Doran E, Gallagher N, Beverland DE. Total Knee Arthroplasty in Severe Valgus Deformity Using a Modified Technique-A 10-Year Follow-Up Study. *J Arthroplasty* 2019; **34**: 40-46.e1 [PMID: 30318254 DOI: 10.1016/j.arth.2018.09.002]
- Shi X, Li H, Zhou Z, Shen B, Yang J, Pei F. Comparison of Postoperative Alignment Using Fixed vs Individual Valgus Correction Angle in Primary Total Knee Arthroplasty With Lateral Bowing Femur. *J Arthroplasty* 2016; **31**: 976-983 [PMID: 26787012 DOI: 10.1016/j.arth.2015.10.040]
- Rahm S, Camenzind RS, Hingsammer A, Lenz C, Bauer DE, Farshad M, Fucentese SF. Postoperative alignment of TKA in patients with severe preoperative varus or valgus deformity: is there a difference between surgical techniques? *BMC Musculoskelet Disord* 2017; **18**: 272 [PMID: 28637433 DOI: 10.1186/s12891-017-1628-8]
- Kim YH, Park JW, Kim JS, Park SD. The relationship between the survival of total knee arthroplasty and postoperative coronal, sagittal and rotational alignment of knee prosthesis. *Int Orthop* 2014; **38**: 379-385 [PMID: 24173677 DOI: 10.1007/s00264-013-2097-9]
- Jeffery RS, Morris RW, Denham RA. Coronal alignment after total knee replacement. *J Bone Joint Surg Br* 1991; **73**: 709-714 [PMID: 1894655 DOI: 10.1302/0301-620X.73B5.1894655]
- Bellemans J, Colyn W, Vandenneucker H, Victor J. The Chitranjan Ranawat award: is neutral mechanical alignment normal for all patients? *Clin Orthop Relat Res* 2012; **470**: 45-53 [PMID: 21656315 DOI: 10.1007/s11999-011-1936-5]
- Oussedik S, Abdel MP, Victor J, Pagnano MW, Haddad FS. Alignment in total knee arthroplasty. *Bone Joint J* 2020; **102-B**: 276-279 [PMID: 32114811 DOI: 10.1302/0301-620X.102B3.BJJ-2019-1729]
- Laende EK, Richardson CG, Dunbar MJ. A randomized controlled trial of tibial component migration with kinematic alignment using patient-specific instrumentation versus mechanical alignment using computer-assisted surgery in total knee arthroplasty. *Bone Joint J* 2019; **101-B**: 929-940 [PMID: 31362561 DOI: 10.1302/0301-620X.101B8.BJJ-2018-0755.R3]
- Hetaimish BM, Khan MM, Simunovic N, Al-Harbi HH, Bhandari M, Zalzal PK. Meta-analysis of navigation vs conventional total knee arthroplasty. *J Arthroplasty* 2012; **27**: 1177-1182 [PMID: 22333865 DOI: 10.1016/j.arth.2011.12.028]
- Shelton TJ, Nedopil AJ, Howell SM, Hull ML. Do varus or valgus outliers have higher forces in the medial or lateral compartments than those which are in-range after a kinematically aligned total knee arthroplasty? *Bone Joint J* 2017; **99-B**: 1319-1328 [PMID: 28963153 DOI: 10.1302/0301-620X.99B10.BJJ-2017-0066.R1]
- Martin JR, Beahrs TR, Stuhlman CR, Trousdale RT. Complex Primary Total Knee Arthroplasty: Long-Term Outcomes. *J Bone Joint Surg Am* 2016; **98**: 1459-1470 [PMID: 27605690 DOI: 10.2106/JBJS.15.01173]

- 20 **Madonna V**, Screpis D, Condello V, Piovan G, Russo A, Guerriero M, Zorzi C. A novel technique for combined medial collateral ligament and posterior oblique ligament reconstruction: technical note. *Knee Surg Sports Traumatol Arthrosc* 2015; **23**: 2814-2819 [PMID: [26233595](#) DOI: [10.1007/s00167-015-3721-6](#)]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA
Telephone: +1-925-3991568
E-mail: bpgoffice@wjgnet.com
Help Desk: <https://www.f6publishing.com/helpdesk>
<https://www.wjgnet.com>

