



Coexistence of diffuse large B-cell lymphoma, acute myeloid leukemia, and untreated lymphoplasmacytic lymphoma/waldenström macroglobulinemia in a same patient: A case report

Liu-Bo Zhang, Lu Zhang, Hong-Lei Xin, Yan Wang, Hong-Yu Bao, Qing-Qi Meng, Su-Yu Jiang, Xue Han, Wan-Ru Chen, Jian-Ning Wang, Xiao-Feng Shi

Specialty type: Hematology

Provenance and peer review:

Unsolicited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's scientific quality classification

Grade A (Excellent): 0

Grade B (Very good): 0

Grade C (Good): C

Grade D (Fair): 0

Grade E (Poor): 0

P-Reviewer: Saito M, Japan

Received: January 4, 2023

Peer-review started: January 4, 2023

First decision: April 19, 2023

Revised: April 25, 2023

Accepted: May 9, 2023

Article in press: May 9, 2023

Published online: June 26, 2023



Liu-Bo Zhang, Lu Zhang, Hong-Lei Xin, Hong-Yu Bao, Qing-Qi Meng, Su-Yu Jiang, Xue Han, Wan-Ru Chen, Jian-Ning Wang, Xiao-Feng Shi, Department of Hematology, Second Affiliated Hospital of Nanjing Medical University, Nanjing 210003, Jiangsu Province, China

Yan Wang, Department of Pathology, Second Affiliated Hospital of Nanjing Medical University, Nanjing 210003, Jiangsu Province, China

Corresponding author: Xiao-Feng Shi, MD, PhD, Chief Physician, Director, Department of Hematology, Second Affiliated Hospital of Nanjing Medical University, No. 262 North Zhongshan Road, Nanjing 210003, Jiangsu Province, China. shixiaofeng1977@163.com

Abstract

BACKGROUND

The Coexistence of myeloid and lymphoid malignancies is rare. Myeloid leukemia occurs more frequently as a secondary event in patients receiving chemotherapy agents for lymphoid malignancies. Synchronous diagnoses of diffuse large B-cell lymphoma (DLBCL), acute myeloid leukemia (AML), and untreated lymphoplasmacytic lymphoma/Waldenström macroglobulinemia (LPL/WM) in the same patient have not been reported. Here we report one such case.

CASE SUMMARY

An 89-year-old man had a chest wall mass histopathologically diagnosed as DLBCL. The bone marrow and peripheral blood contained two groups of cells. One group of cells fulfilled the criteria of AML, and the other revealed the features of small B lymphocytic proliferative disorder, which we considered LPL/WM. Multiple chromosomal or genetic changes were detected in bone marrow mononuclear cells, including *ATM* deletion, *CCND1* amplification, mutations of *MYD88* (L265P) and *TP53*, *WT1* overexpression, and fusion gene of *BIRC2-ARAP1*, as well as complex chromosomal abnormalities. The patient refused chemotherapy because of old age and died of pneumonia 1 mo after the final diagnosis.

CONCLUSION

The coexistence of DLBCL, AML, and untreated LPL/WM in the same patient is extremely rare, which probably results from multiple steps of genetic abnor-

malities. Asymptomatic LPL/WM might have occurred first, then myelodysplastic syndrome-related AML developed, and finally aggressive DLBCL arose. Therefore, medical staff should pay attention to this rare phenomenon to avoid misdiagnoses.

Key Words: Diffuse large B-cell lymphoma; Acute myeloid leukemia; Small B lymphocyte proliferative disorder; Lymphoplasmacytic lymphoma/Waldenström macroglobulinemia; Coexistence; Case report

©The Author(s) 2023. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: The coexistence of myeloid and lymphoid malignancies is rare. Synchronous diagnosis of diffuse large B-cell lymphoma (DLBCL), acute myeloid leukemia (AML), and untreated lymphoplasmacytic lymphoma/Waldenström macroglobulinemia (LPL/WM) in the same patient has not yet been reported. Herein, we report one such case. We assumed that the three diseases have been occurred in a stepwise manner. Asymptomatic indolent LPL/WM might have occurred earlier, then insidious myelodysplastic syndrome developed into AML, while chest wall DLBCL grew separately.

Citation: Zhang LB, Zhang L, Xin HL, Wang Y, Bao HY, Meng QQ, Jiang SY, Han X, Chen WR, Wang JN, Shi XF. Coexistence of diffuse large B-cell lymphoma, acute myeloid leukemia, and untreated lymphoplasmacytic lymphoma/waldenström macroglobulinemia in a same patient: A case report. *World J Clin Cases* 2023; 11(18): 4295-4305

URL: <https://www.wjgnet.com/2307-8960/full/v11/i18/4295.htm>

DOI: <https://dx.doi.org/10.12998/wjcc.v11.i18.4295>

INTRODUCTION

Diffuse large B-cell lymphoma (DLBCL) and acute myeloid leukemia (AML) rarely occur concurrently. Only a few cases of DLBCL after AML chemotherapy[1,2] have been reported. Chemotherapy-induced immunosuppression predisposes patients to develop secondary malignancies, particularly virus-related ones[1,2]. Only two cases with a synchronous diagnoses of DLBCL and AML have been reported[3,4]. The coexistence of AML and small B lymphocytic proliferative disorder (LPD) is also rare. Most cases of AML develop after small B LPD (B-LPD), probably due to immunodeficiency over the long course of LPD[5,6] or after treatment[7]. Only a few cases of newly diagnosed synchronous AML and small B-LPD have been reported[8-10]. Lymphoplasmacytic lymphoma/Waldenström macroglobulinemia (LPL/WM) is a specific type of B-LPD. To the best of our knowledge, synchronous diagnosis of DLBCL, AML, and untreated LPL/WM in a same patient has not been reported yet. Herein, we report one such case.

CASE PRESENTATION

Chief complaints

An 89-year-old man was admitted to our hospital complaining of cough, expectoration, and shortness of breath for half a month.

History of present illness

The patient complained of cough and expectoration for half a month. His blood test results were abnormal: Leukocyte count, 12.6 ($4.0-10.0 \times 10^9/L$); neutrophil count, 1.44 ($2.00-7.50 \times 10^9/L$); lymphocyte count, 4.21 ($0.80-4.00 \times 10^9/L$); monocyte count, 6.9 ($0-0.8 \times 10^9/L$); hemoglobin level, 73 ($120-160g/L$); and platelet count 55 ($100-300 \times 10^9/L$). The patient was therefore admitted to the Department of Hematology.

History of past illness

The patient had chronic bronchitis for 10 years, which often deteriorated in winter.

Physical examination

Physical examination revealed skin pallor, enlarged lymph nodes in the armpit, and moist rales in both lungs, but no hepatosplenomegaly.

Laboratory examinations

CT-guided puncture of the mass in the chest wall was performed for biopsy. The histopathological examination showed a hyperproliferation of large cells with Ki-67 (70%), CD20 (+++), PAX-5 (++), CD10 (+), MUM-1 (+), BCL-6 (+), c-MYC (< 40%), BCL-2 (< 50%), CD3 (-), CK-pan (-), TTF (-), NapsinA (-), CK7 (-), P40 (-), CK5/6 (-), Syn (-), CgA (-), CD56 (-), and EBER (-), suggesting a diagnosis of DLBCL (Figure 1A-O, Table 1). Blood tests were performed, and many abnormalities were found: Procalcitonin levels, 1.42 (< 0.05 ng/mL); erythrocyte sedimentation rate, 97 (0-20 mm/h); C-reactive protein levels, 370 (0-10 mg/L); lactate dehydrogenase levels, 460 (120-250 U/L); serum albumin levels, 34.3 (40.0-55.0 g/L); serum fibrinogen levels, 7.24 (2.00-4.00 g/L); CA 125, 48.2 (0-35.0 U/mL); pro-B-type natriuretic peptide, 1918 (< 300 pg/mL); serum immunoglobulin (Ig) M levels, 6.55 (0.40-2.30 g/L); urine Kappa light chain, 318 (0-50 mg/L); free triiodothyronine levels, 2.61 (3.10-6.80 pmol/L); and triiodothyronine levels, 0.77 (1.30-3.10 nmol/L). Peripheral blood and bone marrow smears were performed. Surprisingly, no large B lymphocytes were found, but there were 71.6% mature small lymphocytes, along with 23.2% myeloid blasts which were peroxidase (++), periodic acid-Schiff stain (+), and non-specific esterase (-/+), cannot be inhibited by NaF (Figure 2A-F, Table 1). The patient's peripheral blood had a high percentage of myeloblasts (50%) and a relatively low percentage of B lymphocytes (29%) (Figure 2A-F). Immunohistochemical analysis of the bone marrow biopsy showed mixed expression of myeloblast and mature B lymphocyte markers with CD10 (+), CD20 (+++), CD34 (++), CD43 (+), CD117 (++), PAX-5(+++), myeloperoxidase (MPO) (+), silver staining (++), BCL-6 (-), CD3 (-), and CD5 (-) (Figure 2G-R, Table 1). To identify the features of these mixed cells, flow cytometry was used to gate the two groups of cells from peripheral blood and bone marrow samples. The immunophenotypic results confirmed the morphological findings of the presence of two discrete abnormal cell populations: 49.64% large blasts with dim CD45 expression and 22.21% small lymphocytes with strong CD45 expression. The large blasts population expressed CD34, CD13, and CD117, and partially expressed CD33, HLA-DR, and MPO, but did not express CD3, CD4, CD5, CD7, CD8, CD10, CD11b, CD14, CD15, CD19, CD20, CD36, CD56, CD64, or CD23, compatible with myeloblasts. In contrast, the small lymphocytes expressed CD19, CD20, and Kappa light chain, but did not express CD103, CD25, CD11c, CD34, CD117, CD5, CD10, CD23 or Lambda light chain, which were considered LPD cells (Figure 3, Table 1). Therefore, AML-M2 along with a small B-LPD were suspected. Furthermore, cytogenetic and molecular biology tests were performed on bone marrow mononuclear cells (BMMNCs). Monoclonal IgM Kappa was identified by immunofixation electrophoresis (Figure 4A), which was consistent with the diagnosis of monoclonal B lymphocytosis or plasmacytosis. Fluorescence *in situ* hybridization (FISH) results showed deletion of 11q22 (*ATM*) (83%) and an increased copy number of 11q13 (*CCND1*) (80%) (Figure 4B and C), but no abnormalities of chromosomes or genes in 13q34, 13q14.3, 12, 17p13.1 (*P53*), or 13q14.2 (*RB1*). The IgH rearrangement was confirmed and t(11; 14) was not detected. Chromosomal karyotype analysis was performed and five metaphases were obtained using the routine method, whereas 20 metaphases were obtained after stimulation with CpG-oligodeoxynucleotides. All metaphases showed a complex abnormality with 46, XY, -2, del(5)(q13q31), -8, add(11)(q13), del(13)(q11q32), +mar1, and +mar2 (Figure 4D). A total of 281 genes associated with malignant lymphoid and myeloid diseases were tested by sequencing (Supplementary material). *MYD88* (14.8%) mutation (p.L265P) and 27.3% of *TP53* mutations (p.N29Kfs*14 and p.S215N) were detected (Table 2). Different kinds of copy number variations were found in chromosome 2p25.3p13.3 (2.56 copies), 5q14.3q34 (1.46 copies), 8p23.3p21.2 (4.30 copies), 8q21.3q24.3 (2.55 copies), 11q12.1q14.1 (3.05 copies), 11q14.1q22.1 (1.43 copies), 11q22.3q23.1 (1.44 copies), 11q23.1q25 (4.85 copies), 13q12.3q14.13 (1.43 copies), 13q21.1q22.1 (2.63 copies), and 13q31.1q31.3 (2.49 copies), which were consistent with the karyotype results (Table 2). AML-associated 56 fusion genes were tested using real-time PCR (Supplementary material) and *WT1* overexpression was also detected (Table 2). Newly discovered fusion genes were also tested using RNA sequencing (Supplementary material), and a new fusion gene, *BIRC2(11q22.2)-ARAP1(11q13.4)*, was detected (Table 2). Gene sequencing of the tissue from the chest wall mass was not available because of the inability to obtain sufficient tissue.

Imaging examinations

Chest and abdominal CT revealed multiple shadows in both lungs, a newborn mass in the left chest wall, and multiple enlarged lymph nodes in the armpits, mediastinum, and hilus pulmonis (Figure 1P-T).

FINAL DIAGNOSIS

The patient was diagnosed with the coexistence of DLBCL, AML, and LPL/WM.

Table 1 Immunophenotype of bone marrow and chest wall tumor

Expression	Molecular marker
The immunophenotype of BMMNCs by flow cytometry	
Group A	49.64%
Expression	CD34, CD13, CD117
Partial expression	CD33, HLA-DR
No expression	CD3, CD4, CD5, CD7, CD8, CD10, CD11b, CD14, CD15, CD19, CD20, CD36, CD56, CD64, CD23
Group B	22.21%
Expression	CD19, CD20, Kappa
Partial expression	
No expression	CD103, CD25, CD11c, CD34, CD117, CD5, CD10, CD23, lambda
The immunohistochemical analysis of bone marrow biopsy	
Expression	CD10, CD20, CD34, CD43, CD117, PAX-5, MPO, silver dyeing
Partial expression	
No expression	BCL-6, CD3, CD5, MPO
The immunohistochemical analysis of chest wall tumor	
Expression	Ki-67 (70%), CD20, PAX-5, CD10, MUM-1, BCL-6, c-MYC (< 40%), BCL-2 (< 50%)
Partial expression	
No expression	CD3, CK-pan, TTF, NapsinA, CK7, P40, CK5/6, Syn, CgA, CD56, EBER

BMMNCs: Bone marrow mononuclear cells; MPO: Myeloperoxidase.

TREATMENT

The patient refused chemotherapy because of old age and performance status.

OUTCOME AND FOLLOW-UP

The patient died of pneumonia 1 mo later. The patient’s family refused an autopsy.

DISCUSSION

This patient was admitted to our hospital because of the chest wall mass, which was histopathologically identified as DLBCL (GCB subtype). The cells were large and strongly positive for CD20, PAX-5, BCL-6, and MUM-1, with high Ki-67expression, suggesting an aggressive feature. The positive expression of BCL-2, BCL-6, and c-MYC suggested that DLBCL was a “triple expressor” type. MPO-negative expression of these large cells ruled out the probability of myeloblast infiltration. Blood cell abnormalities were occasionally observed during regular examinations. The two groups of abnormal cells were found in peripheral blood and bone marrow. Morphology and immunohistochemistry suggested a diagnosis of AML and small B-LPD. The two groups of cells were identified by flow cytometry. The cells expressing CD13, CD34, and CD117 were from myeloid blasts, whereas the others expressing CD19, CD20, and Kappa were from B lymphoid mature cells. Based on the morphology and flow cytometry results, the B-cells were found to be small, consistent with a diagnosis of small B-LPD, which is a chronic monoclonal neoplasm, including chronic lymphocytic leukemia/small lymphocytic lymphoma (CLL/SLL), mantle cell lymphoma (MCL), follicular lymphoma (FL), marginal zone lymphoma (MZL), hairy cell leukemia (HCL), and LPL/WM. Negative expression of CD103, CD25, and CD11b ruled out the diagnosis of HCL or HCLv. The panel of CD5 (-), CD23 (-), and CD10 (-) ruled out the diagnosis of CLL/SLL and FL[11]. Additionally, the lack of t (11;14) ruled out the diagnosis of MCL [11]. Subsequently, LPL/WM and MZL were considered. The patient had monoclonal IgM Kappa and MYD88 L265P mutations, and LPL/WM was diagnosed, whereas, MZL with plasmacytoid differentiation could not be completely ruled out. Both LPL/WM and MZL seldom transform to DLBCL,

Table 2 Abnormalities of genes from bone marrow mononuclear cells


Expression	Abnormality
The mutation of genes	
<i>MYD88</i>	p.L265P (14.8%)
<i>TP53</i>	p.N29Kfs*14 (27.3%)
<i>TP53</i>	p.S215N (27.2%)
The fusion of genes or gene expression	
<i>WT1</i>	Positive
<i>BIRC2-ARAP1</i>	Positive
The copy number variations	
chr2p25.3p13.3	2.56
chr5q14.3q34	1.46
chr8p23.3p21.2	4.30
chr8q21.3q24.3	2.55
chr11q12.1q14.1	3.05
chr11q14.1q22.1	1.43
chr11q22.3q23.1	1.44
chr11q23.1q25	4.85
chr13q12.3q14.13	1.43
chr13q21.1q22.1	2.63
chr13q31.1q31.3	2.49

although MZL often infiltrates extranodal tissues, including the lungs. Whether LPL/WM and DLBCL are from the same clone is still unknown since the genes from chest wall masses were not tested because of the limited tissue obtained through CT-guided puncture. Therefore, we considered that in the bone marrow LPL/WM occurred and the chest wall tumor was DLBCL.

Multiple chromosomal and gene changes were detected in the BMMNCs (Guangzhou KingMed Diagnostics Group Co., Ltd.). The routine karyotype analysis method, by which more metaphases were from myeloblasts, or CPG-oligodeoxynucleotide stimulating method, by which metaphases were from lymphocytes, detected the homogeneous chromosomal changes of 46, XY, -2, del(5)(q13q31), -8, add(11)(q13), del(13)(q11q32), +mar1, and +mar2. Myeloblasts (49.64%) and lymphocytes (22.21%) shared the same chromosomal abnormalities. Moreover, FISH detected 83% of 11q22 deletions, and 80% of 11q13 multiple copies. Based on this homogeneity, we suspected that myeloblasts and lymphocytes might be derived from the same progenitor, although different clones were not separated for molecular analysis due to the limited specimens.


Often, 5q- is found in AML or myelodysplastic syndrome (MDS), 11q- in small B-LPD, and 13q- in MDS, AML, or CLL. Although the copy number of 11q13 (*CCND1*) increases, no chromosomal change of t (11;14) is found; therefore, MCL should be ruled out. The deletion of 11q22.3 (*ATM*) often causes a defect in apoptosis, similar to deletion or mutation of 17p13 (*TP53*), suggesting a poor prognosis[12,13]. The frameshift mutation of *TP53* (p.N29Kfs*14) leads to a change in amino acid sequence or early termination of protein translation, causing a loss of *TP53* function. Missense mutation of *TP53* (p.S215N) can be found in lymphoid or myeloid malignant diseases, such as CLL, DLBCL, MCL, FL, LPL/WM, AML, and MDS, and are associated with poor prognosis. *WT1* is overexpressed in newly diagnosed or relapsed AML, CML in the accelerated and blastic phases, and high-risk MDS. *BIRC2(11q22.2)-ARAP1(11q13.4)* is a novel fusion gene discovered in this patient. Whether it is related with AML or LPD remains unknown. The patient's complex chromosomal abnormality with 5q- and old age led us to speculate that AML may originate from MDS, although dysplasia in erythroblasts or megakaryocytes was not found.

Multiple theories behind the simultaneous development of several malignancies in an untreated patient have been proposed, including immunosuppression mediated by chronic small B-LPDs[14], a common stem cell defect[15], or just a chance. Therefore, we hypothesized that these diseases occurred in a stepwise manner. Asymptomatic indolent LPL/WM might have occurred earlier, then insidious MDS developed into AML, while chest wall DLBCL grew separately.




DOI: 10.12998/wjcc.v11.i18.4295 Copyright ©The Author(s) 2023.


Figure 1 The biopsy and imaging of chest wall mass. A-O: The HE and immunohistochemistry of chest wall mass; A: Hematoxylin-Eosin (HE) staining $\times 20$ objective; B: HE staining $\times 40$ objective; C-O: The immunohistochemistry shows Ki-67 (++, approximately 70%), CD20 (++), PAX-5 (++), CD10 (+), MUM-1 (++), BCL-6 (++), c-MYC (+, < 40%), BCL-2 (+, < 50%), CD3 (T cells few +), CK-pan (-), myeloperoxidase (MPO) (-), CD 56 (-), and EBER (-). P: Enlarged axillary lymph node (arrow) in the chest computed tomography (CT) scan; Q and R: Infiltration of masses (arrows) to the lung and chest wall on mediastinum window (Q) and lung window (R). S and T: Unenhanced (S) and contrast-enhanced (T) CT images shows the infiltration of masses (arrows) to the lung and chest wall on mediastinal window.



DOI: 10.12998/wjcc.v11.i18.4295 Copyright ©The Author(s) 2023.

Figure 2 The cytomorphology of peripheral blood and bone marrow. A: Peripheral blood smear (Wright staining, $100 \times$ objective); B: Bone marrow smear (Wright staining, $100 \times$ objective); C-F: Histochemistry staining for bone marrow smear. Peroxidase (52%, 99 points), periodic acid-Schiff stain (86%, 158 points), non-specific esterase (NSE) (32%, 35 points), and NSE + NaF (25%, 26 points) staining for bone marrow smear (Blue arrows direct myeloid blasts, while green arrows direct small lymphocytes); G-R: Bone marrow biopsy ($40 \times$ objective); G: Hematoxylin-Eosin staining; H-R: Immunohistochemistry shows myeloperoxidase (myelocytes +), CD34 (myelocytes ++), CD117 (myelocytes +), CD20 (B cells ++), PAX-5 (B cells ++), CD3 (T cells, few +), CD10 (few +), BCL-6(-), CD43(partial +), CD5 (T cells -), and silver staining (+++). POX: Peroxidase; PAS: Periodic acid-Schiff; NSE: Non-specific esterase; MPO: Myeloperoxidase.






DOI: 10.12998/wjcc.v11.i18.4295 Copyright ©The Author(s) 2023.

Figure 3 Immunophenotype of bone marrow mononuclear cells. Cells of group A (red color) are myeloid blasts expressing CD34, CD13, and CD117, and partially expressing CD33, HLA-DR, and myeloperoxidase (MPO). Cells of group B (blue color) are clonal B-lymphocytes expressing CD19, CD20, and Kappa,

but no expressing CD103, CD25, CD11c, CD34, CD117, CD5, CD10, CD23, and Lambda. The results out the red box were from bone marrow sample obtained by bone marrow aspiration. The results in the red box were from peripheral blood gotten by venipuncture since the patient refused frequent bone marrow puncture.



DOI: 10.12998/wjcc.v11.i18.4295 Copyright ©The Author(s) 2023.

Figure 4 The immunofixation electrophoresis of serum and cytogenetic analysis of bone marrow mononuclear cells. A: The monoclonal IgM Kappa is found by serum immunofixation electrophoresis; B and C: Fluorescence *in situ* hybridization of bone marrow mononuclear cells (BMMNCs) shows a deletion of ATM gene at 11q22 (2G1R 80%, 2G2R 17%, 1G1R 3%) and 4 copies of CCND1 gene at 11q13 (80%); D: The karyotype analysis of BMMNCs detects 46, XY, -2, del(5)(q13q31), -8, add(11)(q13), del(13)(q11q32), +mar1, +mar2 [5] or 46, XY, -2, del(5)(q13q31), -8, add(11)(q13), del(13)(q11q32), +mar1, +mar2 [20] under the stimulation of CpG-oligodeoxynucleotide.

CONCLUSION

The coexistence of DLBCL, AML, and untreated LPL/WM in a same patient is extremely rare. Herein, we reported one such case that might have resulted from multiple steps of gene mutations. In the bone marrow, asymptomatic indolent LPL/WM might have occurred earlier, then insidious MDS developed into AML. Finally, aggressive DLBCL in the chest wall grew. Hematologists should pay close attention to this extremely rare case to avoid misdiagnoses. However, this report has some limitations. Ideally, different clones from BMMNCs should have been separated for molecular analysis, genes from chest wall masses should have been tested, and an autopsy should have been performed.

FOOTNOTES

Author contributions: Zhang LB collected the sample and analyzed the clinical data; Zhang L analyzed the morphology of bone marrow; Xin HL collected the sample and performed gene sequencing; Wang Y analyzed the

histopathology of tissue; Bao HY, Meng QQ, Jiang SY, Han X, and Chen WR analyzed the clinical data; Wang JN and Shi XF provided the conception, analyzed the data, and wrote the manuscript.

Supported by the National Natural Science Foundation of China, No. 81700130; and Nanjing Medical University Science and Technology Development Fund.

Informed consent statement: Informed written consent was obtained from the patient's family for the publication of this report and any accompanying images.

Conflict-of-interest statement: The authors declare that they have no conflict of interest.

CARE Checklist (2016) statement: The authors have read the CARE Checklist (2016), and the manuscript was prepared and revised according to the CARE Checklist (2016).

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Country/Territory of origin: China

ORCID number: Liu-Bo Zhang 0000-0001-8571-4480; Lu Zhang 0000-0002-3707-2011; Hong-Lei Xin 0000-0001-7924-4865; Yan Wang 0000-0003-2057-0097; Hong-Yu Bao 0000-0002-1892-6958; Qing-Qi Meng 0000-0002-8092-8719; Su-Yu Jiang 0000-0002-2516-949X; Xue Han 0000-0002-8756-4209; Wan-Ru Chen 0000-0002-8016-2602; Jian-Ning Wang 0000-0001-9466-2272; Xiao-Feng Shi 0000-0002-8547-711X.

S-Editor: Chen YL

L-Editor: A

P-Editor: Ji MX

REFERENCES

- 1 **Higuchi M**, Sasaki S, Kawadoko S, Uchiyama H, Yasui T, Kamihira T, Aoki K, Sasaguri T, Nakano R, Uchiyama A, Muta T, Ohshima K. Epstein-Barr virus-positive diffuse large B-cell lymphoma following acute myeloid leukemia: a common clonal origin indicated by chromosomal translocation t(3;4)(p25;q21). *Int J Hematol* 2015; **102**: 482-487 [PMID: 25953309 DOI: 10.1007/s12185-015-1802-4]
- 2 **Ririe MR**, Florell SR, Miles RR, Duffy KL. Secondary diffuse large B-cell lymphoma after chemotherapy for acute myeloid leukemia: looking for the unexpected diagnosis. *Am J Dermatopathol* 2014; **36**: e125-e128 [PMID: 24950422 DOI: 10.1097/DAD.0000000000000027]
- 3 **Kunitomi A**, Kotani S, Ukyo N, Ono K, Nakamine H, Nohgawa M. Epstein-Barr virus-positive diffuse large B-cell lymphoma of the elderly complicated by the onset of acute myeloid leukemia. *Intern Med* 2014; **53**: 51-56 [PMID: 24390529 DOI: 10.2169/internalmedicine.53.0219]
- 4 **Bumbea H**, Popov VM, Tomuleasa C, Omer M, Dobrea C, Manea I, Zurac S, Popp C, Dumitru I, Simoiu M, Mastalier B. Coexistence of Trisomy 8 and 13 in a Newly Diagnosed Patient With Diffuse Large B Cell Non-Hodgkin Lymphoma and Acute Myeloid Leukemia Secondary to Primary Myelofibrosis. *Cureus* 2022; **14**: e22217 [PMID: 35186608 DOI: 10.7759/cureus.22217]
- 5 **Ito S**, Fujiwara SI, Mashima K, Umino K, Minakata D, Nakano H, Yamasaki R, Kawasaki Y, Sugimoto M, Ashizawa M, Yamamoto C, Hatano K, Okazuka K, Sato K, Oh I, Ohmine K, Suzuki T, Muroi K, Kanda Y. Development of acute myeloid leukemia in patients with untreated chronic lymphocytic leukemia. *Ann Hematol* 2017; **96**: 719-724 [PMID: 28144729 DOI: 10.1007/s00277-017-2933-x]
- 6 **Stern N**, Shemesh J, Ramot B. Chronic lymphatic leukemia terminating in acute myeloid leukemia: review of the literature. *Cancer* 1981; **47**: 1849-1851 [PMID: 6939479 DOI: 10.1002/1097-0142(19810401)47:7<1849::aid-cnrcr2820470722>3.0.co;2-x]
- 7 **Milosevic I**. Coexistence of Chronic Lymphocytic Leukemia and Acute Myeloid Leukemia. *Turk J Haematol* 2016; **33**: 353-354 [PMID: 27751969 DOI: 10.4274/tjh.2016.0106]
- 8 **Al Mussaed E**, Osman H, Elyamany G. Simultaneous existence of acute myeloid leukemia and chronic lymphocytic leukemia: a case report. *BMC Cancer* 2016; **16**: 739 [PMID: 27643996 DOI: 10.1186/s12885-016-2780-5]
- 9 **Chen RR**, Zhu LX, Wang LL, Li XY, Sun JN, Xie MX, Zhu JJ, Zhou D, Li JH, Huang X, Xie WZ, Ye XJ. Synchronous diagnosis and treatment of acute myeloid leukemia and chronic lymphocytic leukemia: Two case reports. *World J Clin Cases* 2021; **9**: 9144-9150 [PMID: 34786398 DOI: 10.12998/wjcc.v9.i30.9144]
- 10 **Shoye O**, Gupta G. Synchronous Diagnosis of De Novo Acute Myeloid Leukemia with inv(16)(p13q22) and Chronic Lymphocytic Leukemia: A Case Report and Review of the Literature. *Ann Clin Lab Sci* 2018; **48**: 790-796 [PMID: 30610052 DOI: 10.1016/j.medcle.2019.03.004]
- 11 **Lynch RC**, Gratzinger D, Advani RH. Clinical Impact of the 2016 Update to the WHO Lymphoma Classification. *Curr Treat Options Oncol* 2017; **18**: 45 [PMID: 28670664 DOI: 10.1007/s11864-017-0483-z]

- 12 **te Raa GD**, Malcikova J, Pospisilova S, Trbusek M, Mraz M, Garff-Tavernier ML, Merle-Béral H, Lin K, Pettitt AR, Merkel O, Stankovic T, van Oers MH, Eldering E, Stilgenbauer S, Zenz T, Kater AP; European Research Initiative on CLL (ERIC). Overview of available p53 function tests in relation to TP53 and ATM gene alterations and chemoresistance in chronic lymphocytic leukemia. *Leuk Lymphoma* 2013; **54**: 1849-1853 [PMID: [23614766](#) DOI: [10.3109/10428194.2013.796058](#)]
- 13 **Skowronska A**, Parker A, Ahmed G, Oldreive C, Davis Z, Richards S, Dyer M, Matutes E, Gonzalez D, Taylor AM, Moss P, Thomas P, Oscier D, Stankovic T. Biallelic ATM inactivation significantly reduces survival in patients treated on the United Kingdom Leukemia Research Fund Chronic Lymphocytic Leukemia 4 trial. *J Clin Oncol* 2012; **30**: 4524-4532 [PMID: [23091097](#) DOI: [10.1200/JCO.2011.41.0852](#)]
- 14 **Gómez-Arbonés J**, Gallart MA, Mellado A, Marco V, Panadés MJ, Macià JM. Concomitant diagnosis of acute myeloid leukemia (AML) and chronic lymphocytic leukemia (CLL). Importance of flow cytometry in the diagnosis of CLL without lymphocytosis accompanying AML. *Eur J Haematol* 1997; **59**: 335-337 [PMID: [9414649](#) DOI: [10.1111/j.1600-0609.1997.tb01698.x](#)]
- 15 **Lima M**, Porto B, Rodrigues M, Teixeira MA, Coutinho J, Ribeiro AC, Malheiro MI, Justiça B. Cytogenetic findings in a patient presenting simultaneously with chronic lymphocytic leukemia and acute myeloid leukemia. *Cancer Genet Cytogenet* 1996; **87**: 38-40 [PMID: [8646738](#) DOI: [10.1016/0165-4608\(95\)00262-6](#)]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

Telephone: +1-925-3991568

E-mail: bpgoffice@wjgnet.com

Help Desk: <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

