

Expression and location of Smad2, 4 mRNAs during and after liver fibrogenesis of rats

Yang Liu, Li-Feng Wang, Hai-Feng Zou, Xiao-Yan Song, Hua-Feng Xu, Ping Lin, Hai-Hong Zheng, Xiao-Guang Yu

Yang Liu, Hai-Feng Zou, Hua-Feng Xu, Ping Lin, Xiao-Guang Yu, Department of Biochemistry and Molecular Biology, Harbin Medical University, Harbin 150086, Heilongjiang Province, China

Li-Feng Wang, Department of Pathology of the First Clinical College, Harbin Medical University, Harbin 150001, Heilongjiang Province, China

Xiao-Yan Song, Master degree student of Department of Pathology of Harbin Medical University, Harbin 150086, Heilongjiang Province, China

Hai-Hong Zheng, Department of Experimental Animal of the Second Clinical College, Harbin Medical University, Harbin 150086, Heilongjiang Province, China

Supported by the Research Foundation of Department of Education of Heilongjiang Province, No.10551131

Correspondence to: Dr. Xiao-Guang Yu, Department of Biochemistry and Molecular Biology, Harbin Medical University, Harbin 150086, Heilongjiang Province, China. coollya@gmail.com

Telephone: +86-451-86671684

Received: 2005-03-22

Accepted: 2005-07-21

Abstract

AIM: To investigate the location alteration of Smad2 and Smad4 mRNAs in the liver during and after fibrogenesis in rats.

METHODS: Eighty male Wistar rats weighing approximately 200 g each were used. The rat models of experimental hepatic fibrosis were established by injection with carbon tetrachloride (CCl₄), normal rats and rats were injected with olive oil and served as control groups. *In situ* hybridization (ISH) was used to detect the Smad2 and Smad4 mRNA in liver.

RESULTS: *In situ* hybridization showed Smad2 and Smad4 mRNA expressions in the cytoplasm of hepatic stellate cells (HSC), fibroblasts and myofibroblasts around the central vein and hepatic sinus during and after fibrogenesis. Expression of Smad2, 4 mRNA was higher than that in normal and control rats.

CONCLUSION: In the process of and after hepatic fibrosis formation, HSC, fibroblasts and myofibroblasts are the major cells that express Smad2 and Smad4. The more serious the hepatic fibrosis is in the injured liver, the higher the level of Smad2 and Smad4 gene expression is during and after fibrogenesis respectively.

© 2006 The WJG Press. All rights reserved.

Key words: Smad2; Smad4; mRNA; Liver fibrogenesis

Liu Y, Wang LF, Zou HF, Song XY, Xu HF, Lin P, Zheng HH, Yu XG. Expression and location of Smad2, 4 mRNA during and after liver fibrogenesis of rats. *World J Gastroenterol* 2006; 12(10): 1577-1582

<http://www.wjgnet.com/1007-9327/12/1577.asp>

INTRODUCTION

Fibrosis is associated with many liver diseases, including hepatitis C virus infection, iron deposition, alcohol consumption, and nonalcoholic fatty liver disease. Hepatic fibrosis results from a net increased synthesis and decreased degradation of extracellular matrix (ECM) proteins. Type I collagen is the most prevalent ECM protein deposited^[1], with activated hepatic stellate cells (HSCs) serving as the primary source. Following a fibrogenic stimulus, HSCs are activated from their normal quiescent state, whereby they increase synthesis of procollagen type I protein^[1,2], and increase cellular proliferation, migration, and contractility^[3,4]. Excess ECM accumulation results in scarring within the tissue^[5].

In the progression of HSCs activation, many kinds of cytokines play important roles, including transforming growth factor-beta (TGF- β)^[6]. TGF- β is a multifunctional cytokine with diverse effects on development, growth, and homeostasis in most tissues^[7]. Signaling by TGF- β occurs through type I and type II transmembrane serine/threonine kinase receptors and intracellular Smad transduction molecules. The TGF- β ligand binds to its type II receptor, either directly or via coreceptors. Once activated by TGF- β , type II receptors recruit, bind, and transphosphorylate type I receptors, thereby stimulating protein kinase activity of the later. The activated type I receptors phosphorylate Smad2 or Smad3, which then bind to Smad4. The resulting Smad complex then moves into the nucleus, where it interacts in a cell-specific manner with various transcription factors to regulate the transcription of target genes^[8].

It has been reported that there is a correlation between TGF- β and the formation of liver fibrosis^[9,10]. TGF- β has a very important role in ECM production by HSC and many investigations have been done^[11,12]. Antagonism of cytokines as a route to antifibrotic therapy, by blocking the interaction between TGF- β and its receptors, and the signal transduction mechanism for TGF- β , has been reported for the treatment of fibrosis^[13,14]. Thus the investigation of the Smads signal transduction is indispensable for the study

of fibrosis and the development of its therapy. Until now, there have been no reports on the location and expression alteration of Smad2,4 during and after fibrogenesis. In this article, the expression and location of Smad2,4 mRNA were measured in rat livers with cRNA probes of Smad2,4 by *in situ* hybridization.

MATERIALS AND METHODS

Animal experiments

Eighty male Wistar rats weighing approximately 200 g each were used. Chronic liver injury was produced in 10 rats in each group by subcutaneous injections of 50% CCl₄ in olive oil at a volume of 2 ml/kg body weight twice weekly for 3, 6, 9, 12, 15, and 18 wk. Ten control rats similarly received olive oil alone as control group, and 10 other rats were not treated as normal group. All rats were maintained on a standard diet with water *ad libitum*. On the sixth day after the last treatment, 10 rats in each group were decapitated. Their livers were removed immediately, fixed in 4% paraformaldehyde at room temperature and snap-frozen in liquid nitrogen. The tissue specimens were stored at -80 °C until analysis.

Cloning of Smad2 and Smad4 cDNAs

Smad2 and Smad4 cDNAs were cloned in order to synthesize ISH riboprobes. The two cDNAs were obtained from rat liver total RNA by reverse transcription coupled to PCR by the use of the primers Smad2-up (5' GGTGGATCCGTGTCTCATCGGAAAGGG-3'), Smad2-GTGAATTCGGAATGGAGTGGGTATAG-3') and Smad4-up (5'-AACCGGATCCATCTTTCAGC ACCACCC-3'), Smad4-down (5'CGAATTCCTTGCCTA TGTGCAACCTTGC-3'), respectively. The two resulting fragments were digested with *Bam*H I and *Eco*R I and cloned into the pSPT18 vector (Boehringer Mannheim) to obtain the pSPT18+/Smad2 and pSPT18+/Smad4 plasmids.

ISH riboprobe synthesis

The cDNA-containing plasmids were linearized as follows: pSPT18+/Smad2 and pSPT18+/Smad4 with *Bam*H I and *Eco*R I. The linearized plasmids were then gel-isolated (Hua Shun Corp., China) and used as templates for anti-sense and sense digoxigenin (DIG)-labeled riboprobe synthesis (Boehringer Mannheim). The transcription mixture (20 µL) included 1 µg of linearized template cDNA, ATP, GTP and CTP at 1 mmol/L each, UTP 0.7 mmol/L, DIG-UTP 0.3 mmol/L, DTT 10 mmol/L, RNase inhibitor (1 U/µL of transcription mix), and T3 or T7 RNA polymerase (1 U/µL of transcription mix). Transcription was performed for at least 2 hr at 37 °C. The template cDNAs were then digested by RNase-free DNase (2 µL at 1 U/µL, 15 min at 37 °C), and all reactions were stopped by adjusting the reaction volume to 100 µL with EDTA (pH8.0). The riboprobes were then purified through two precipitation steps by addition of 100 µL LiCl (4 mmol/L) and 500 µL EtOH (100%), and centrifugation for 30 min at 4 °C in a microfuge. The pellet was resuspended in 100 µL DECP-treated water.

In situ hybridization

The serial paraffin sections (thickness 4 µm) were dried at 55 °C, then deparaffinized by xylene and rehydrated in graded ethanol, incubated for 2×5 min in TBS containing 0.1% active DEPC (Fluka), acidified in 0.2 N HCl for 20 min at room temperature, and incubated for 2×5 min in TBS before digestion with proteinase K for 15 min, and then incubated for 2×5 min in TBS. After postfixation in 4% paraformaldehyde-PBS, sections were incubated for 2×5 min in TBS. And then sections were dehydrated with graded ethanol. After prehybridization at 42 °C for 2 h, the labeled RNA probes were added to the hybridization mix (Smad2 3.7 ng/µL, Smad4 2.3 ng/µL). The hybridization was carried out at 42 °C for 18 h with 20µL of hybridization mix on each section. The sections were washed with 2×SSC, 1×SSC, and TBST (TBS containing Tween20) respectively. After washed, the sections were incubated for 15 min at room temperature with Buffer I containing 1% blocking reagent (Boehringer Mannheim), and then at 37 °C with alkaline phosphatase-coupled anti-digoxigenin antibody (Boehringer Mannheim) diluted at 1:50 in Buffer I. Excess antibody was removed by 5×4 min washes in TBST, and the sections were equilibrated for 2×5 min in Buffer I (Tris-HCl 100mmol/L, NaCl 100mmol/L, and MgCl₂ 100mmol/L, pH9.5). Color development was performed at 37 °C for 8 h in Buffer I containing NBT and BCIP (Promega). Staining was stopped by a 10 min washing in tap water, and non-special staining was removed in EtOH 95% with gentle agitation. Then the sections were rehydrated through successive baths of EtOH (70, 95, and 100%) and xylol (2×15 min each) and mounted in gum for microscopic examination and photography. Blank control: Smad2 and Smad4 cRNA probes for positive hepatic tissues were replaced by prehybridization solution. Negative control: *in situ* hybridization was performed in normal liver tissues. False positive control: *in situ* hybridization was performed in liver tissues injected with olive oil.

Pathology observation

The whole liver sections were stained with hematoxylin and eosin, and examined under microscope.

Statistic analysis

The *t* test was used to determine statistical significance between groups. *P* value of less than 0.05 was considered statistically significant.

RESULTS

Pathologic findings

During the hepatic fibrosis formation in the experimental rats, in liver tissues at 3 wk, infiltration of inflammatory cells was found around the portal area and central vein; at 6 wk, fatty degeneration and infiltration of inflammatory cells was detected; at 9 wk, hyperplasia of the lattice fibers and collagenous fibers was observed in portal area; at 12 wk, hyperplasia of the lattice fibers and collagenous fibers was observed in portal area and extended outwards, hyperplasia surrounding the central vein observed was




Figure 1 Pathology observation of the experimental rat liver sections stained with hematoxylin and eosin ($\times 100$). **A**: normal rats; **B**: control rats; **C**: CCl₄ treated rats at 3 wk; **D**: CCl₄ treated rats at 6 wk; **E**: CCl₄ treated rats at 9 wk; **F**: CCl₄ treated rats at 12 wk; **G**: CCl₄ treated rats at 15 wk; **H**: CCl₄ treated rats at 18 wk.




Figure 2 Pathology observation of the experimental rat liver sections stained with hematoxylin and eosin ($\times 400$). The arrow shows atypical hyperplasia in the epithelium of bile capillary. **A, B, C, D, E, F, G, H**, see Figure 1.

distributed along hepatic sinus and connected with each other, the hepatic lobules were encysted and separated by collagen bundles, the normal structure of lobules was destroyed, and pseudolobules formed, infiltration of inflammatory cells was found around the portal area and central vein; at 15 and 18 wk, excluding the findings of 12 wk, atypical hyperplasia was found in the epithelium of biliary capillary. The structure of liver tissues was normal in normal and control groups (Figures 1 and 2).

Smad2 and Smad4 location

Smad2 and Smad4 mRNAs in liver from experimental rats were detected mainly in the HSCs, fibroblasts and myofibroblasts around the portal area and central vein. Expressions of Smad2 and Smad4 mRNAs exhibited as blue particles in cytoplasm. No positive expression was found in nuclei. There was only mild positive expression in normal and control groups. Image pattern analysis showed

that the expression in the experimental group was much stronger than that in the normal and control groups (Table 1, Figures 3 and 4).

DISCUSSION

To establish the model of liver fibrosis successfully is the key for detection of Smad2 and Smad4 mRNAs in the liver tissue during and after fibrogenesis. Researches for establishing the model of liver fibrosis with CCl₄ began in 1936. After that many methods to establish the model of liver fibrosis have been tried. Among them, liver fibrosis models induced by immune response or CCl₄ have been generally accepted mainly because they are more close to the disease characteristics in terms of the distinct stages of the disease of human body and a low mortality. In this study, the method of injecting CCl₄ was used to establish the model of liver fibrosis, and it was proved successful

Table 1 Expression of Smad2 and Smad4 mRNA during and after liver fibrogenesis (mean \pm SD)

Group	n	Nor	Con	3wk	6wk	9wk	12wk	15wk	18w
Smad2	80	120.20 \pm 9.00	113.30 \pm 15.18	150.50 \pm 8.13 ^{ce}	213.40 \pm 13.92 ^{ace}	310.30 \pm 25.43 ^{ace}	510.80 \pm 7.53 ^{ace}	523.50 \pm 17.35 ^{ace}	568.40 \pm 7.64 ^{ace}
Smad4	80	99.40 \pm 3.65	105.40 \pm 10.89	143.50 \pm 6.47 ^{ce}	165.00 \pm 32.10 ^{ace}	305.10 \pm 18.53 ^{ace}	485.00 \pm 4.08 ^{ace}	500.40 \pm 10.94 ^{ace}	529.00 \pm 6.78 ^{ace}

^a $P < 0.05$ vs a previous time point, ^c $P < 0.05$ vs normal group, ^e $P < 0.05$ vs control group.




Figure 3 Expression of Smad2 mRNA in experimental rats liver sections during and after fibrosis (*in situ* hybridization $\times 400$). A, B, C, D, E, F, G, H, see Figure 1.




Figure 4 Expression of Smad4 mRNA in experimental rats liver sections during and after fibrosis (*in situ* hybridization $\times 400$). A, B, C, D, E, F, G, H, see Figure 1.

by histochemical analysis of the sections stained with hematoxylin and eosin. According to the pathologic findings of this study, the exact time for establishment of rat hepatic cirrhosis by injecting CCl₄ may be at 12 wk. From 15 and 18 wk on, atypical hyperplasia was found in the epithelium of bile capillary. It may indicate that injecting CCl₄ may cause hepatic carcinoma.

TGF- β consists of a big molecular family that has a common structure including BMP and activin. These proteins have many roles in the cell structure and function,

such as cell growth, differentiation, apoptosis and regulation of ECM production^[15,16]. The main functions of the TGF- β super family are morphogenesis, inflammation, tissue recovery and oncogenesis^[17]. Smads are intracellular signal transductive molecules of the TGF- β super family. These molecules were discovered as Mothers against dpp (Mad)^[18,19], or sma^[20] and named Smad in 1996^[21]. According to the differences of structure and function, nine Smads have been reported and classified into three groups^[17,22]. Smads 2 and 3 are named R-Smads in the

pathway and Smad 4 a Co-Smads for all these pathways. Smads 6, 7, 8 and Xsmad 8 are inhibitory factors of these Smads. When TGF- β binds to its receptor, Smad 2/3 is phosphorylated and binds with Smad 4, together they move into the nucleus for translation and expression of the target gene^[7,23]. These mechanisms are thought to play a role in the process of liver damage, recovery, as well as liver fibrosis. In a damaged liver including chronic inflammation, in hepatitis B, TGF- β signaling was reported^[24]. Not only Kupffer cells but also the HSC produce TGF- β by the autocrine system^[25-28]. TGF- β 1 stimulation evokes ECM production, which develops into liver fibrosis.

In this study, Smad2 and Smad4 mRNAs were investigated in rat liver during and after fibrogenesis caused by CCl₄ *in vivo*. With the method of *in situ* hybridization, we found Smad2 and Smad4 mRNAs were expressed obviously in HSC, myofibroblasts and fibroblasts of livers of liver fibrosis group, mostly in portal area and fibrous septum, while very mildly in vascular endothelial cells of livers from normal and control groups. While with the development of liver fibrosis, expressions of Smad2 and Smad4 mRNAs were increased also. These results indicate that Smad2 and Smad4 play an important role in the progression of liver fibrosis. In the sections of liver fibrosis at 15 and 18 wk, expressions of Smad2 and Smad4 mRNAs were higher in the epithelium of bile capillary than in normal and control groups. At this stage, there was atypical hyperplasia in the epithelium of bile capillary, indicating early stage of carcinoma. Therefore, Smad2 and Smad4 may be associated with cancerogenesis.

After stimulation of TGF- β , Smad2, 3 and 4 proteins translocate from cytoplasm to nucleus in MV1Lu cells, analyzed by immunofluorescence using specific antiserum^[22]. Moreover, Smad3 seems to play the most important role in liver fibrosis, because Smad3 interacts with SP1 which increases expression of ECM by regulating COL1A2 gene^[29]. By transfection of Smad3 expression plasmid in CF37 cells, ECM also increases by regulating COL1A2 gene. Nevertheless, COL1A2 transcription in CF37 cells is markedly increased by TGF- β after transfection with a Smad2 expression plasmid^[30]. So Smad2 also plays an important role in liver fibrosis. Our findings are consistent with the above results. The difference is perhaps due to use of different cell lines or experimental conditions. Further researches are needed to confirm the results.

As the liver fibrosis is a complex process, involving many cytokines, cells, and signal transduction pathways, the mechanism cannot be fully explained by these experiments. However, it is clear that TGF/Smads signal transduction pathway plays an important role in liver fibrosis which has been proved by many researches. In our study, Smad2 and Smad4 mRNAs were detected in the plasma of HSC, fibroblasts and myofibroblasts around the portal area and central vein by *in situ* hybridization. Expressions of Smad2 and Smad4 mRNAs were higher during and after liver fibrogenesis than normal and control groups, indicating that they play an important role in liver fibrosis.

REFERENCES

- Brenner DA, Waterboer T, Choi SK, Lindquist JN, Stefanovic B, Burchardt E, Yamauchi M, Gillan A, Rippe RA. New aspects of hepatic fibrosis. *J Hepatol* 2000; **32**: 32-38
- Stefanovic B, Hellerbrand C, Holcik M, Briendl M, Aliehbhaber S, Brenner DA. Posttranscriptional regulation of collagen alpha1(I) mRNA in hepatic stellate cells. *Mol Cell Biol* 1997; **17**: 5201-5209
- Pinzani M, Marra F, Carloni V. Signal transduction in hepatic stellate cells. *Liver* 1998; **18**:2-13
- Reeves HL, Friedman SL. Activation of hepatic stellate cells--a key issue in liver fibrosis. *Front Biosci* 2002; **7**: d808-826
- Benyon RC, Arthur MJ. Extracellular matrix degradation and the role of hepatic stellate cells. *Semin Liver Dis* 2001; **21**: 373-384
- Zhang LJ, Yu JP, Li D, Huang YH, Chen ZX, Wang XZ. Effects of cytokines on carbon tetrachloride-induced hepatic fibrogenesis in rats. *World J Gastroenterol* 2004; **10**: 77-81
- Heldin CH, Miyazono K, ten Dijke P. TGF-beta signalling from cell membrane to nucleus through SMAD proteins. *Nature* 1997; **390**: 465-471
- Shi Y, Massague J. Mechanisms of TGF-beta signaling from cell membrane to the nucleus. *Cell* 2003; **113**: 685-700
- Friedman SL. Seminars in medicine of the Beth Israel Hospital, Boston. The cellular basis of hepatic fibrosis. Mechanisms and treatment strategies. *N Engl J Med* 1993; **328**: 1828-1835
- Burt AD. C. L. Oakley Lecture (1993). Cellular and molecular aspects of hepatic fibrosis. *J Pathol* 1993; **170**: 105-114
- Tseng SC, Lee PC, Ells PF, Bissell DM, Smuckler EA, Stern R. Collagen production by rat hepatocytes and sinusoidal cells in primary monolayer culture. *Hepatology* 1982; **2**: 13-18
- Knittel T, Schuppan D, Meyer zum Buschenfelde KH, Ramadori G. Differential expression of collagen types I, III, and IV by fat-storing (Ito) cells in vitro. *Gastroenterology* 1992; **102**: 1724-1735
- Casini A, Pinzani M, Milani S, Grappone C, Galli G, Jezequel AM, Schuppan D, Rotella CM, Surrenti C. Regulation of extracellular matrix synthesis by transforming growth factor beta 1 in human fat-storing cells. *Gastroenterology* 1993; **105**: 245-253
- Border WA, Noble NA, Yamamoto T, Harper JR, Yamaguchi Y, Pierschbacher MD, Ruoslahti E. Natural inhibitor of transforming growth factor-beta protects against scarring in experimental kidney disease. *Nature* 1992; **360**: 361-364
- Zhang K, Rekhter MD, Gordon D, Phan SH. Myofibroblasts and their role in lung collagen gene expression during pulmonary fibrosis. A combined immunohistochemical and *in situ* hybridization study. *Am J Pathol* 1994; **145**: 114-125
- Border WA, Noble NA. Transforming growth factor beta in tissue fibrosis. *N Engl J Med* 1994; **331**: 1286-1292
- Derynck R, Feng XH. TGF-beta receptor signaling. *Biochim Biophys Acta* 1997; **1333**: F105-150
- Rafferty LA, Twombly V, Wharton K, Gelbart WM. Genetic screens to identify elements of the decapentaplegic signaling pathway in *Drosophila*. *Genetics* 1995; **139**: 241-254
- Sekelsky JJ, Newfeld SJ, Rafferty LA, Chartoff EH, Gelbart WM. Genetic characterization and cloning of mothers against dpp, a gene required for decapentaplegic function in *Drosophila melanogaster*. *Genetics* 1995; **139**: 1347-1358
- Savage C, Das P, Finelli AL, Townsend SR, Sun CY, Baird SE, Padgett RW. *Caenorhabditis elegans* genes sma-2, sma-3, and sma-4 define a conserved family of transforming growth factor beta pathway components. *Proc Natl Acad Sci USA* 1996; **93**: 790-794
- Derynck R, Gelbart WM, Harland RM, Heldin CH, Kern SE, Massague J, Melton DA, Mlodzik M, Padgett RW, Roberts AB, Smith J, Thomsen GH, Vogelstein B, Wang XF. Nomenclature: vertebrate mediators of TGF-beta family signals. *Cell* 1996; **87**: 173
- Nakao A, Imamura T, Souchelnytskyi S, Kawabata M, Ishisaki A, Oeda E, Tamaki K, Hanai J, Heldin CH, Miyazono K, ten Dijke P. TGF-beta receptor-mediated signalling through

- Smad2, Smad3 and Smad4. *EMBO J* 1997; **16**: 5353-5362
- 23 **Nakao A**, Afrakhte M, Moren A, Nakayama T, Christian JL, Heuchel R, Itoh S, Kawabata M, Heldin NE, Heldin CH, ten Dijke P. Identification of Smad7, a TGFbeta-inducible antagonist of TGF-beta signalling. *Nature* 1997; **389**: 631-635
- 24 **Lee DK**, Park SH, Yi Y, Choi SG, Lee C, Parks WT, Cho H, de Caestecker MP, Shaul Y, Roberts AB, Kim SJ. The hepatitis B virus encoded oncoprotein pX amplifies TGF-beta family signaling through direct interaction with Smad4: potential mechanism of hepatitis B virus-induced liver fibrosis. *Genes Dev* 2001; **15**: 455-466
- 25 **Matsuzaki K**, Date M, Furukawa F, Tahashi Y, Matsushita M, Sugano Y, Yamashiki N, Nakagawa T, Seki T, Nishizawa M, Fujisawa J, Inoue K. Regulatory mechanisms for transforming growth factor beta as an autocrine inhibitor in human hepatocellular carcinoma: implications for roles of smads in its growth. *Hepatology* 2000; **32**: 218-227
- 26 **Bissell DM**, Wang SS, Jarnagin WR, Roll FJ. Cell-specific expression of transforming growth factor-beta in rat liver. Evidence for autocrine regulation of hepatocyte proliferation. *J Clin Invest* 1995; **96**: 447-455
- 27 **Milani S**, Herbst H, Schuppan D, Stein H, Surrenti C. Transforming growth factors beta 1 and beta 2 are differentially expressed in fibrotic liver disease. *Am J Pathol* 1991; **139**: 1221-1229
- 28 **Dooley S**, Delvoux B, Lahme B, Mangasser-Stephan K, Gressner AM. Modulation of transforming growth factor beta response and signaling during transdifferentiation of rat hepatic stellate cells to myofibroblasts. *Hepatology* 2000; **31**: 1094-1106
- 29 **Flanders KC**. Smad3 as a mediator of the fibrotic response. *Int J Exp Pathol* 2004; **85**: 47-64
- 30 **Inagaki Y**, Mamura M, Kanamaru Y, Greenwel P, Nemoto T, Takehara K, Ten Dijke P, Nakao A. Constitutive phosphorylation and nuclear localization of Smad3 are correlated with increased collagen gene transcription in activated hepatic stellate cells. *J Cell Physiol* 2001; **187**: 117-123

S- Editor Wang J L- Editor Zhu LH E- Editor Bi L