

全部

图片

视频

新闻

更多

设置

工具

找到约 60 条结果 (用时 0.80 秒)

Design and Validation of a Thermoreversible Material for ... - NCBI

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - 翻译此页

作者: A Johnson - 2013 - 被引用次数: 9 - 相关文章

Such a material can be injected as a liquid which then forms a semi-solid gel at ... Ex vivo liver ablations showed that P407 protects neighboring tissues, but may ... Keywords: ablation, poloxamer 407, hydrodissection, D5W, thermoreversible ... One possible side effect of more aggressive treatment is thermal damage to ...

[Effect of thermo-sensitive Matrigel on minimization of thermal injury to ...

www.ncbi.nlm.nih.gov/pubmed/26792607 - 翻译此页

作者: JJ Fu - 2016

2016年1月5日 - [Effect of thermo-sensitive Matrigel on minimization of thermal injury to the nearby structures in radiofrequency ablation of subcapsular hepatic tumors in a rat model]. ... (90 ± 2) °C, 41 hepatic ablations was performed abutting the diaphragm, ... in vials at temperature 37 °C in 1 min, presenting gel formation.

缺少字词: poloxamer 407 based microwave

[PDF] Final Report 1.95 MB Wednesday, December 8, 2010 - UW BME De...

bmedesign.engr.wisc.edu/projects/file/?fid=1534 - 翻译此页

2010年12月6日 - The poloxamer 407 (Lutrol F-127) solution was most favorable for future ... Damage to the diaphragm may result in ... During thermal ablation procedures, the hydrodissection fluid is ... the past 15 years; these include: RF ablation, microwave ablation, ... because the gelation is not based on temperature.

[PDF] Midsemester Paper 1.74 MB Wednesday, March 9 ... - UW BME Des...

bmedesign.engr.wisc.edu/projects/file/?fid=1780 - 翻译此页

2011年3月9日 - Thus, a 19.0 %w/w poloxamer 407 solution that would gel at 32 o ... Seventy to ninety percent of these liver ... Damage to the diaphragm may result in slight pain while ... During thermal ablation procedures, the hydrodissection fluid is ... the past 15 years; these include: RF

Name of journal: *World Journal of Gastroenterology*

ESPS Manuscript NO: 31877

Manuscript Type: ORIGINAL ARTICLE

Basic Study

Effect of a poloxamer 407-based thermosensitive gel on minimization of thermal injury to diaphragm during microwave ablation of the liver

Li-Li Zhang, Gui-Min Xia, Yu-Jiang Liu, Rui Dou, John Eisenbrey, Ji-Bin Liu,

Xiao-Wei Wang, Lin-Xue Qian

Abstract

Match Overview

- | Match Number | Source | Words | Similarity |
|--------------|---|----------|------------|
| 1 | Crossref | 62 words | 2% |
| | Zhang, Dedong, Daolin Xie, Xiaoli Wei, Dakun Zhang, Min Chen, Xiaoling Yu, and Ping Liang. "Microwave ablation of t | | |
| 2 | Internet | 14 words | <1% |
| | crawled on 30-Mar-2010 www.ajronline.org | | |

Effect of a poloxamer 407-based thermosensitive gel on minimization of thermal i

全部 图片 视频 新闻 地图 图书

找到约 82 条结果

时间不限

过去 1 小时内
过去 24 小时内
过去 1 周内
过去 1 个月内
过去 1 年内

所有结果

精确匹配

[Design and Validation of a Thermoreversible Material for ...](#)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4281175/>

However, currently used fluids such as normal saline and 5% dextrose in ... The P407 gel was similar to other hydrodissection fluids at ultrasound and CT imaging. ... Keywords: **ablation**, **poloxamer 407**, hydrodissection, D5W, thermoreversible ... One possible side **effect** of more aggressive treatment is **thermal damage** to ...

[PDF] [Midsemester Paper 1.74 MB Wednesday, March ... - UW BME Design](#)

bmedesign.engr.wisc.edu/projects/file/?fid=1780

9 Mar 2011 ... for the treatment of malignant tumors of the **liver**, lungs, and kidneys. ... Thus, a 19.0 %w/w **poloxamer 407** solution that would **gel** at 32 o. C ... **Damage** to the **diaphragm** may result in slight pain while ... **During thermal ablation** procedures, the hydrodissection fluid is injected between the ... be **minimized**.

[PDF] [Final Paper, Final PDS at end 1.81 MB Tuesday ... - UW BME Design](#)

bmedesign.engr.wisc.edu/projects/file/?fid=1839

4 May 2011 ... for treatment of malignant tumors of the **liver**, lungs, and kidneys. ... **poloxamer 407** solution that would **gel** at 32 ... determine their **effect** on the gelation temperature and kinematic viscosity **During thermal ablation** procedures, the hydrodissection fluid is D5W **minimizes** unwanted tissue **damage** most.

[PDF] [Final Report 1.95 MB Wednesday, December 8 ... - UW BME Design](#)

bmedesign.engr.wisc.edu/projects/file/?fid=1534

6 Dec 2010 ... The **poloxamer 407** (Lutrol F-127) solution was most favorable for future ... **Damage** to the **diaphragm** may result in ... **During thermal** ablation procedures, the hydrodissection fluid is ... the past 15 years, these include: RF ablation, **microwave ablation**, because the gelation is not **based** on temperature.

[Diaphragmatic rupture - Wikipedia](#)

[全部](#)[图片](#)[视频](#)[购物](#)[新闻](#)[更多](#)[设置](#)[工具](#)

找到约 82 条结果 (用时 0.91 秒)

[Design and Validation of a Thermoreversible Material for ... - NCBI - NIH](#)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4281175/> - [翻译此页](#)

作者: A Johnson - 2013 - 被引用次数: 9 - [相关文章](#)

However, currently used fluids such as normal saline and 5% dextrose in ... The P407 gel was similar to other hydrodissection fluids at ultrasound and CT imaging. ... Keywords: ablation, poloxamer 407, hydrodissection, D5W, thermoreversible ... One possible side effect of more aggressive treatment is thermal damage to ...

[\[PDF\] Midsemester Paper 1.74 MB Wednesday, March 9 ... - UW BME Des...](#)

bmedesign.engr.wisc.edu/projects/file/?fid=1780 - [翻译此页](#)

2011年3月9日 - for the treatment of malignant tumors of the liver, lungs, and kidneys. ... Thus, a 19.0 %w/w poloxamer 407 solution that would gel at 32 o. C ... Damage to the diaphragm may result in slight pain while ... During thermal ablation procedures, the hydrodissection fluid is injected between the ... be minimized.

[\[PDF\] Final Paper, Final PDS at end 1.81 MB Tuesday ... - UW BME Design](#)

bmedesign.engr.wisc.edu/projects/file/?fid=1839 - [翻译此页](#)

2011年5月4日 - for treatment of malignant tumors of the liver, lungs, and kidneys. ... poloxamer 407 solution that would gel at 32 ... determine their effect on the gelation temperature and kinematic viscosity During thermal ablation procedures, the hydrodissection fluid is D5W minimizes unwanted tissue damage most.

[\[PDF\] Final Report 1.95 MB Wednesday, December 8, 2010 - UW BME De...](#)

bmedesign.engr.wisc.edu/projects/file/?fid=1534 - [翻译此页](#)

2010年12月6日 - The poloxamer 407 (Lutrol F-127) solution was most favorable for future ... Damage to the diaphragm may result in ... During thermal ablation procedures, the hydrodissection fluid is ... the past 15 years; these include: RF ablation, microwave ablation, because the gelation is not