

1

Name of journal World Journal of Gastroenterology

ESPS Manuscript NO: 16558

Column: ORIGINAL ARTICLE

Prospective Study

Evolution of nonspecific duodenal lymphocytosis over 2 years of follow-up

Giuseppe Leonardo, Domenico Piccirilli, Antonio Giangaspero, Mariabastrika Principi, Francesca Buffelli, Flotiana Giorgio, Lucia Montenegro, Claudia Sorrentino, Annachiara Amoroso, Enzo Ierardi, Alfredo Di Leo

Abstract

AIM To assess the evolution of duodenal lymphocytosis (DL), a condition characterized

Match Overview





Evolution of nonspecific duodenal lymphocytosis over 2 years of follow-u



网页

图片

新闻

视频

更多 ▾

搜索工具

找到约 37,400 条结果 (用时 0.59 秒)

Modern Pathology - Duodenal intraepithelial lymphocytosis ...

www.nature.com > [Journal home](#) > [Archive](#) > [Original Articles](#) - [翻译此页](#)

作者: L Memeo - 2005 - 被引用次数: 78 - [相关文章](#)

2005年4月1日 - [Duodenal intraepithelial lymphocytosis](#) with normal villous ... Lorenzo Memeo, Jeffrey Jhang, Hanina Hibshoosh, Peter H Green, ... infecting up to 30% of individuals in developed countries and >50% of the ... characterized as either ulcer-associated or [nonspecific](#). ... Earth sciences - [Evolution](#) & Ecology ...

Increasing duodenal intraepithelial lymphocytosis found at ...

www.ncbi.nlm.nih.gov/pubmed/24565068 - [翻译此页](#)

作者: E Schmidt - 2014 - 被引用次数: 5 - [相关文章](#)

2014年2月22日 - Schmidt E(1), Smyrk TC(2), Boswell CL(3), Enders FT(4), Oxentenko AS(5). ... BACKGROUND: Increased [intraepithelial lymphocytosis](#) (IEL) with preserved villous architecture is a common yet [nonspecific](#) finding on [duodenal](#) biopsies. ... of isolated IEL on [duodenal](#) biopsy over time and determine whether ...

缺少字词: [evolution](#)

An approach to duodenal biopsies

www.ncbi.nlm.nih.gov > ... > [PubMed Central \(PMC\)](#) - [翻译此页](#)

作者: S Serra - 2006 - 被引用次数: 48 - [相关文章](#)

2006年5月10日 - The [development](#) of the intestinal tract begins as a tube that joins the ... [Duodenal](#) biopsy specimens can be taken in [two](#) ways: by suction or by ... be reported as borderline [intraepithelial lymphocytosis](#) and that ≥30 ... needs to be assessed by long-term [follow-up](#) or gluten challenge. ... Age >2 [years](#).

[PDF] Biopsy Approach to Duodenal Mucosal Biopsies Dr Golds...

www2.massgeneral.org/.../Postgraduate%20GI%20Course%20... - [翻译此页](#)

作者: HMSP Course - [相关文章](#)

Web

Images

Videos

News

Shopping

Maps

Books

About 64,600 results

Any time

Past hour

Past 24 hours

Past week

Past month

Past year

All results

Verbatim

[Modern Pathology - Duodenal intraepithelial lymphocytosis with ...](#)

www.nature.com/articles/3800404 ▾

Apr 1, 2005 ... Our study highlights frequent **duodenal intraepithelial lymphocytosis** in individuals ... pathogens, infecting **up** to 30% of individuals in developed countries and >50% of ... characterized as either ulcer-associated or **nonspecific**. All **duodenal** biopsies had at least **two** well-oriented pieces (range **2–9**, mean ...

[An approach to duodenal biopsies](#)

www.ncbi.nlm.nih.gov/pmc/articles/PMC1860495/ ▾

May 10, 2006 ... The **development** of the intestinal tract begins as a tube that joins the **Duodenal** biopsy specimens can be taken in **two** ways: by suction or by be reported as borderline **intraepithelial lymphocytosis** and that ≥ 30 ... needs to be assessed by long-term **follow-up** or gluten challenge. Age **>2 years**.

[Increasing duodenal intraepithelial lymphocytosis found at upper ...](#)

www.ncbi.nlm.nih.gov/pubmed/24565068 ▾

Feb 22, 2014 ... Shmidt E(1), Smyrk TC(2), Boswell CL(3), Enders FT(4), Oxentenko AS(5). ... BACKGROUND: Increased **intraepithelial lymphocytosis** (IEL) with preserved villous architecture is a common yet **nonspecific** finding on **duodenal** biopsies. ... of isolated IEL on **duodenal** biopsy **over** time and determine whether ...

[\[PDF\] Biopsy Approach to Duodenal Mucosal Biopsies - Massachusetts ...](#)

www2.massgeneral.org/.../Postgraduate%20GI%20Course%20Duodenal%20Disease%20Dr%20... ▾

duodenal villi can be reduced significantly **over** normally present lymphoid nodules ... **years**. Celiac disease always occurs in persons who have inherited HLA ... positive **intraepithelial lymphocytosis** characteristic of celiac disease remains unknown results from **follow-up** of a large-scale population survey of antibodies to ...

[Diagnosis and Management of Celiac Disease | American College ...](#)