

5,650 Results Any time ▾

MRI characterization of focal liver lesions in non ...<https://insightsimaging.springeropen.com/articles/10.1186/s13244-020-00894-3> ▾

Sep 22, 2020 To evaluate the added value of the hepatobiliary (HFB) phase in gadoxetic acid-enhanced magnetic resonance imaging (MRI) in characterizing newly discovered indeterminate focal liver lesions in non-cirrhotic patients. One-hundred and twenty-five non-cirrhotic patients (median age, 46 years; range, 20–85 years; 100 females) underwent gadoxetic acid-enhanced MRI, including the ...

Author: Bardia Moosavi, Anuradha S. Shenoy...

Publish Year: 2020

Estimated Reading Time: 12 mins

Search Tools

[Turn off Hover Translation \(关闭翻译\)](#)**Nonhypervascular Hypointense Nodules at Gadoxetic Acid ...**<https://pubs.rsna.org/doi/10.1148/radiol.15141350>

< Introduction Materials and Methods Results Discussion At >

The recently introduced hepatocyte-specific contrast agent gadoxetic acid (Primovist, Bayer Healthcare, Berlin, Germany) has opened new horizons for liver magnetic resonance (MR) imaging, with promising results for the detection and characterization of **hepatocellular carcinoma (HCC)**. This dual-acting imaging agent provides hemodynamic information during early dynamic phases and enhances HCC detection in the hepatobiliary phase (1–3). Results of extensive studies have shown that most HCCs at...

[See more on pubs.rsna.org](#)

Cited by: 62


Author: Jiyoung Hwang, Young Kon Kim, Woo Ky...

Publish Year: 2015

03-Aug-2021 06:11PM

3262 words matched • 29 sources

FAQ

 iThenticate®

64821_Auto_Edited.docx

Quotes Excluded
Bibliography Excluded

16%
Matched

Name of Journal: *World Journal of Hepatology*

Manuscript NO: 64821

Manuscript Type: MINIREVIEWS

Indeterminate liver lesions on Gadoteric acid-enhanced magnetic resonance imaging of the liver: case-based radiologic-pathologic review

Indeterminate liver lesions radiological-pathological correlation.

Jurate Noreikaite, Dekan Albasha, Vijay Chidambaram, Ankur Arora, Ashok Katti

Match Overview

1

Internet 48 words
crawled on 26-Dec-2019
www.gonline.org

1%

2

Crossref 28 words
George Miller, Lawrence H. Schwartz, Michael D'Angelica "The Use of Imaging in the Diagnosis and Staging of Hepa ...

1%

3

Crossref 28 words
Pei-qiang Cai, Yao-pan Wu, Chuan-miao Xie, Wei-dong Zhang, Hai Han, Pei-hong Wu. "Hepatic angiosarcoma: C ...

1%

4

Internet 27 words
crawled on 27-Aug-2019
www.karger.com

1%

5

Internet 27 words
crawled on 16-Nov-2020
www.researchgate.net

1%

6

Internet 26 words
crawled on 13-Mar-2016
tpp.uah.edu

1%

7

Crossref 25 words
Kuma Ozaki, Hiroshi Ikeda, Yasuharu Kozaki, Kazuya Maeda, Shohhei Higuchi, Nobuyuki Kozaka, Hirohiko Kimura, Toshi

1%

8

Crossref 22 words
Hiroki Kanno, Yuchiro Maruyama, Toshihiro Sato, Toru Hira

1%

1

2

PAGE 1 OF 12

1

2

1

2

Text-Only Report

国内版

国际版

Indeterminate liver lesions on gadoxetic acid-enhanced magnetic r



ALL

IMAGES

VIDEOS

6,140 Results

Any time ▾

MRI characterization of focal liver lesions in non ...

<https://insightsimaging.springeropen.com/articles/10.1186/s13244-020-00894-3> ▾

Sep 22, 2020 · To evaluate the added value of the hepatobiliary (HPB) phase in gadoxetic acid-enhanced magnetic resonance imaging (MRI) in characterizing newly discovered indeterminate focal liver lesions in non-cirrhotic patients. One-hundred and twenty-five non-cirrhotic patients (median age, 46 years; range, 20–85 years; 100 females) underwent gadoxetic acid-enhanced MRI, including the ...

Author: Bardia Moosavi, Anuradha S. Shenoy-... Publish Year: 2020

Estimated Reading Time: 12 mins

Approach to the Solitary Liver Lesion: Imaging and When to ...

<https://www.sciencedirect.com/science/article/pii/S0846537115000893>

May 01, 2016 · Magnetic resonance imaging of the lesion (arrows) with gadoxetic acid demonstrating T1 isointensity to liver with a central hypointense nodule on unenhanced images (A), heterogenous mild arterial enhancement (B), portal venous washout (C), and hypointensity to liver on hepatocyte phase (D).

Cited by: 5

Author: Emily H.T. Pang, Alison C. Harris, Silvia ...

Publish Year: 2016

PEOPLE ALSO ASK

What kind of imaging is used for liver lesions? ▾

How is positron emission tomography used to diagnose liver disease? ▾

How big is a lesion on the liver? ▾

Feedback

Indeterminate 1-2-cm Nodules Found on Hepatocellular ...

<https://www.researchgate.net/publication/51774073...>

Purpose: To investigate the patterns of imaging appearance of hepatocellular carcinoma (HCC) on gadoxetic acid-enhanced magnetic resonance (MR) images and to assess their potential value as ...

Estimated Reading Time: 10 mins