

[Ulcerative colitis with mucosal lesions in duodenum: Two ...](#)

<https://pubmed.ncbi.nlm.nih.gov/30946342>

Rationale: Ulcerative colitis (UC) is a chronic, nonspecific, inflammatory disease of the colon. Colorectal is the main target organ of UC, while other digestive tract involvement is rare. This report describes 2 rare cases of duodenal mucosa lesions in patients with UC after total colectomy.

[Evaluation of the Upper Gastrointestinal Tract in ...](#)

<https://pubmed.ncbi.nlm.nih.gov/29674758>

Forty-two patients (19.4%) had upper gastrointestinal lesions, including multiple erosions (n=18), bamboo joint-like appearance (n=17), mucosa with white spots (n=4), friable mucosa (n=2), ulcer (n=1), and purulent deposits within the mucosa (n=1) in the stomach and/or duodenum.

Cited by: 3

Author: Ryo Kato, Masaya Iwamuro, Sakiko Hira...

Publish Year: 2018

PEOPLE ALSO ASK

What is ulcerative colitis (UC)?



How to diagnose ulcerative proctitis?



What are the endoscopic features of the colon?



What are erosions and ulcers?



Feedback

[Ulcerative colitis with mucosal lesions in duodenum: Two ...](#)

<https://journals.lww.com/md-journal/Fulltext/2019/...>

UC often presents with abdominal pain, diarrhea, and hematochezia. It is important to exclude infection as the etiology of UC. Colorectal is the main target organ of UC, while other digestive tract involvement is rare. In recent years, it has been reported that patients with UC may also develop complications with gastric and duodenal mucosal lesions. These complications may be associated with colorectal resection ...

(PDF) [Ulcerative colitis with mucosal lesions in duodenum ...](#)

ALL IMAGES VIDEOS

993,000 Results Any time ▾

Evaluation of the Upper Gastrointestinal Tract in ...

<https://pubmed.ncbi.nlm.nih.gov/29674758>

Forty-two patients (19.4%) had upper gastrointestinal lesions, including multiple erosions (n=18), bamboo joint-like appearance (n=17), mucosa with white spots (n=4), friable mucosa (n=2), ulcer (n=1), and purulent deposits within the mucosa (n=1) in the stomach and/or duodenum.

Cited by: 4 Author: Ryo Kato, Masaya Iwamura, Sakiko Hiraoka...
Publish Year: 2018

Prospective evaluation of upper gastrointestinal mucosal ...

<https://www.ncbi.nlm.nih.gov/pubmed/7815240>

Granulomas were detected in 10 patients in the upper gastrointestinal tract: one esophageal, eight gastric, and three duodenal. Of the ulcerative colitis patients, seven had esophagitis, one had esophageal ulcer, 17 had nonspecific gastritis, two had gastric ulcers, two had duodenal ulcers, and five had H. pylori infection; altogether 30 (75%) yielded pathological findings.

Cited by: 211 Author: T. Ruuska, P. Vaajalahti, P. Arajarvi, M. Maki...
Publish Year: 1994

Evaluation of the Upper Gastrointestinal Tract in ...

<https://www.ncbi.nlm.nih.gov/pubmed/29674758>

Forty-two patients (19.4%) had upper gastrointestinal lesions, including multiple erosions (n=18), bamboo joint-like appearance (n=17), mucosa with white spots (n=4), friable mucosa (n=2), ulcer (n=1), and purulent deposits within the mucosa (n=1) in the stomach and/or duodenum. Compared to the patients without upper gastrointestinal lesions, those with upper gastrointestinal lesions showed significantly ...

Cited by: 4 Author: Ryo Kato, Masaya Iwamura, Sakiko Hiraoka...
Publish Year: 2018

Histopathology of duodenal mucosal lesions in pediatric ...

<https://pubmed.ncbi.nlm.nih.gov/25207874>

Histopathologic lesions of the upper gastrointestinal tract (UGT) are common in inflammatory bowel disease (IBD) patients. Pediatric patients have a higher incidence of IBD-associated gastritis and duodenitis than do adults.

Cited by: 32 Author: Steven Hardee, Arik Alper, Dinesh S. Pasha...
Publish Year: 2014

Upper gastrointestinal tract lesions in patients with ...

<https://www.ncbi.nlm.nih.gov/pubmed/12856568>

The endoscopic examination revealed erosions of the abdominal esophagus in 13.8% cases, erosive-ulcerative alterations in the stomach and duodenum in 27.5% of NUC patients. The most marked changes occurred in patients with severe NUC. Alterations in the upper GIT were not related with either Hp or the acidity.

Cited by: 2 Author: Gidaiatov Aa, Alieva Sa...
Publish Year: 2003

Ulcerative colitis with mucosal lesions in duodenum: Two ...

<https://journals.lww.com/md-journal/Fulltext/2019/...>

UC often presents with abdominal pain, diarrhea, and hematochezia. It is important to exclude infection as the etiology of UC. Colorectal is the main target organ of UC, while other digestive tract involvement is rare. In recent years, it has been reported that patients with UC may also develop complications with gastric and duodenal mucosal lesions. These complications may be associated with colorectal ...


Search Tools

Turn off Hover Translation (X)

Add the Olive with Bing extension >

Make a difference for a nonprofit, simply by searching on Bing

MAYBE LATER YES



Name of Journal: *World Journal of Gastroenterology*

Manuscript NO: 63662

Manuscript Type: MINIREVIEWS

Mucosal lesions of the upper gastrointestinal tract in patients with ulcerative colitis:
A review

Sun Y *et al.* Mucosal lesions of UGI in UC

Yan Sun, Zhe Zhang, Chang-Qing Zheng, Li-Xuan Sang

Abstract

Ulcerative colitis (UC) is a chronic, nonspecific, relapsing inflammatory bowel disease. The colorectum is considered the chief target organ of UC, whereas upper gastrointestinal (UGI) tract manifestations are infrequent. Recently, emerging evidence has suggested that UC presents complications in esophageal, stomachic and duodenal mucosal injuries. However, UC-related UGI tract manifestations are varied and are

Match Overview

1	Crossref 85 words Jingmei Lin, Barbara J. McKenna, Henry D. Appelman. "Mor- phologic Findings in Upper Gastrointestinal Biopsies of Pa...	2%
2	Crossref 79 words Serena, Alessandra, Paolo Biancheri, Philip Allan, Oino R. Corazza, and Antonio Di Sabatino. "Immune and adaptive in...	2%
3	Crossref Posted Content 15 words Shihui Sun, Hongqing Gu, Lei Cao, Qi Chen et al. "Charact... ization and structural basis of a lethal mouse-adapted SARS	<1%
4	Internet 13 words crawled on 21-Dec-2017 link.springer.com	<1%
5	Crossref 12 words Xin Cao, Kang Li, Xue-Li Xu, Karen M von Deneen, Guo-Hu a Geng, Xue-Li Chen. "Development of tomographic reconstr	<1%
6	Crossref 12 words Giancarlo Maria Lumbroso. "A Case Report of Long-term ... emission of Ulcerative Colitis After Lymphocyto-plasmapher	<1%
7	Internet 10 words crawled on 08-Mar-2019 www.abjournals.org	<1%

Mucosal lesions of the upper gastrointestinal tract in patients with u



ALL IMAGES VIDEOS

781,000 Results Any time ▾

Prospective evaluation of upper gastrointestinal mucosal ...

<https://www.ncbi.nlm.nih.gov/pubmed/7815240>

Granulomas were detected in 10 patients in the upper gastrointestinal tract: one esophageal, eight gastric, and three duodenal. Of the ulcerative colitis patients, seven had esophagitis, one had esophageal ulcer, 17 had nonspecific gastritis, two had gastric ulcers, two had duodenal ulcers, and five had H. pylori infection; altogether 30 (75%) yielded pathological findings.

Cited by: 210 Author: T. Ruuska, P. Vaajalahti, P. Arajärvi, M. M...
Publish Year: 1994

Evaluation of the Upper Gastrointestinal Tract in ...

<https://pubmed.ncbi.nlm.nih.gov/29674758>

Forty-two patients (19.4%) had upper gastrointestinal lesions, including multiple erosions (n=18), bamboo joint-like appearance (n=17), mucosa with white spots (n=4), friable mucosa (n=2), ulcer (n=1), and purulent deposits within the mucosa (n=1) in the stomach and/or duodenum.

Cited by: 4 Author: Ryo Kato, Masaya Iwamuro, Sakiko Hira...
Publish Year: 2018

PEOPLE ALSO ASK

How much of an oral ulcer is painless? ▾

Is there a prevalence of oral ulcers in SLE? ▾

How many rheumatoid arthritis patients have oral ulceration? ▾

What kind of lesions are found on the hard palate? ▾

Feedback

Evaluation of the Upper Gastrointestinal Tract in ...

<https://www.ncbi.nlm.nih.gov/pubmed/29674758>

Forty-two patients (19.4%) had upper gastrointestinal lesions, including multiple erosions (n=18), bamboo joint-like appearance (n=17), mucosa with white spots (n=4), friable mucosa (n=2), ulcer (n=1), and purulent deposits within the mucosa (n=1) in the stomach and/or duodenum. Compared to the patients