

Match Overview

[x] [y]

1	CrossCheck (7 words) Shimizu, Ichiro, Tsubota, Kazuo, Yoshida, Toshiyuki, Iida, Shiro, Iwano, Takanori, Nakamura, Kazuo, Ito, Jun-ichi, Ito, Jun-ichi	1%
2	Internet (17 words) Shimizu, Ichiro, Tsubota, Kazuo, Yoshida, Toshiyuki, Iida, Shiro, Iwano, Takanori, Nakamura, Kazuo, Ito, Jun-ichi, Ito, Jun-ichi	1%
3	Internet (11 words) Shimizu, Ichiro, Tsubota, Kazuo, Yoshida, Toshiyuki, Iida, Shiro, Iwano, Takanori, Nakamura, Kazuo, Ito, Jun-ichi, Ito, Jun-ichi	<1%
4	CrossCheck (11 words) Shimizu, Ichiro, Tsubota, Kazuo, Yoshida, Toshiyuki, Iida, Shiro, Iwano, Takanori, Nakamura, Kazuo, Ito, Jun-ichi, Ito, Jun-ichi	<1%
5	Publication (11 words) Shimizu, Ichiro, Tsubota, Kazuo, Yoshida, Toshiyuki, Iida, Shiro, Iwano, Takanori, Nakamura, Kazuo, Ito, Jun-ichi, Ito, Jun-ichi	<1%
6	CrossCheck (11 words) Shimizu, Ichiro, Tsubota, Kazuo, Yoshida, Toshiyuki, Iida, Shiro, Iwano, Takanori, Nakamura, Kazuo, Ito, Jun-ichi, Ito, Jun-ichi	<1%
7	Internet (11 words) Shimizu, Ichiro, Tsubota, Kazuo, Yoshida, Toshiyuki, Iida, Shiro, Iwano, Takanori, Nakamura, Kazuo, Ito, Jun-ichi, Ito, Jun-ichi	<1%

Name of journal: World Journal of Gastroenterology

ESPS Manuscript NO: 14842

Column: ORIGINAL ARTICLE

Randomized Clinical Trial

Effectiveness of probiotic therapy for the prevention of relapse in patients with inactive ulcerative colitis

Yasushi Yoshimatsu, Akitsuo Yamada, Kyuichi Furukawa, Koji Sano, Atsuko

Chamura, Kenjiro Nakamura, Hiroshi Aoki, Yukiko Tsuda, Nobuo Hosoe,

Nobuo Takada, Yasuo Suzuki



Effectiveness of probiotic therapy for the prevention of relapse in patients with



学术搜索

找到约 1,780 条结果 (用时0.11秒)

我的著作引用情况

文章

我的图书馆

时间不限

2015以来

2014以来

2011以来

自定义范围...

按相关性排序

按日期排序

搜索所有网页

中文网页

简体中文网页

☒ 包括专利☐ 包含引用

创建快讯

小提示: 只搜索中文(简体)结果, 可在 学术搜索设置 指定搜索语言

Double-blind comparison of an oral Escherichia coli preparation and mesalazine in maintaining remission of ulcerative colitis

W Kruis, E Schütz, P Fric, B Fixa... - Alimentary ..., 1997 - Wiley Online Library

... Conclusions: From the results of this preliminary study, probiotic treatment appears to offer another option for ... pathogenic E. coli is of experimental character and, as far as effective standard therapy is ... 06: K5: H1) showed in this short-term pilot study similar efficacy in maintaining ...

被引用次数: 715 相关文章 所有 5 个版本 引用 保存

wiley.com 中的 [PDF]

Therapeutic manipulation of the enteric microflora in inflammatory bowel diseases: antibiotics, probiotics, and prebiotics

RB Sartor - Gastroenterology, 2004 - Elsevier

... isolated ileal disease), perianal fistulae and pouchitis, whereas selected probiotic preparations prevent ... considerable logic, unfortunately, there is quite limited documentation of efficacy of these the rationale, available evidence for the use of antibiotics, probiotics, and prebiotics ...

被引用次数: 805 相关文章 所有 12 个版本 引用 保存

hmphanmi.cn 中的 [PDF]

Non-pathogenic Escherichia coli versus mesalazine for the treatment of ulcerative colitis: a randomised trial

BJ Rembacken, AM Snelling, PM Hawkey... - The Lancet, 1999 - Elsevier

... database, Sutherland and colleagues found that 5-aminosalicylic acid was significantly more effective than placebo but there was no difference in efficacy between different ... The manipulation of the colonic bacteria with antibiotics and probiotic agents may prove more ...

被引用次数: 1067 相关文章 所有 8 个版本 引用 保存

nutrition.org 中的 [HTML]

Protection from gastrointestinal diseases with the use of probiotics

PR Marteau, M de Vrese, CJ Cellier... - The American journal ..., 2001 - Am Soc Nutrition

... Two studies (88, 89) compared the efficacy of an oral E. coli preparation [E. coli strain Nissle (Mutaflor, Ardeypharm GmbH, Herdecke, Germany)] and ... The second study included 116 patients and also showed that the probiotic preparation was as effective as mesalazine in ...

被引用次数: 582 相关文章 所有 11 个版本 引用 保存

wignot.com 中的 [HTML]

[网页](#)[新闻](#)[购物](#)[图片](#)[视频](#)[更多 ▾](#)[搜索工具](#)

找到约 763,000 条结果 (用时 0.69 秒)

Google 学术 : Effectiveness of probiotic therapy for the prevention of relapse in patients with inactive ulcerative colitis

... in maintaining remission of ulcerative colitis - Kruis - 被引用次数 : 716

Therapeutic manipulation of the enteric microflora in ... - Sartor - 被引用次数 : 806

... versus mesalazine for the treatment of ulcerative colitis ... - Rembacken - 被引用次数 : 1068

Intestinal microbiota, probiotics and prebiotics in ...

www.ncbi.nlm.nih.gov > ... > Literature > PubMed Central (PMC) - 翻译此页

作者 : TK Trop - 2014 - 被引用次数 : 2 - 相关文章

2014年9月7日 - For treatment of active ulcerative colitis, as well as for maintenance ...

A pathogenic role of luminal constituents is suggested by the prevention and treatment of Thus, generalizations concerning the efficacy of a whole species or even

Relapse was reported in 40.1% of patients treated with probiotics ...

Double-blind comparison of an oral Escherichia coli ...

www.ncbi.nlm.nih.gov/pubmed/9354192 ▾ 翻译此页

作者 : W Kruis - 1997 - 被引用次数 : 717 - 相关文章

METHODS: A total of 120 patients with inactive ulcerative colitis were included in ...

K5: H1) for 12 weeks with regard to their efficacy in preventing a relapse of the

disease. ... CONCLUSIONS: From the results of this preliminary study, probiotic ...

Maintaining remission of ulcerative colitis with the probiotic ...

www.pubfacts.com/.../Maintaining-remission-of-ulcerative-colit... ▾ 翻译此页

The effectiveness of probiotic treatment further underlines the pathogenetic ... coli

(Nissle 1917) was as effective as mesalazine in preventing relapse of ulcerative colitis.

... A total of 120 patients with inactive ulcerative colitis were included in a ...