

[全部](#)[图片](#)[新闻](#)[视频](#)[购物](#)[更多](#)[设置](#)[工具](#)

找到约 86,500 条结果 (用时 0.73 秒)

Mo1441 Lymph Node Ratio and Para-Aortic Node ... - Gastroenterology

[www.gastrojournal.org/article/S0016-5085\(16\)34165-8/abstract](http://www.gastrojournal.org/article/S0016-5085(16)34165-8/abstract) - 翻译此页

作者: V Beltrame - 2016

IBD · **Pancreatic** Diseases · Viral Hepatitis · AGA · AGA Home · AGA Mobile · Join AGA · Login close.

Email/Username: Password: Remember me. Forgot password ...

SSAT - Lymph Node Ratio and Para-Aortic Node Involvement Are Not ...

meetings.ssat.com/abstracts/2016/Mo1441.cgi ▼ 翻译此页

2016 Annual Meeting: Lymph **Node** Ratio and **Para-Aortic Node** Involvement Are **Not Independent Predictors of Survival After Resection for Pancreatic Cancer**.

Mo1441 Lymph Node Ratio and Para-Aortic Node Involvement Are ...

https://www.researchgate.net/.../301504975_Mo1441_Lymph_Node_Ratio_... - 翻译此页

Mo1441 Lymph **Node** Ratio and **Para-Aortic Node** Involvement Are **Not Independent Predictors of Survival After Resection for Pancreatic Cancer** on ...

Prognostic value of the lymph node ratio after resection of ... - NCBI

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4048080/> - 翻译此页

作者: SG Farid - 2014 - 被引用次数: 4 - 相关文章

2012年11月19日 - A LNR > 0.2, and **not** lymph node status, is an **independent** prognostic ... The American Joint Committee on **Cancer** (AJCC) TNM classification and high rates of LN **metastasis** to **para-aortic nodal** stations in periampullary tumours' **nodes** as **predictors of survival** for **resected pancreatic adenocarcinoma**.



32681-Review-check.docx

Quotes Excluded
Bibliography Excluded

6%
SIMILAR

Name of Journal: *World Journal of Gastroenterology*

ESPS Manuscript NO: 32681

Manuscript Type: ORIGINAL ARTICLE

Para-aortic node involvement is not an independent predictor of survival after resection for pancreatic cancer

Retrospective Cohort Study

Sperti C *et al.* Pancreatic cancer and para-aortic nodes

Cosimo Sperti, Mario Gruppo, Stella Blandamura, Michele Valmasoni, Gioia Pozza, Nicola Passuello, Valentina Beltrame, Lucia Moletta

Abstract

AIM

To analyze the importance of para-aortic node status in a series of patients who underwent pancreaticoduodenectomy (PD) in a single Institution.

Match Overview

1	Crossref 69 words Pedrazzoli, Sergio. "Extent of lymphadenectomy to associate with pancreaticoduodenectomy in patients with pa ...	2%
2	Crossref 24 words van der Gaag, N.A.. "Chylous Ascites after Pancreaticoduodenectomy: Introduction of a Grading System", Journal	1%
3	Crossref 19 words Nappo, Gennaro, Domenico Borzomati, Giuseppe Perro ne, Sergio Valeri, Michela Amato, Tommasangelo Petitti,	1%
4	Crossref 18 words Tol, Johanna A.M.G., Dirk J. Gouma, Claudio Bassi, Christos Derveniz, Marco Montorsi, Mustapha Adham, Ake A	1%
5	Crossref 13 words Kono, Toshiaki, Yoshiaki Murakami, Naru Kondo, Kenichiro Uemura, Yasushi Hashimoto, Naoya Nakagawa, Ka	<1%
6	Crossref 13 words "posters", <i>Annals of Oncology</i> , 06/01/2009	<1%
7	Internet 12 words crawled on 15-Mar-2012 www.biomedcentral.com	<1%

[全部](#)[图片](#)[新闻](#)[视频](#)[购物](#)[更多](#)[设置](#)[工具](#)

找到约 203,000 条结果 (用时 0.66 秒)

Mo1441 Lymph Node Ratio and Para-Aortic Node ... - Gastroenterology

[www.gastrojournal.org/article/S0016-5085\(16\)34165-8/abstract](http://www.gastrojournal.org/article/S0016-5085(16)34165-8/abstract) - 翻译此页

作者: V Beltrame - 2016

IBD · **Pancreatic** Diseases · Viral Hepatitis · AGA · AGA Home · AGA Mobile · Join AGA · Login close.

Email/Username: Password: Remember me. Forgot password ...

SSAT - Lymph Node Ratio and Para-Aortic Node Involvement Are Not ...

meetings.ssat.com/abstracts/2016/Mo1441.cgi ▼ 翻译此页

2016 Annual Meeting: Lymph **Node** Ratio and **Para-Aortic Node** Involvement Are **Not Independent Predictors of Survival After Resection for Pancreatic Cancer**.

The Role of Paraaortic Lymph Node Involvement on Early Recurrence ...

[www.journalacs.org/article/S1072-7515\(06\)00824-6/abstract](http://www.journalacs.org/article/S1072-7515(06)00824-6/abstract) ▼ 翻译此页

作者: K Shimada - 2006 - 被引用次数: 66 - 相关文章

Department of Hepatobiliary and **Pancreatic Surgery**, National **Cancer** Center ... **after** operation and those whose recurrence did **not** develop within 1 year. ... **involvement** and positive washing cytology were **independent predictors** of early recurrence. The median **survival** time in 26 patients with **paraaortic nodal involvement** ...

Mo1441 Lymph Node Ratio and Para-Aortic Node Involvement Are ...

https://www.researchgate.net/.../301504975_Mo1441_Lymph_Node_Ratio_... - 翻译此页

Mo1441 Lymph **Node** Ratio and **Para-Aortic Node** Involvement Are **Not Independent Predictors of Survival After Resection for Pancreatic Cancer** on ...

[All](#)[Images](#)[News](#)[Videos](#)[Shopping](#)[More](#)[Settings](#)[Tools](#)

About 257,000 results (0.78 seconds)

SSAT - Lymph Node Ratio and Para-Aortic Node Involvement Are Not ...

meetings.ssat.com/abstracts/2016/Mo1441.cgi ▼

2016 Annual Meeting: Lymph Node Ratio and Para-Aortic Node Involvement Are Not Independent Predictors of Survival After Resection for Pancreatic Cancer.

Mo1441 Lymph Node Ratio and Para-Aortic Node ... - Gastroenterology

[www.gastrojournal.org/article/S0016-5085\(16\)34165-8/abstract](http://www.gastrojournal.org/article/S0016-5085(16)34165-8/abstract)

by V Beltrame - 2016

IBD · **Pancreatic** Diseases · Viral Hepatitis. AGA. AGA Home · AGA Mobile · Join AGA. Login close.

Email/Username: Password: Remember me. Forgot password ...

The Role of Paraaortic Lymph Node Involvement on Early Recurrence ...

<https://www.ncbi.nlm.nih.gov/labs/articles/16931307/> ▼

21 Jul 2006 - The median **survival** time in 26 patients with **paraaortic nodal** ... Ratio Is the Strongest Prognostic Factor **After Resection of Pancreatic Cancer**.

Mo1441 Lymph Node Ratio and Para-Aortic Node Involvement Are ...

[metcorner.gastro.org/article/S0016-5085\(16\)34165-8/abstract](https://metcorner.gastro.org/article/S0016-5085(16)34165-8/abstract) ▼

Mo1441 Lymph Node Ratio and Para-Aortic Node Involvement Are Not Independent Predictors of Survival After Resection for Pancreatic Cancer. Valentina ...

Mo1441 Lymph Node Ratio and Para-Aortic Node Involvement Are ...

https://www.researchgate.net/.../301504975_Mo1441_Lymph_Node_Ratio_and_Para-A...

Mo1441 Lymph Node Ratio and Para-Aortic Node Involvement Are Not Independent Predictors of Survival After Resection for Pancreatic Cancer on ...

Postoperative prognosis of pancreatic cancer with para-aortic lymph ...

https://www.researchgate.net/.../267745206_Postoperative_prognosis_of_pancreatic_can...

... cancer with **para-aortic** lymph node metastasis: a multicenter study on 822 patients ... static PAIN may survive for longer than expected after ... Keywords **Pancreatic cancer** **Para-aortic**