

# World Journal of *Clinical Cases*

*World J Clin Cases* 2023 September 26; 11(27): 6318-6669



**MINIREVIEWS**

- 6318 Characteristics of amino acid metabolism in colorectal cancer  
*Xu F, Jiang HL, Feng WW, Fu C, Zhou JC*

**ORIGINAL ARTICLE****Clinical and Translational Research**

- 6327 Exploring the pharmacological mechanism of Wuzhuyu decoction on hepatocellular carcinoma using network pharmacology  
*Ouyang JY, Lin WJ, Dong JM, Yang Y, Yang HK, Zhou ZL, Wang RQ*
- 6344 Identification of potential diagnostic and prognostic biomarkers for breast cancer based on gene expression omnibus  
*Zhang X, Mi ZH*

**Retrospective Cohort Study**

- 6363 Treatment of proximal humeral fractures accompanied by medial calcar fractures using fibular autografts: A retrospective, comparative cohort study  
*Liu N, Wang BG, Zhang LF*

**Retrospective Study**

- 6374 Effectiveness of out-fracture of the inferior turbinate with reduction nasal bone fracture  
*Kim SY, Nam HJ, Byeon JY, Choi HJ*
- 6383 Prognostic model of hepatocellular carcinoma based on cancer grade  
*Zhang GX, Ding XS, Wang YL*
- 6398 Oncologic efficacy of gonadotropin-releasing hormone agonist in hormone receptor-positive very young breast cancer patients treated with neoadjuvant chemotherapy  
*Choi HJ, Lee JH, Jung CS, Ryu JM, Chae BJ, Lee SK, Yu JH, Kim SW, Nam SJ, Lee JE, Jung YJ, Kim HY*
- 6407 Correlation analysis of serum thyroglobulin, thyroid-stimulating hormone levels, and thyroid-cancer risk in thyroid nodule surgery  
*Shuai JH, Leng ZF, Wang P, Ji YC*
- 6415 Closed thoracic drainage in elderly patients with chronic obstructive pulmonary disease complicated with spontaneous pneumothorax: A retrospective study  
*Wang W, Zhu DN, Shao SS, Bao J*

**Observational Study**

- 6424 *Helicobacter pylori* eradication treatment for primary gastric diffuse large B-cell lymphoma: A single-center analysis  
*Saito M, Mori A, Kajikawa S, Yokoyama E, Kanaya M, Izumiyama K, Morioka M, Kondo T, Tanei ZI, Shimizu A*

**Prospective Study**

- 6431** Effect of polyene phosphatidylcholine/ursodeoxycholic acid/ademetionine on pregnancy outcomes in intrahepatic cholestasis  
*Dong XR, Chen QQ, Xue ML, Wang L, Wu Q, Luo TF*

**SYSTEMATIC REVIEWS**

- 6440** Maternal diaphragmatic hernia in pregnancy: A systematic review with a treatment algorithm  
*Augustin G, Kovač D, Karadjole VS, Zajec V, Herman M, Hrbač P*

**META-ANALYSIS**

- 6455** Laparoscopic vs open radical resection in management of gallbladder carcinoma: A systematic review and meta-analysis  
*He S, Yu TN, Cao JS, Zhou XY, Chen ZH, Jiang WB, Cai LX, Liang X*

**CASE REPORT**

- 6476** Acute acquired concomitant esotropia with congenital paralytic strabismus: A case report  
*Zhang MD, Liu XY, Sun K, Qi SN, Xu CL*
- 6483** Tumor recurrence after pathological complete response in locally advanced gastric cancer after neoadjuvant therapy: Two case reports  
*Xing Y, Zhang ZL, Ding ZY, Song WL, Li T*
- 6491** Acute peritonitis secondary to post-traumatic appendicitis: A case report and literature review  
*Habachi G, Aziza B, Ben-Ammar S, Maherzi O, Houas Y, Kerkeni Y, Sahli S, Jouini R*
- 6498** Fournier's gangrene after insertion of thermo-expandable prostatic stent for benign prostatic hyperplasia: A case report  
*Jung HC, Kim YU*
- 6505** Methyl-CpG-Binding protein 2 duplication syndrome in a Chinese patient: A case report and review of the literature  
*Xing XH, Takam R, Bao XY, Ba-alwi NA, Ji H*
- 6515** Blood purification for treatment of non-liquefied multiple liver abscesses and improvement of T-cell function: A case report  
*Tang ZQ, Zhao DP, Dong AJ, Li HB*
- 6523** Eosinophilic granulomatosis with polyangiitis, asthma as the first symptom, and subsequent Loeffler endocarditis: A case report  
*He JL, Liu XY, Zhang Y, Niu L, Li XL, Xie XY, Kang YT, Yang LQ, Cai ZY, Long H, Ye GF, Zou JX*
- 6531** Left atrium veno-arterial extra corporeal membrane oxygenation as temporary mechanical support for cardiogenic shock: A case report  
*Lamastra R, Abbott DM, Degani A, Pellegrini C, Veronesi R, Pelenghi S, Dezza C, Gazzaniga G, Belliato M*

- 6537** Successful treatment of eyebrow intradermal nevi by shearing combined with electrocautery and curettage: Two case reports  
*Liu C, Liang JL, Yu JL, Hu Q, Li CX*
- 6543** Amniotic membrane mesenchymal stromal cell-derived secretome in the treatment of acute ischemic stroke: A case report  
*Lin FH, Yang YX, Wang YJ, Subbiah SK, Wu XY*
- 6551** Managing spindle cell sarcoma with surgery and high-intensity focused ultrasound: A case report  
*Zhu YQ, Zhao GC, Zheng CX, Yuan L, Yuan GB*
- 6558** Triplet regimen as a novel modality for advanced unresectable hepatocellular carcinoma: A case report and review of literature  
*Zhao Y, He GS, Li G*
- 6565** Acute diquat poisoning case with multiorgan failure and a literature review: A case report  
*Fan CY, Zhang CG, Zhang PS, Chen Y, He JQ, Yin H, Gong XJ*
- 6573** Fungal corneal ulcer after repair of an overhanging filtering bleb: A case report  
*Zhao J, Xu HT, Yin Y, Li YX, Zheng YJ*
- 6579** Combination therapy with toripalimab and anlotinib in advanced esophageal squamous cell carcinoma: A case report  
*Chen SC, Ma DH, Zhong JJ*
- 6587** Removal of a pulmonary artery foreign body during pulse ablation in a patient with atrial fibrillation: A case report  
*Yan R, Lei XY, Li J, Jia LL, Wang HX*
- 6592** Delayed-onset *micrococcus luteus*-induced postoperative endophthalmitis several months after cataract surgery: A case report  
*Nam KY, Lee HW*
- 6597** Anesthetic management of a pregnant patient with Eisenmenger's syndrome: A case report  
*Zhang Y, Wei TT, Chen G*
- 6603** Recurrence of unilateral angioedema of the tongue: A case report  
*Matsuhisa Y, Kenzaka T, Shimizu H, Hirose H, Gotoh T*
- 6613** Transverse mesocolic hernia with intestinal obstruction as a rare cause of acute abdomen in adults: A case report  
*Zhang C, Guo DF, Lin F, Zhan WF, Lin JY, Lv GF*
- 6618** Compound heterozygous mutations in tripeptidyl peptidase 1 cause rare autosomal recessive spinocerebellar ataxia type 7: A case report  
*Liu RH, Wang XY, Jia YY, Wang XC, Xia M, Nie Q, Guo J, Kong QX*

- 6624** Treatment of posterior interosseous nerve entrapment syndrome with ultrasound-guided hydrodissection: A case report  
*Qin LH, Cao W, Chen FT, Chen QB, Liu XX*
- 6631** Rapidly growing extensive polypoid endometriosis after gonadotropin-releasing hormone agonist discontinuation: A case report  
*Zhang DY, Peng C, Huang Y, Cao JC, Zhou YF*
- 6640** Preserving finger length in a patient with symmetric digital gangrene under local anesthesia: A case report  
*Kim KH, Ko IC, Kim H, Lim SY*
- 6646** Reconstruction of the lower back wound with delayed infection after spinal surgery: A case report  
*Kim D, Lim S, Eo S, Yoon JS*
- 6653** Solitary intraosseous neurofibroma in the mandible mimicking a cystic lesion: A case report and review of literature  
*Zhang Z, Hong X, Wang F, Ye X, Yao YD, Yin Y, Yang HY*
- 6664** Complete response of metastatic *BRAF* V600-mutant anaplastic thyroid cancer following adjuvant dabrafenib and trametinib treatment: A case report  
*Lee SJ, Song SY, Kim MK, Na HG, Bae CH, Kim YD, Choi YS*

**ABOUT COVER**

Editorial Board Member of *World Journal of Clinical Cases*, Alexandru Corlateanu, MD, PhD, Reader (Associate Professor), Department of Respiratory Medicine, Nicolae Testemitanu State University of Medicine and Pharmacy, Chisinau 2001, Moldova. alexandru\_corlateanu@yahoo.com

**AIMS AND SCOPE**

The primary aim of *World Journal of Clinical Cases* (*WJCC*, *World J Clin Cases*) is to provide scholars and readers from various fields of clinical medicine with a platform to publish high-quality clinical research articles and communicate their research findings online.

*WJCC* mainly publishes articles reporting research results and findings obtained in the field of clinical medicine and covering a wide range of topics, including case control studies, retrospective cohort studies, retrospective studies, clinical trials studies, observational studies, prospective studies, randomized controlled trials, randomized clinical trials, systematic reviews, meta-analysis, and case reports.

**INDEXING/ABSTRACTING**

The *WJCC* is now abstracted and indexed in Science Citation Index Expanded (SCIE, also known as SciSearch®), Journal Citation Reports/Science Edition, Current Contents®/Clinical Medicine, PubMed, PubMed Central, Reference Citation Analysis, China National Knowledge Infrastructure, China Science and Technology Journal Database, and Superstar Journals Database. The 2023 Edition of Journal Citation Reports® cites the 2022 impact factor (IF) for *WJCC* as 1.1; IF without journal self cites: 1.1; 5-year IF: 1.3; Journal Citation Indicator: 0.26; Ranking: 133 among 167 journals in medicine, general and internal; and Quartile category: Q4.

**RESPONSIBLE EDITORS FOR THIS ISSUE**

Production Editor: *Ying-Yi Yuan*; Production Department Director: *Xiang Li*; Editorial Office Director: *Jin-Lei Wang*.

**NAME OF JOURNAL**

*World Journal of Clinical Cases*

**ISSN**

ISSN 2307-8960 (online)

**LAUNCH DATE**

April 16, 2013

**FREQUENCY**

Thrice Monthly

**EDITORS-IN-CHIEF**

Bao-Gan Peng, Jerzy Tadeusz Chudek, George Kontogeorgos, Maurizio Serati, Ja Hyeon Ku

**EDITORIAL BOARD MEMBERS**

<https://www.wjgnet.com/2307-8960/editorialboard.htm>

**PUBLICATION DATE**

September 26, 2023

**COPYRIGHT**

© 2023 Baishideng Publishing Group Inc

**INSTRUCTIONS TO AUTHORS**

<https://www.wjgnet.com/bpg/gerinfo/204>

**GUIDELINES FOR ETHICS DOCUMENTS**

<https://www.wjgnet.com/bpg/GerInfo/287>

**GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH**

<https://www.wjgnet.com/bpg/gerinfo/240>

**PUBLICATION ETHICS**

<https://www.wjgnet.com/bpg/GerInfo/288>

**PUBLICATION MISCONDUCT**

<https://www.wjgnet.com/bpg/gerinfo/208>

**ARTICLE PROCESSING CHARGE**

<https://www.wjgnet.com/bpg/gerinfo/242>

**STEPS FOR SUBMITTING MANUSCRIPTS**

<https://www.wjgnet.com/bpg/GerInfo/239>

**ONLINE SUBMISSION**

<https://www.f6publishing.com>

## Reconstruction of the lower back wound with delayed infection after spinal surgery: A case report

DoWon Kim, SooA Lim, SuRak Eo, Jung Soo Yoon

**Specialty type:** Medicine, research and experimental

**Provenance and peer review:** Unsolicited article; Externally peer reviewed.

**Peer-review model:** Single blind

**Peer-review report's scientific quality classification**

Grade A (Excellent): 0  
Grade B (Very good): B  
Grade C (Good): C, C  
Grade D (Fair): 0  
Grade E (Poor): 0

**P-Reviewer:** Collart-Dutilleul PY, France; Long X, China; Mostafavinia A, Iran

**Received:** July 18, 2023

**Peer-review started:** July 18, 2023

**First decision:** August 16, 2023

**Revised:** August 22, 2023

**Accepted:** September 1, 2023

**Article in press:** September 1, 2023

**Published online:** September 26, 2023



**DoWon Kim, SooA Lim, SuRak Eo, Jung Soo Yoon**, Department of Plastic and Reconstructive Surgery, DongGuk University Ilsan Hospital, Goyang 10326, South Korea

**Corresponding author:** Jung Soo Yoon, MD, PhD, Doctor, Professor, Surgeon, Department of Plastic and Reconstructive Surgery, DongGuk University Ilsan Hospital, 814 SikSa-dong, IISanDong-gu, GoYang-si, GyeongGi-do, Goyang 10326, South Korea. [crsboys@naver.com](mailto:crsboys@naver.com)

### Abstract

#### BACKGROUND

Orthopedic surgeries after device implantation are susceptible to infections and may require device removal in the worst cases. For this reason, many efforts are being made to control infections after spinal surgery; however, the number of infection cases is increasing owing to the increasing number of elderly citizens.

#### CASE SUMMARY

A 75-year-old male with a chronic spinal defect due to previous spine surgery underwent reconstruction using a perforator-based island flap. After bursectomy and confirmation that there was no connection with the deep tissue, reconstruction was performed. However, wound disruption occurred with abscess formation on postoperative day 29, which led to an imaging workup revealing delayed deep tissue infection.

#### CONCLUSION

Infection is one of the most common causes of surgical wound dehiscence and is associated with devastating results if not controlled promptly and definitively. Surgeons should always suspect delayed infections when reconstructing chronic soft tissue defects.

**Key Words:** Surgical wound dehiscence; Surgical wound infection; Lumbar spine; Complications; Abscess; Case report

©The Author(s) 2023. Published by Baishideng Publishing Group Inc. All rights reserved.

**Core Tip:** Reasonably, several patients with surgical site dehiscence may require flap surgery after spinal surgery. Although surgeons pay close attention to signs of infection during reconstruction, there can be neglected delayed infection around deep devices. Reconstructive surgeons should be cautious when planning flap surgery after implant surgery.

**Citation:** Kim D, Lim S, Eo S, Yoon JS. Reconstruction of the lower back wound with delayed infection after spinal surgery: A case report. *World J Clin Cases* 2023; 11(27): 6646-6652

**URL:** <https://www.wjgnet.com/2307-8960/full/v11/i27/6646.htm>

**DOI:** <https://dx.doi.org/10.12998/wjcc.v11.i27.6646>

## INTRODUCTION

Surgical wound dehiscence (SWD) is an uncommon but inevitable complication for surgeons. Various factors such as poor general condition, old age, repetitive surgeries, obesity, diabetes, and poor nutrition may contribute to SWD. Among these, surgical site infections (SSIs) are the most frequently reported[1]. A previous study on SWD suggested that longer healing times require more topical antimicrobial dressings or treatment solutions for infections[2].

Recently, the commonly applied management of SSI cases is based on preliminary wound swab culture to detect bacterial infection. Surgical debridement is performed, and intravenous antibiotics are administered. Modern technological developments suggest treating SWD using negative-pressure wound therapy and biosynthetic materials. Vacuum-assisted closure not only cleanses necrotic tissues, but also helps in granulation tissue formation and gradual skin margin approximation[3-5].

Orthopedic surgery, including spinal surgery, is one of the most common causes of SSI. If the infection is related to foreign bodies such as implants, it is associated with high morbidity and may require secondary surgery and delay recovery. Spinal infections with implants are reported to be approximately 8.5%, but < 1% in open surgery without devices[6]. The treatment course is challenging considering that the surgical and medical multidisciplinary approach is essentially based on the type and location of the implant and the patient's comorbidities[7].

By sharing a case of unexpected SSI after reconstruction of a chronic SWD in the lumbar region, the authors want to emphasize a more careful approach to SSI management and the usefulness of perforator flaps for median or paramedian spinal defects. Written informed consent was obtained from the patient.

## CASE PRESENTATION

### Chief complaints

A 75-year-old male was referred to our unit from a local clinic complaining of chronic wounds on his back after undergoing spinal surgery.

### History of present illness

There seemed to be no problem postoperatively for 2 years; however, approximately 2 mo prior, he noticed wound discharge from the back. The neurosurgeon who performed his spine surgery at the local clinic tried debridement and wound revision three times. Radiology at a local clinic revealed no evidence of deep wound infection, and in wound culture, *Pseudomonas* spp. were cultivated. This led to the use of piperacillin/tazobactam; however, the patient developed a severe allergic reaction, and ciprofloxacin was administered instead. The patient was referred to us because of no improvement in the clinical course.

### History of past illness

He had herniation of the nucleus pulposus and had undergone open lumbar-assisted microdiscectomy 3 years previously. At that time, the patient underwent wound revision for SWD. He also underwent anterior lumbar interbody fusion 2 years prior and direct lateral interbody fusion 3 mo later. SWD was observed again in both the second and third surgeries.

### Personal and family history

His body mass index was 27.67 kg/m<sup>2</sup> and he had been taking diabetic medications for approximately 20 years. There were no other remarkable comorbidities.

### Physical examination

On physical examination, approximately 5 cm × 4 cm sized elliptical soft tissue defect was discovered with bursa formation (Figure 1).



DOI: 10.12998/wjcc.v11.i27.6646 Copyright ©The Author(s) 2023.

Figure 1 A 75-year-old man with a chronic wound on the lower back.

### Laboratory examinations

C-reactive protein (CRP) level at the time of admission was 0.50 mg/dL, within the normal range. In addition, no abnormalities were observed in other routine blood or urine analyses.

### Imaging examinations

Simple spine X-ray revealed degenerative spondylosis and further imaging study was not necessary at the time of visit.

---

## FINAL DIAGNOSIS

Soft tissue defects with chronic infections were diagnosed after spinal surgery. After debridement with curettage of the infected granulation tissue, negative-pressure wound therapy was planned for one week, and a perforator-based island flap was scheduled under general anesthesia.


---

## TREATMENT

After a thorough bursectomy, no connection with the deep tissue was confirmed based on gross findings, and reconstruction was performed. Two healthy freestyle perforators of the dorsal branch of the left lumbar artery within 5 cm lateral to the midline were detected using handheld Doppler imaging. The perforators were marked as pivot points and an island flap was designed. The flap was designed to be slightly longer than the defect. Finally, a 4 cm × 9 cm sized perforator-based flap was elevated and rotated toward the wound. Along with defatting, an island-type inset was completed and the authors confirmed the absence of tension on the perforator (Figure 2). Tissue culture during surgery reported *Pseudomonas aeruginosa* growth. Because the patient had a history of allergic reactions to piperacillin-tazobactam, intravenous ciprofloxacin 400 mg every 12 h was administered during an infectious disease consultation. No immediate post-operative complications were observed.


On postoperative day 29, when stitches were removed partially, skin thinning with bulging was observed in the flap's 5 o'clock region. A pus-like discharge was made out of the lesion using a slit incision. A betadine-soaked gauze packing dressing with massive irrigation was applied for 10 d, and wound closure with 4-0 nylon was attempted. Nine days after resuturing, another abscess formation was observed in the 11 o'clock region of the flap (Figure 3A).

Therefore, we rechecked the imaging results. L-spine computed tomography and enhanced L-spine magnetic resonance imaging revealed a suspicious soft tissue phlegmon along the interspinous reconstruction wire (Figure 3B). The spine surgeon strongly suspected deep tissue infection related to the wire and planned the incision and drainage with wire removal. There was no evidence of infection with the interbody devices and screws; therefore, the removal of another instrumentation was not considered. With an incision in the previous surgical scar and the proximal region of the flap, the bilateral pedicle screw head was exposed. The pus gushed out above the interspinous wire from a deeper region of the midline. Liquefactive tissue with an infectious tract was observed around the wire, which made the source of the infection evident. The interspinous wire was successfully removed by undermining the distal region of the wire



DOI: 10.12998/wjcc.v11.i27.6646 Copyright ©The Author(s) 2023.

**Figure 2 Intraoperative photographs.** A: A 4 cm × 9 cm sized perforator-based island flap was designed; B: Island-type inset was tried after defatting; C: Authors confirmed no tension to the perforator before skin closure.



DOI: 10.12998/wjcc.v11.i27.6646 Copyright ©The Author(s) 2023.

**Figure 3 Delayed infection was noticed.** A: Another abscess formation was noticed in the 11 o'clock region of the flap; B: 1.5T enhanced L-spine magnetic resonance imaging T2 imaging study revealed suspicious soft tissue phlegmon (orange arrow) along the interspinous reconstruction wire.

(Figure 4). Meticulous curettage along the infected tract was performed with massive irrigation, along with layer-by-layer closure. Tissue culture during the removal surgery also resulted in *Pseudomonas aeruginosa* growth.

## OUTCOME AND FOLLOW-UP

On postoperative day 18, all sutures were removed after the wire removal, and the patient was discharged without further complications (Figure 5). The Infectious Diseases Department recommended further oral ciprofloxacin 750 mg twice daily for more than 3 mo. At the 5-mo follow-up, the wound had healed well without any complications.

## DISCUSSION

Delayed wound healing and compromised granulation tissue formation result from numerous causes. The host factors include age, tobacco use, comorbidities, and obesity[7]. Mechanical factors include preoperative skin antisepsis, suture breakage, infection, hematoma, seroma, and mechanical stress[1,7,8]. Wound healing requires sufficient oxygen and nutrients from an appropriate diet and vascular supply. Therefore, any factor that impedes them might compromise wound healing and increase the risk of dehiscence (SWD). They deteriorate quality of life, prolong hospital stays, and increase medical costs. Infection (SSI) is one of the most frequent causes of all the factors. Some surgeons consider SSD as an SSI because of its inseparable association[9].



DOI: 10.12998/wjcc.v11.i27.6646 Copyright ©The Author(s) 2023.

Figure 4 Infected interspinous reconstruction wire was removed by the spine surgeon.



DOI: 10.12998/wjcc.v11.i27.6646 Copyright ©The Author(s) 2023.

Figure 5 The patient was discharged without further complications.

Orthopedic implants are increasingly being used by senior citizens and those with moderate-to-severe medical problems[9]. Orthopedic implants are classified into two categories. One is a prosthesis designed to replace joints, and the other is hardware, such as plates, screws, and wires, designed to fix broken bones. Both are highly susceptible to infection, and despite endless efforts to overcome this, infection still occurs in 5% of cases, and 12%-38% recur[10,11]. In addition, the absolute number of infections increases because of the growing number of older adults and the lifelong risk of microbial seeding on implants[10]. If bacterial infections of these implants are not treated thoroughly and immediately, they may develop into fungal infections, in addition to significant morbidity and increased expenditure[12]. Severe cases involving chronic osteomyelitis, spine instability, and spreading can result in septic conditions[13].

*Staphylococcus aureus* and *Staphylococcus epidermidis* are the most commonly reported pathogens associated with orthopedic device-related infections[14]. *Pseudomonas aeruginosa* is known to develop sturdy biofilms that easily lead to chronic infections. It resides in a deeper layer of wounds than *Staphylococcus aureus*, and the infection tends to be more severe, broader, and harder to recover from[15]. This mechanism involves interactions between the host, implant, and the microorganism[10]. During surgery, the bacteria on the patient's skin enter through the incision site and establish colonies on the device[14]. Implants lack microcirculation, which is required for host defense and antibiotic distribution. Biofilms formed by bacterial colonization consist of aggregated microbial cells that continuously produce a self-made polymeric matrix, including host components[2]. It can be observed in chronic wounds and most implant-associated infections. Biofilms defeat antibiotics and host defenses and help bacteria prosper on the device[2,14].

Diagnosis is made considering all clinical, laboratory, histopathological, microbiological, and imaging findings. If the patient has no symptoms such as continuous back pain, fever, or shivers, the surgeon should carefully inspect the wound for any local inflammatory signs[13]. Repetitive postoperative CRP measurements can provide valuable information during the postoperative period. Histopathological analysis of periprosthetic tissue is reported to have more than 80% of sensitivity and more than 90% of specificity in the literature[10]. The rapid detection of infection is critical for preventing osteomyelitis and device loss. Radiological assessment is crucial for detecting mechanical complications, such as spinal instability and inflammation, abscess formation, or fluid collection[13].

Despite the devastating results, the optimal management of SSI after spinal instrumentation remains controversial[13]. A multidisciplinary approach involving surgeons, microbiologists, and infectious disease specialists is essential to achieve the best outcome. When the microorganism causing the infection is of low virulence and is reactive to antibiotics, mostly in postoperative three to six weeks, device retention may be possible. Surgical debridement may be needed, and intravenous antibiotics for more than 6 wk are recommended, along with subsequent oral antibiotics. In cases of more than postoperative 6 wk, additional surgery is inevitable to reduce the inoculum size, which may be an explanation of the device or exchanging it depending on the spine condition. Debridement should ensure complete resection of necrotic tissue, dead bone, abscess membranes, and unhealthy granulation tissue[11]. Plastic and reconstructive interventions are required in cases of chronic osteomyelitis or large soft tissue defects. Prolongation of antibiotic use should be planned, and fine adjustment is required regarding the patient's underlying disease, type of implant and pathogen, and most importantly, clinical presentation[13].

Flap reconstruction of the back is not commonly compared with other regions; therefore, related literature is limited [16]. Surgical reconstruction in this region remains challenging because the lower back has low skin laxity and is securely attached to underlying vital structures. Pressure wounds, tumor resection, and spine surgeries such as laminectomy, discectomy, and spinal arthrodesis can cause soft tissue defects in the lower back. When bones or implants are exposed, because there are few bulky muscles available, reconstruction options are traditionally limited to the paraspinal muscle flap, turnover latissimus dorsi muscle flap, bipediced paraspinal muscle flap, and free flaps[16]. Perforator flaps have gained popularity due to advances in anatomy. They are widely used in back-defect repair, in preference to local random or free flaps. Even large lumbar defects can be reconstructed using intercostal artery, lumbar artery, and superior gluteal artery perforators, and pedicled flaps[3]. A single perforator supplies a wide base of tissue[16]. It is advantageous in many ways, including mobility, desired shape, and shorter operative time. The biggest advantage is low donor site morbidity and preservation of the major back muscles, which preserves shoulder and arm function[16]. It also decreases the operation time compared to traditional flaps, which is essential considering that the surgical position is inevitably limited to the prone or lateral positions. Other options include lumbar artery and superior gluteal artery perforator flaps. They are the two major perforators in the lumbosacral region and have large interactions[4].

In the aforementioned case, a delayed chronic infection occurred after spinal surgery. Chronic infection was not easily noticed because the patient had no symptoms, including pain or fever, and all laboratory results were within the normal range. It might have been better if the authors suspected a neglected or subclinical infection, and the wire was removed before flap surgery. However, evidence supporting the removal of deep device was insufficient at that time. We assume that previous hematoma, wound infection at the first surgery, and repetitive spine surgery may have induced biofilm infection in the deep tissue and implant devices. Consequently, biofilm formation may have caused the chronic infection observed in our case.

---

## CONCLUSION

Repetitive surgical wound infections and dehiscence can induce delayed infection, making complete wound healing challenging. Imaging studies and other laboratory results do not reveal significant infection; however, deep tissue - should always be strongly suspected when wound dehiscence recurs. In our case, we reconstructed chronic lumbar spinal wound dehiscence with a perforator-based flap, and as for the recurred deep tissue infection, successful salvage was achieved through infected wire removal. In cases of inappropriately diagnosed or recently treated surgical wound infections, devastating results in need of aggressive surgical treatment, including free flaps, is required. The authors recommend suspicion of SSI, even deep device infection, when surgeons encounter chronic SWD.

---

## FOOTNOTES

**Author contributions:** Kim D and Yoon JS contributed to manuscript writing and editing, and data collection; Yoon JS contributed to conceptualization and supervision; and all authors have read and approved the final manuscript.

**Informed consent statement:** Informed written consent was obtained from the patient for publication of this report and any accompanying images.

**Conflict-of-interest statement:** All the authors report no relevant conflicts of interest for this article.

**CARE Checklist (2016) statement:** The authors have read the CARE Checklist (2016), and the manuscript was prepared and revised according to the CARE Checklist (2016).

**Open-Access:** This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

**Country/Territory of origin:** South Korea

**ORCID number:** DoWon Kim 0000-0003-2089-9896; SooA Lim 0000-0003-3845-780X; SuRak Eo 0000-0002-4221-2613; Jung Soo Yoon 0000-0003-2462-5702.

**S-Editor:** Wang JJ

**L-Editor:** A

**P-Editor:** Wang JJ

---

## REFERENCES

---

- 1 **Sandy-Hodgetts K**, Carville K, Leslie GD. Surgical wound dehiscence: a conceptual framework for patient assessment. *J Wound Care* 2018; **27**: 119-126 [PMID: 29509107 DOI: 10.12968/jowc.2018.27.3.119]
- 2 **Breder JSC**, Tsukumo DML, Pereira E, Lima MH. Surgical Wound Dehiscence Treatment With Low-Level Laser Therapy and Barbatimão: A Case Report. *Wound Manag Prev* 2021; **67**: 18-22 [PMID: 35030091]
- 3 **Ahmed O**, Storey CM, Zhang S, Chelly MR, Yeoh MS, Nanda A. Vacuum-assisted closure of necrotic and infected cranial wound with loss of dura mater: A technical note. *Surg Neurol Int* 2015; **6**: 11 [PMID: 25657864 DOI: 10.4103/2152-7806.149846]
- 4 **Ploumis A**, Mehbod AA, Dressel TD, Dykes DC, Transfeldt EE, Lonstein JE. Therapy of spinal wound infections using vacuum-assisted wound closure: risk factors leading to resistance to treatment. *J Spinal Disord Tech* 2008; **21**: 320-323 [PMID: 18600140 DOI: 10.1097/BSD.0b013e318141f99d]
- 5 **Labler L**, Keel M, Trentz O, Heinzelmann M. Wound conditioning by vacuum assisted closure (V.A.C.) in postoperative infections after dorsal spine surgery. *Eur Spine J* 2006; **15**: 1388-1396 [PMID: 16835734 DOI: 10.1007/s00586-006-0164-2]
- 6 **Trampuz A**, Widmer AF. Infections associated with orthopedic implants. *Curr Opin Infect Dis* 2006; **19**: 349-356 [PMID: 16804382 DOI: 10.1097/01.qco.0000235161.85925.e8]
- 7 **Zimmerli W**. Clinical presentation and treatment of orthopaedic implant-associated infection. *J Intern Med* 2014; **276**: 111-119 [PMID: 24605880 DOI: 10.1111/joim.12233]
- 8 **Chue AL**, Moran E, Atkins BL, Byren I. Orthopaedic device associated Candidal infection - a retrospective review of patient characteristics and outcome. *J Infect* 2011; **63**: 167-171 [PMID: 21669224 DOI: 10.1016/j.jinf.2011.05.018]
- 9 **Barrey C**, Launay O, Freitas E, Michel F, Laurent F, Chidiac C, Perrin G, Ferry T. The follow-up of patients with postoperative infection of the spine. *Eur J Orthop Surg Traumatol* 2013; **23** Suppl 1: S29-S34 [PMID: 23712673 DOI: 10.1007/s00590-013-1243-9]
- 10 **Stocks G**, Janssen HF. Infection in patients after implantation of an orthopedic device. *ASAIO J* 2000; **46**: S41-S46 [PMID: 11110293 DOI: 10.1097/00002480-200011000-00036]
- 11 **Lee K**, Yoon SS. Pseudomonas aeruginosa Biofilm, a Programmed Bacterial Life for Fitness. *J Microbiol Biotechnol* 2017; **27**: 1053-1064 [PMID: 28301918 DOI: 10.4014/jmb.1611.11056]
- 12 **Zimmerli W**. Orthopaedic device-associated infection. *Clin Microbiol Infect* 2012; **18**: 1160-1161 [PMID: 23148446 DOI: 10.1111/j.1469-0691.2012.04011.x]
- 13 **Durgun M**, Baş S, Aslan C, Canbaz Y, Işık D. Use of dorsal intercostal artery perforator flap in the repair of back defects. *J Plast Surg Hand Surg* 2016; **50**: 80-84 [PMID: 26542196 DOI: 10.3109/2000656X.2015.1102737]
- 14 **Komagoe S**, Watanabe T, Komatsu S, Kimata Y. Pedicled Flaps Versus Free Flaps for Back Reconstruction. *Ann Plast Surg* 2018; **81**: 702-707 [PMID: 30247187 DOI: 10.1097/SAP.0000000000001590]
- 15 **Kato H**, Hasegawa M, Takada T, Torii S. The lumbar artery perforator based island flap: anatomical study and case reports. *Br J Plast Surg* 1999; **52**: 541-546 [PMID: 10658107 DOI: 10.1054/bjps.1999.3120]
- 16 **Di Rienzo A**, Colasanti R, Liverotti V, Benigni R, Paracino R, Bizzocchi G, Scerrati M, Iacoangeli M. On-ward surgical management of wound dehiscence: report of a single neurosurgical center experience and comparison of safety and effectiveness with conventional treatment. *Neurosurg Rev* 2020; **43**: 131-140 [PMID: 30120610 DOI: 10.1007/s10143-018-1022-5]



Published by **Baishideng Publishing Group Inc**  
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA  
**Telephone:** +1-925-3991568  
**E-mail:** [bpgoffice@wjgnet.com](mailto:bpgoffice@wjgnet.com)  
**Help Desk:** <https://www.f6publishing.com/helpdesk>  
<https://www.wjgnet.com>

