

90388_Auto_Edited.docx

Name of Journal: *World Journal of Clinical Cases*

Manuscript NO: 90388

Manuscript Type: CASE REPORT

Cardiac remodeling in patients with atrial fibrillation reversing bradycardia-induced cardiomyopathy: A case report

AF reversing BIC Cardiac remodeling

Abstract

BACKGROUND

Bradycardia-induced cardiomyopathy (BIC), a disease resulting from bradycardia, is characterized by cardiac chamber enlargement and diminished cardiac function. The correction of bradycardia has demonstrated significant improvements in both cardiac function and structure; however, this disease has been infrequently documented. In this case, we conducted a longitudinal follow-up of a patient who endured BIC for **more than 40 years** to heighten awareness and prompt timely diagnosis and rational intervention for BIC.

CASE SUMMARY

A woman presented with post-activity fatigue and dyspnea was diagnosed with bradycardia at the age of 7. Since she had no obvious symptoms, she did not receive any treatment to improve her bradycardia during the 42-year follow-up except for the implantation of a temporary pacemaker during labor induction surgery. As time went by, the patient's heart gradually expanded due to low ventricular rate, and she was diagnosed with bradycardia-induced cardiomyopathy (BIC). In 2014, the patient developed atrial fibrillation (AF), and her ventricular rate gradually increased, and then her heart shape gradually returned to normal. This report described the cardiac morphological changes caused by heart rate changes in the BIC patient **over 40 years**, introduced another possible outcome of BIC, and also emphasized the importance of early intervention in BIC.

CONCLUSION

BIC can induce AF, causing increased ventricular rate and leading to positive cardiac remodeling.

Key Words: Bradycardia cardiomyopathy; Heart rate; Atrial fibrillation; Cardiac dilatation; Case report

Gao D, Ye X, Duan Z, Zhang H, Xiong T, Li Z, pei H. Cardiac remodeling in patients with atrial fibrillation reversing bradycardia-induced cardiomyopathy: A case report. *World J Clin Cases* 2024; In press

Core Tip: Bradycardia-induced cardiomyopathy (BIC), is a disease caused by bradycardia and characterized by enlargement of the heart chambers and decreased cardiac function; increasing the ventricular rate significantly improved the associated symptoms and cardiac remodeling, however, this disease has rarely been reported. This case described a woman with BIC who, with little therapeutic intervention to improve bradycardia, had an increase in ventricular rate due to the onset of atrial fibrillation and then her cardiac enlargement due to low ventricular rate was reversed. This case also emphasized the importance of early intervention for BIC.

INTRODUCTION

Prolonged tachyarrhythmia and bradyarrhythmia can significantly impact cardiac function. While tachyarrhythmia-induced cardiomyopathy (TIC) has garnered considerable attention in recent studies, evidence suggests that bradycardia also plays a role in cardiomyopathy progression. Fabritz *et al*^[1]. were the pioneers in introducing the concept of "Bradycardiomyopathy." The pathogenesis of bradycardia-induced cardiomyopathy (BIC) may be linked to a prolonged ventricular diastolic phase, volume overload, abnormal atrial and ventricular excitation and contraction sequences, adverse myocardial perfusion, neuroendocrine function alterations, among other factors. Despite these insights, reports on bradycardia cardiomyopathy remain scarce. In our longitudinal follow-up spanning over 40 years, we tracked a female patient with bradycardia-induced cardiomyopathy, documenting the transformation of her cardiac chambers and morphology from abnormal to normal as heart rate increased, predominantly without pacing therapy. This endeavor aims to heighten awareness and prompt timely diagnosis and rational intervention for BIC.

CASE PRESENTATION

Chief complaints

A 7-year-old female presented with post-activity cardiac fatigue and dyspnea and was admitted to the general medical of our hospital in December 1980. After being discharged from the hospital, she was followed up in our hospital for a long time because of arrhythmia.

History of present illness

The female presented with post-activity fatigue and dyspnea at the age of 7, electrocardiographic (ECG) assessment revealed arrhythmia and bradycardia, culminating in a diagnosis of "myocarditis and sick sinus syndrome (SSS) ". Her symptoms of fatigue and dyspnea improved after treatment with atropine and prednisone (details unknown), but her ventricular rate remained at a low level. Throughout the course of treatment, there were no episodes of syncope or amaurosis. After discharge, she was followed up in our hospital for a long time because of low heart rate. The follow-up ECG, echocardiography (UCG) and chest X-ray showed that she had bradycardia and cardiac enlargement. Nine years ago (2014), the patient developed atrial fibrillation, the ventricular rate gradually increased while the dilated heart gradually returned to normal, during which the patient did not show any cardiac symptoms.

History of past illness

The patient has no history of congenital heart disease and has never developed hypertension, diabetes, valvular heart disease, hyperthyroidism, hypothyroidism, hyperlipidemia or other diseases related to cardiac remodeling. In 2001, the patient could not bear pregnancy because of bradycardia, so she chose induced abortion and temporary pacemaker was installed before operation. In November, 2012, the patient was admitted to the hospital for uterine fibroids. Due to bradycardia, the patient again

had to install a temporary pacemaker before surgery. She does not use specific drugs or drugs, has no history of smoking or drinking, and has no history of radiation or toxin exposure.

Personal and family history

The female's menstruation was reported as normal (the patient had menarche at the age of 12, with a menstrual cycle of 28-30 days, and each menstrual period lasted for 3-5 days. She became menopausal in February 2020. The periods were regular, and the menstrual flow was moderate, without dysmenorrhea). No familial occurrences of a similar disease were identified.

Physical examination

none

Laboratory examinations

The female's Laboratory examinations did not reveal any abnormalities.

Imaging examinations

On June 2, 1981 after prednisone and atropine treatment, Echocardiography (UCG) showed enlargement of left atrium and left ventricle (details unknown).

On November.30,2001, UCG suggested that the ventricular rate was 39BPM, right atrium and right ventricle were slightly enlarged, Right Atrial Transverse Diameter (RA): 53mm; Aortic Valve Peak Systolic Flow Velocity (AV): 1.73m/s.

On November.9,2012, UCG hint: right heart enlargement, RA:49mm, AV:2.19m/s; chest X-ray hint: cardiothoracic ratio 0.65.

On July 02, 2014, UCG suggested that all chambers of the heart were normal, and RA:40mm, AV:1.14m/s; chest X-ray showed that the cardiothoracic ratio was 0.53.

On May.10, 2016, UCG hint: atrial enlargement, RA:52mm, AV:1.40m/s; on May 16 chest X-ray indicated that the heart was enlarged, the cardiothoracic ratio was 0.59.

On September.27, 2020, UCG indicated that the atrium was slightly enlarged, RA:46mm, AV:1.31m/s.

On May.06,2022, UCG suggested: Atrial enlargement, RA:47mm,AV:0.87m/s.

UCG on December.20,2022: Heart morphology was normal, RA:39mm, AV:0.82m/s, with minimal trace regurgitation in the bicuspid, tricuspid, and pulmonary valves.

ECG EXAMINATIONS

The patient's early ECG manifestations included sinus arrest, junctional escape rhythm with a rate of approximately 40 beats per minute (BPM), sporadic frequent ventricular premature beats, brief bursts of ventricular tachycardia, and infrequent sinus arrest with first-degree atrioventricular block (AVB) at 8 years old. Notably, two prolonged gaps were observed, with the lengthiest RR interval reaching 4680ms. The suggestion for an artificial pacemaker implantation was made due to these findings, but the recommendation was declined by the patient.

In 2001, the patient, unable to endure the pregnancy due to bradycardia, opted for induced abortion with preoperative installation of a temporary pacemaker. The unipolar electrocardiogram derived from the temporary pacemaker electrodes, situated in the elevated right atrium, revealed sinus arrest with intermittent atrial electrical activity (various A-waves), near-complete atrioventricular dissociation, and junctional escape rhythm. Stimulation of the elevated right atrium with S₁S₁ (940ms) induced atrial agitation, with a 1:1 conduction to the ventricle, while S₁R prolonged to 310ms.

Upon admission for uterine fibroids on November 9, 2012, the patient exhibited an ECG depicting junctional escape rhythm at 45 BPM.

ECG assessments on July 2, 2014 showed atrial fibrillation with atrial rates of 250 BPM and a mean ventricular rate of 74 BPM.

The ECG measured on May 10, 2016 showed: atrial fibrillation, with a mean ventricular rate 125BPM.

On September 27, 2020, the ECG indicated atrial fibrillation with a mean heart rate of 90 BPM.

On July 8, 2022, the ECG revealed atrial flutter with a ventricular rate of 83 BPM.

FINAL DIAGNOSIS

The patient was diagnosed with a BIC.

TREATMENT

The patient was treated with prednisone and atropine for a short period when symptoms became apparent at the age of 7 years, and subsequently received a temporary pacemaker implant during induction of labor and uterine fibroid surgery. She did not undergo any treatment to increase the ventricular rate, including pacemaker implantation, thereafter. Even after atrial fibrillation occurred, not only did radiofrequency ablation combined with pacemaker implantation not be performed, but anticoagulation treatment was even refused.

OUTCOME AND FOLLOW-UP

We organized the ECG and UCG data collected spanning over 20 years chronologically (Figure 1). Our analysis revealed that the patient exhibited bradycardia until the age of 40 (before 2014), marked by an enlarged cardiac morphology and dilated chambers as indicated by UCG and chest radiographs. Subsequently, the patient experienced a transition to atrial flutter/atrial fibrillation, leading to a gradual increase in ventricular rate. Remarkably, without any specific treatment, the cardiac morphology gradually normalized over this period. Notably, significant alterations in the right atrial transverse diameter and peak systolic flow velocity of the aortic valve were observed, both of which normalized with the increase in ventricular rate. This observation suggests that an elevated ventricular rate can effectively reverse the dilated state of the heart in patients with BIC, leading to improvements in hemodynamics and cardiac

function. Considering the patient's age-related high risk of developing tachyarrhythmia-induced cardiomyopathy, we strongly advocate for cardiac radiofrequency ablation.

DISCUSSION

In recent years, multiple studies have underscored a significant correlation between cardiomyopathies and arrhythmias, with the well-established concept of tachyarrhythmia-induced cardiomyopathy (TIC). Radiofrequency ablation has emerged as an efficacious intervention for a growing number of patients experiencing tachyarrhythmias, fostering improvements in heartbeat and the restoration of cardiac morphology and function. Conversely, investigations into bradycardia-induced cardiomyopathy (BIC), a disease resulting from abnormal heart rates characterized by cardiac chamber enlargement and diminished cardiac function, are limited. Historical research in animals revealed that chronic complete high-grade atrioventricular block led to ventricular cardiomyocyte hypertrophy^[3]. Sasaki^[4] documented a case of restrictive cardiomyopathy and paroxysmal atrial fibrillation, wherein the transition to sinus rhythm induced right heart failure, later alleviated by pacemaker implantation and pharmacological maintenance of sinus rhythm. Kovács T^[5] reported sinus arrest with escape rhythm (40 BPM) in a hypertensive patient on β -blockers, highlighting elevated NT-proBNP (N-terminal B-type brain natriuretic peptide precursor), which normalized upon discontinuation of β -blockers. Fabritz^[1] proposed "bradycardiomyopathy" in an animal study, linking severe bradycardia to dilatation, fibrosis, and atrioventricular dysfunction. Caliskan K^[6] reported significant improvement in a male patient with sinus bradycardia and non-dense cardiomyopathy after pacemaker implantation, establishing a causal link. Our documented case of BIC, spanning over 40 years with a gradual shift in cardiac morphology from abnormal to normal, represents unique empirical data, offering the first extended observational insight into the natural evolution of BIC.

Similar to the diagnosis of TIC, the retrospective diagnosis of BIC hinges on specific criteria. Diagnostic parameters for BIC include: 1. a history of prolonged bradycardia; 2.

manifestation of pertinent symptoms and signs, encompassing weakness, dizziness, amaurosis, and, in severe cases, fainting, along with signs and symptoms of heart failure; 3. ultrasound confirmation of enlarged cardiac chambers in the context of prolonged bradycardia, while maintaining a generally normal left ventricular ejection fraction; and 4. normalization of cardiac chambers and function following pacing treatment. The augmentation of ventricular rate through pacemaker implantation or alternative methods has been demonstrated to yield substantial improvements in cardiac function and structure in such patients^[7]. Notably, the patient in question initially exhibited symptoms of cardiac dysfunction during the early stages of bradycardia diagnosis. The disease's evolution paralleled that of other arrhythmia-related cardiomyopathies, with the increasing ventricular rate facilitating a partial recovery of heart morphology and function. This correlation underscores the close association between cardiomyopathy-like changes and bradycardia, substantiating the diagnosis of BIC.

In this case, beginning with the childhood diagnosis of sick sinus node syndrome, the patient progressively exhibited manifestations of bradycardia, sinus arrest, junctional escape beats, and first-degree atrioventricular block. Despite meeting criteria for pacemaker implantation, as evidenced by the need for implantation after atrial fibrillation radiofrequency ablation^[8], the patient's refusal of radiofrequency ablation and subsequent prolonged period without cardiac-related symptoms allowed for the observation of cardiac morphologic changes associated with increased heart rate. Chronic bradycardia in this patient resulted in a relative increase in ventricular filling time, causing elevated ventricular wall tension, cardiac off-center hypertrophy, and an increase in the transverse diameter of the right atrium. Additionally, prolonged bradycardia activated the sympathetic nervous system, inducing vasoconstriction, increased myocardial contractility, elevated resistance to cardiac efflux, raised peak flow velocity of the aortic valve, and promoted myocardial fibrosis, potentially triggering atrial fibrillation^[9]. Following the onset of atrial fibrillation, the patient's ventricular rate gradually increased, a strategy shown to mitigate deleterious cardiac

remodeling by maintaining a heart rate below 110 BPM in patients with early atrial fibrillation^[10,11]. Remarkably, without intervention, cardiac remodeling reversed progressively with the rising ventricular rate, culminating in the restoration of normal cardiac morphology and function. The right atrial transverse diameter and peak aortic systolic flow velocity gradually normalized, suggesting that pacing therapy or an increased ventricular rate due to atrial fibrillation/atrial flutter can ameliorate abnormal cardiac morphology and function in patients with BIC. As the patient ages, the potential for the recurrence of atrial enlargement and ventricular modifications in response to heart rate changes, possibly progressing towards TIC, poses a notable risk of atrial fibrillation-induced stroke. Consequently, we strongly advocate for cardiac radiofrequency ablation and the routine use of non-vitamin K antagonist oral anticoagulants before and after resuscitation^[10]. Increased awareness of BIC and early intervention holds the potential to positively impact the prognosis of this disease.

The landscape of bradycardia treatment underwent a paradigm shift in 1958 when Swedish cardiac surgeon Ake Senning pioneered the first pacemaker implantation in human history^[12]. This landmark development has significantly limited opportunities for observing the progression of bradycardia to BIC. While the evolution from arrhythmia to arrhythmogenic cardiomyopathy typically spans months to years, recurrent arrhythmias can hasten the onset of heart failure, accompanied by a rapid decline in ventricular function. Early diagnosis and effective arrhythmia management have shown considerable potential in improving ventricular function and alleviating clinical symptoms. The prompt identification of electromechanical dysfunction in individuals with a normal left ventricular ejection fraction is now achievable through the assessment of cardiac electrical and mechanical function using acoustic cardiography (ACG)^[13,14]. ACG serves as a valuable early screening tool for heart failure.

The presented case is subject to certain limitations. The first limitation is the lack of ECGs prior to the patient reaching 8 years of age, which precludes a definitive determination of congenital sinus node dysfunction and subtle cardiac insufficiency

during that period. The second limitation is the absence of empirical evidence from the initial hospitalization supporting the diagnosis of myocarditis. This report described the cardiac morphological changes caused by heart rate changes in the BIC patient over 40 years, introduced another possible outcome of BIC. But we still need to remain vigilant about BIC.

CONCLUSION

The documented cardiac structural changes in this case indicates BIC can induce AF, causing increased ventricular rate and leading to positive cardiac remodeling, which are believed to be inherent due to no therapeutic interventions aimed at cardiac remodeling implemented in response to the detected heart rate abnormality, with few similar instances reported in the literature. The etiology of BIC remains uncertain, further complicated by the scarcity of prospective studies. The gradual, often unnoticed development of this condition may lead to a potential underestimation of its incidence. Therefore, maintaining an enhanced level of vigilance for BIC is of paramount importance in clinical practice.

ORIGINALITY REPORT

0%

SIMILARITY INDEX

PRIMARY SOURCES

EXCLUDE QUOTES	ON	EXCLUDE SOURCES	< 12 WORDS
EXCLUDE BIBLIOGRAPHY	ON	EXCLUDE MATCHES	< 12 WORDS