

1 / 58

Name of journal: *World Journal of Gastroenterology*

Manuscript NO: 54927

Manuscript Type: SYSTEMATIC REVIEWS

Differential diagnosis of diarrhoea in patients with neuroendocrine tumours: A systematic review

Mohid S Khan, Thomas Walter, Amy Buchanan-Hughes, Emma Worthington, Lucie Keeber, Marion Feuilly, Enrique Grande

Abstract

BACKGROUND

Match Overview

| | | |
|---|--|-----|
| 1 | Internet 27 words crawled on 27-Mar-2020 www.magonlinelibrary.com | <1% |
| 2 | Crossref 16 words Jenni Antikainen, Anu Kantele, Sari H. Pakkanen, Tinja Lä äveri, Jukka Riutta, Martti Vaara, Juha Kirveskari. "A Quant | <1% |
| 3 | Internet 14 words crawled on 14-Jun-2020 mafiadoc.com | <1% |
| 4 | Internet 13 words crawled on 15-Jan-2018 www.karger.com | <1% |
| 5 | Internet 13 words crawled on 16-Nov-2017 hal.archives-ouvertes.fr | <1% |
| 6 | Internet 13 words crawled on 01 May 2020 | <1% |



国内版

国际版

Differential Diagnosis of Diarrhoea in Patients with Neurc



ALL

IMAGES

VIDEOS

27,100 Results

Any time ▾

Prevalence of non-carcinoid syndrome (non-CS) diarrhoea in ...

<https://www.endocrine-abstracts.org/ea/0063/ea0063p446> ▾

Prevalence of non-carcinoid syndrome (non-CS) **diarrhoea in patients with neuroendocrine tumours** (NETs): **a systematic literature review** (SLR) Mohid S Khan 1, Thomas Walter 2, Amy Buchanan-Hughes 3, Lucie Keeber 4, Marion Feuille 5 & Enrique Grande 6

Author: Mohid S Khan, Thomas Walter, Amy ... **Publish Year:** 2019

[PDF] Framework Synthesis (FS) of Qualitative and Quantitative ...

https://www.costellomedical.com/wp-content/uploads/2019/08/PMU90_ISPOR_US_2019.pdf

to explore evidence (identified via **systematic literature review** [SLR]) relating to DDX of **diarrhea in patients** with GEP-NETs, for which little quantitative data are available. **Framework Synthesis** (FS) of Qualitative and Quantitative Evidence to Explore the **Differential Diagnosis** (DDX) of **Diarrhea in Neuroendocrine Tumor** (NET) **Patients**

Differential Diagnosis of Neuroendocrine Tumors | Request ...

<https://www.researchgate.net/publication/307522548...>

As compared to NSCLCs, only little research has been done with respect to new molecular targets as well as improving the classification and **differential diagnosis of neuroendocrine** tumors of the lung.

[PDF] Differential Diagnosis (DDX) of Carcinoid Syndrome ...

<https://nanets.net/abstracts-archive/2019/1345-p20-differential-diagnosis-ddx-of...>

tests was conflicting. If **diarrhea** is misdiagnosed, treatment may be deemed ineffective and underlying conditions could remain undetected. P-20 **Differential Diagnosis** (DDX) of Carcinoid Syndrome **Diarrhea** (CSD): **A Systematic Literature Review** (SLR)

Chat with Bing



You know, sometimes I really don't understand you human...

Say something...

国内版

国际版

Differential diagnosis of diarrhoea in patients with neuroendocrine



ALL

IMAGES

VIDEOS

42,700 Results

Any time ▼

[Chronic Diarrhea in Adults: Evaluation and Differential ...](#)

<https://www.aafp.org/afp/2020/0415/p472.html> ▼

Apr 15, 2020 · Chronic **diarrhea** is defined as a predominantly loose stool lasting longer than four weeks. A **patient** history and physical examination with a complete blood count, C-reactive protein, anti-tissue ...

[Management of Diarrhea in Patients With Carcinoid Syndrome](#)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6867674>

Aug 19, 2019 · Khan MA, Walter **TB**, Buchanan-Hughes AC, et al. **Differential diagnosis (DDx) of carcinoid syndrome diarrhea (CSD): a systematic** literature review (SLR). Abstract J08. Barcelona, Spain: Presented at the 16th Annual **European Neuroendocrine Tumor Society Conference**; March 6–8; 2019. [Google Scholar]

[Images of Differential Diagnosis of Diarrhoea In Patients with ...](#)

<bing.com/images>





Differential diagnosis of diarrhoea in patients with neuro



ALL

IMAGES

VIDEOS

MAPS

NEWS

SHOPPING

43,500 Results

Any time ▾

[Chronic Diarrhea in Adults: Evaluation and Differential ...](https://www.aafp.org/afp/2020/0415/p472.html)

<https://www.aafp.org/afp/2020/0415/p472.html>

Apr 15, 2020 · Chronic **diarrhea** is defined as a predominantly loose stool lasting longer than four weeks. A **patient** history and physical examination with a complete blood count, C-reactive protein, anti-tissue ...

[Prevalence of non-carcinoid syndrome \(non-CS\) diarrhoea in ...](https://www.endocrine-abstracts.org/ea/0063/ea0063p446)

<https://www.endocrine-abstracts.org/ea/0063/ea0063p446>

Prevalence of non-carcinoid syndrome (non-CS) **diarrhoea** in **patients with neuroendocrine tumours (NETs): a systematic literature review** (SLR) Mohid S Khan 1, Thomas Walter 2, Amy Buchanan-Hughes 3, Lucie Keeber 4, Marion Feuilly 5 & Enrique Grande 6

Author: Mohid S Khan, Thomas Walter, Amy B... **Publish Year:** 2019

[Management of Diarrhea in Patients With Carcinoid Syndrome](https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6867674)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6867674>

Aug 19, 2019 · **Differential Diagnosis.** In assessing a **patient** with suspected CS **diarrhea**, it is important to differentiate the GI symptoms of NETs from the **diarrhea**, abdominal pain, and dyspepsia related to other causes (eg, irritable bowel syndrome) by eliciting a thorough **patient** history that details age, sleep patterns, and nonspecific symptoms. 67 ...

[Differential Diagnosis of Neuroendocrine Tumors | Request PDF](https://www.researchgate.net/publication/307522548...)

<https://www.researchgate.net/publication/307522548...>

The accurate **diagnosis** and appropriate treatment of GEP-NET currently gives most **patients** with this **tumor** a good prognosis, as long as it is discovered early, and **differential diagnosis** of