

1,810 Results Any time ▾

[The determination of sensation-dependent bladder emptying ...](#)<https://www.nature.com/articles/3102102/tables>

Retrospective investigation using urodynamic studies and medical records. To evaluate the safety of sensation-dependent bladder emptying in complete spinal cord injury (SCI) patients, based on the ...

Cited by: 5

Author: J C Shin, W H Chang, T H Jung, J H Yoo, S ...

Publish Year: 2008

[The determination of sensation-dependent bladder emptying ...](#)<https://www.semanticscholar.org/paper/The...>

Study design: Retrospective investigation using urodynamic studies and medical records. Objective: To evaluate the safety of sensation-dependent bladder emptying in complete spinal cord injury (SCI) patients, based on the preservation of the desire to void. Setting: Spinal Cord Injury Unit, Yonsei Rehabilitation Hospital, Seoul, Korea. Methods: This study was performed retrospectively on 79 complete ...

[The determination of sensation-dependent bladder emptying ...](#)<https://www.pubfacts.com/detail/17646839/The...>

Study Design: Retrospective investigation using urodynamic studies and medical records. Objective: To evaluate the safety of sensation-dependent bladder emptying in complete spinal cord injury (SCI) patients, based on the preservation of the desire to void. Setting: Spinal Cord Injury Unit, Yonsei Rehabilitation Hospital, Seoul, Korea. Methods: This study was performed retrospectively on 79 ...

[Bladder-filling sensation in patients with spinal cord ...](#)https://www.academia.edu/28182092/Bladder_filling...

Bladder-filling sensation in patients with spinal cord injury and the potential for sensation-dependent bladder emptying. Spinal Cord, 2004. Mufit Akyuz. Download PDF. Download Full PDF Package. This paper. A short summary of this paper. 37 Full PDFs related to this paper. READ PAPER.

[Bladder-filling sensation in patients with spinal cord ...](#)<https://www.nature.com/articles/3101525?error=...>

Study design: Investigation of bladder-filling sensation in 73 consecutive traumatic spinal cord injury (SCI) patients during laboratory cystometry. Objectives: To determine the frequencies of ...

Cited by: 26

Author: M Ersoz, M Akyuz

Publish Year: 2004

[\[PDF\] ORIGINAL ARTICLE The determination of sensation ...](#)https://www.researchgate.net/profile/Won-Hvuk-Chang/publication/6191544_The...

1,780 Results Any time ▾

The determination of sensation-dependent bladder emptying ...

<https://www.nature.com/articles/3102102/tables>

Retrospective investigation using urodynamic studies and medical records. To evaluate the safety of sensation-dependent **bladder emptying** in **complete spinal cord injury** (SCI) patients, based on the ...

Cited by: 5

Author: J C Shin, W H Chang, T H Jung, J H Yoo, S ...

Publish Year: 2008

The determination of sensation-dependent bladder emptying ...

<https://www.semanticscholar.org/paper/The...>

Study design:Retrospective investigation using urodynamic studies and medical records.Objective:To evaluate the safety of sensation-dependent **bladder emptying** in **complete spinal cord injury** (SCI) patients, based on the preservation of the desire to void.Setting:**Spinal Cord Injury** Unit, Yonsei Rehabilitation Hospital, Seoul, Korea.Methods:This study was performed retrospectively on 79 **complete** ...

The determination of sensation-dependent bladder emptying ...

<https://www.pubfacts.com/detail/17646839/The...> ▾

Study Design: Retrospective investigation using urodynamic studies and medical records. Objective: To evaluate the safety of sensation-dependent **bladder emptying** in **complete spinal cord injury** (SCI) patients, based on the preservation of the desire to void. Setting: **Spinal Cord Injury** Unit, Yonsei Rehabilitation Hospital, Seoul, Korea. Methods: This study was performed retrospectively on 79 ...

Name of Journal: *World Journal of Clinical Cases*

Manuscript NO: 69737

Manuscript Type: CASE REPORT

**Disruption of sensation-dependent bladder emptying due to bladder overdistension
in a complete spinal cord injury: A case report**

Yoon JY *et al.* Detrimental effect of bladder overdistension

Match Overview

1	Crossref 13 words K Furusawa, A Tokuhito, H Sugiyama, A Ikeda, F Tajima, E Genda, R Uchida, T Tominaga, H Tanaka, A Magara, M Sum	1%
2	Internet 12 words crawled on 29-Sep-2020 www.ejgnet.com	1%

国内版

国际版

Disruption of sensation-dependent bladder emptying due to blad



ALL

IMAGES

VIDEOS

4,010 Results

Any time ▾

Bladder-filling sensation in patients with spinal cord ...

<https://www.researchgate.net/publication/8884622...>

Bladder-filling sensation was present to some degree in all incomplete SCI patients, in 82.4% of the patients with complete lesions below T10, and 38.9% of the patients with complete lesions above ...

Adult Neurogenic Lower Urinary Tract Dysfunction and ...

[https://www.hindawi.com/journals/au/2019/2757862 ▾](https://www.hindawi.com/journals/au/2019/2757862...)

With any spinal cord injury, the disruption of the normal somatic and autonomic neurological control of bladder causes bladder dysfunction; in higher suprasacral injuries, a micturition reflex may emerge after the acute phase to cause detrusor overactivity and sometimes detrusor-sphincter dyssynergia (DSD) with ineffective bladder emptying and clinical significant large postvoid residual volumes.

Cited by: 11

Author: Michael Kennelly, Nikesh Thiruchelvam, ...

Publish Year: 2019

PEOPLE ALSO ASK

What are the risks of neurogenic bladder dysfunction? ▾

Can a spinal cord injury cause urinary tract dysfunction? ▾

What happens to the spinal cord during urine voiding? ▾

What is the term for postoperative urinary retention? ▾

Feedback

The management of neurogenic lower urinary tract ...

<https://www.nature.com/articles/nrurol.2016.206>

Oct 25, 2016 · Ersoz, M. & Akyuz, M. Bladder-filling sensation in patients with spinal cord injury and the potential for sensation-dependent bladder emptying. Spinal Cord 42 , 110–116 (2004).

Cited by: 54

Author: Jean-Jacques Wyndaele

Publish Year: 2016

Lower urinary tract dysfunction in patients with ...