

Name of Journal: *World Journal of Gastroenterology*

Manuscript NO: 64319

Manuscript Type: ORIGINAL ARTICLE

Retrospective Study

Idiopathic mesenteric phlebosclerosis associated with long-term oral intake of geniposide

Wen Y *et al.* Idiopathic mesenteric phlebosclerosis associated with geniposide

Yang Wen, You-Wei Chen, Ai-Hong Meng, Ming Zhao, Song-Hua Fang, Yan-Qing Ma

Match Overview

1	Internet 46 words crawled on 04-Apr-2017 www.wjgnet.com	1%
2	Internet 36 words crawled on 06-Jun-2020 journals.lww.com	1%
3	Crossref 22 words Naoki Ohmiya, Tomomitsu Tahara, Mitsuo Nagasaka, Yoshihito Nakagawa, Tomoyuki Shibata. "Chapter 12 Endoscopy"	1%
4	Crossref 18 words So Min Lee, Jung Wook Seo. "Phlebosclerotic colitis: case report and literature review focused on the radiologic findings"	<1%
5	Crossref 16 words Wei-Ching Lin, Jeon-Hor Chen, Antonio Carlos Westphalen, Chun-Han Liao, Cheng-Hong Chen, Chun-Ming Chen, C	<1%
6	Crossref 14 words Yosuke Sasaki, Manabu Saito, Yumi Koshihara, Hiroaki Zai, Yoshihisa Urita. "Idiopathic mesenteric phlebosclerosis associated with long-term oral intake of geniposide"	<1%
7	Crossref 14 words Hozumi, Hideaki, Ryota Hokari, Motonori Shimizu, Koji Maruta, Kazuyuki Narimatsu, Hirokazu Sato, Shingo Sato, Tosh	<1%
8	Crossref 13 words Fumihiko Tamamoto, Hideyuki Ishizaki, Tadayuki Maehara. "Phlebosclerotic colitis", <i>Radiation Medicine</i> , 2008	<1%
9	Crossref 12 words Tsong-Shuo Yen. "Relationship between severity of venous calcifications and symptoms of phlebosclerotic colitis", <i>World Journal of Gastroenterology</i>	<1%



ALL

IMAGES

VIDEOS

Add the Give

3,690 Results

Any time

Idiopathic mesenteric phlebosclerosis associated with long ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6322607>

Hiramatsu K, Sakata H, Horita Y, et al. **Mesenteric phlebosclerosis associated with long term oral intake of geniposide**, an ingredient of herbal medicine. Aliment Pharmacol Ther. 2012; 36 :575–86.

Cited by: 1

Author: Hsing Jung Yeh, Pao Ying Lin, Wei Yu Ka...

Publish Year: 2018

Search Tools

Turn off Hover Translation (关闭取词)

Idiopathic mesenteric phlebosclerosis associated with use ...

<https://journals.lww.com/md-journal/Fulltext/2020/...>

Idiopathic mesenteric phlebosclerosis (IMP) is a rare form of ischemic colitis. It is more common in the Asian population people with Asian ancestry. Disease pathogenesis and etiology are not fully elucidated

Idiopathic mesenteric phlebosclerosis associated with long-term oral intake of herb drug



ALL

IMAGES

VIDEOS

4,850 Results

Any time ▾

Idiopathic mesenteric phlebosclerosis associated with long-term oral intake of herb drug

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6322607>

The purple mucosa might be related to geniposide and the change of enterobacterial flora. Because genipin damages colonic mucosa, these patients have mesenteric phlebosclerosis, purple colonic mucosa, and colonic ulcers. No effective medicine for IMP, except stop using herbal medicines to avoid disease progression.

Cited by: 1

Author: Hsing Jung Yeh, Pao Ying Lin, Wei Yu K...

Publish Year: 2018

Idiopathic mesenteric phlebosclerosis associated with long-term oral intake of herb drug

<https://pubmed.ncbi.nlm.nih.gov/32565200>

Idiopathic mesenteric phlebosclerosis associated with long-term oral intake of herb drug occurring in a mother and her daughter Clin Res Hepatol Gastroenterol . 2020 Jun 18;S2210-7401(20)30160-1. doi: 10.1016/j.clinre.2020.05.015.

Author: Qiaoli Zheng, Bohui Zhang, Zhixing Ca... Publish Year: 2021

Mesenteric phlebosclerosis associated with long-term oral intake of geniposide, an ingredient of herbal medicine

<https://pubmed.ncbi.nlm.nih.gov/22817400>

Mesenteric phlebosclerosis associated with long-term oral intake of geniposide, an ingredient of herbal medicine. The long-term use of geniposide in herbal medicines appears to be associated with mesenteric phlebosclerosis.

Cited by: 70

Author: K. Hiramatsu, H. Sakata, Y. Horita, N. Ori...

Publish Year: 2012

Idiopathic mesenteric phlebosclerosis associated with use of herbal medicine

<https://journals.lww.com/md-journal/Fulltext/2020/...>

Idiopathic mesenteric phlebosclerosis (IMP) is a rare form of ischemic colitis. It is more common in the Asian population people with Asian ancestry. Disease pathogenesis and etiology are not fully elucidated but may be associated with the long-term intake of toxins ...

Mesenteric phlebosclerosis associated with long-term oral intake of herb drug

<https://onlinelibrary.wiley.com/doi/full/10.1111/j.1365-2036.2012.05221.x>

Jul 23, 2012 · At least a subgroup of cases of mesenteric phlebosclerosis may be due to oral intake

ALL

IMAGES

VIDEOS

MAPS

NEWS

SHOPPING

TOOLS ▾

Idiopathic mesenteric phlebosclerosis associated with long ...

<https://pubmed.ncbi.nlm.nih.gov/32565200>

Idiopathic mesenteric phlebosclerosis associated with long-term oral intake of herb drug occurring in a mother and her daughter Clin Res Hepatol Gastroenterol . 2020 Jun 18;S2210-7401(20)30160-1. doi: 10.1016/j.clinre.2020.05.015.

Author: Qiaoli Zheng, Bohui Zhang, Zhixing Cao,... **Publish Year:** 2021

[PDF] Mesenteric phlebosclerosis associated with long-term oral ...

<https://onlinelibrary.wiley.com/doi/epdf/10.1111/j.1365-2036.2012.05221.x>

Idiopathic mesenteric phlebosclerosis (IMP) is a rare disease, characterised by thickening of the wall of the right hemicolon with calcification of **mes-enteric** veins. However, the aetiology remains unknown. Aim To investigate the possible association of herbal medicines with IMP. Method The clinical data of four of our own patients were collected.

Mesenteric phlebosclerosis associated with long-term oral ...

<https://pubmed.ncbi.nlm.nih.gov/22817400>

Mesenteric phlebosclerosis associated with long-term oral intake of geniposide, an ingredient of herbal medicine. The **long-term use of geniposide in herbal medicines** appears to be associated with mesenteric phlebosclerosis.

Cited by: 70 **Author:** K. Hiramatsu, H. Sakata, Y. Horita, N. Orita, ...

Publish Year: 2012

Idiopathic mesenteric phlebosclerosis associated with long ...

<https://europepmc.org/articles/PMC6322607> ▾

Jan 01, 2018 · **Geniposide** could be transferred to **genipin**. **Genipin** affects right hemi-colonic bacteria flora and vascular system. The **purple** mucosa might be **related** to **geniposide** and the change of enterobacterial flora. Because **genipin** damages colonic mucosa, these patients have **mesenteric phlebosclerosis, purple** colonic mucosa, and colonic ulcers.

Cited by: 1 **Author:** Hsing Jung Yeh, Pao Ying Lin, Wei Yu Kao, ...

Publish Year: 2018

Idiopathic mesenteric phlebosclerosis associated with use ...

<https://journals.lww.com/md-journal/Fulltext/2020/...>

Idiopathic mesenteric phlebosclerosis (IMP) is a rare form of ischemic colitis. It is more common in the Asian population people with Asian ancestry. Disease pathogenesis and etiology are not fully elucidated