# World Journal of *Clinical Cases*

World J Clin Cases 2022 April 16; 10(11): 3321-3638





Published by Baishideng Publishing Group Inc

W J C C World Journal of Clinical Cases

#### Contents

#### Thrice Monthly Volume 10 Number 11 April 16, 2022

#### **REVIEW**

3321 Encouraging specific biomarkers-based therapeutic strategies for hepatocellular carcinoma Yao M, Yang JL, Wang DF, Wang L, Chen Y, Yao DF

#### **ORIGINAL ARTICLE**

#### **Clinical and Translational Research**

Autophagy-related long non-coding RNA prognostic model predicts prognosis and survival of melanoma 3334 patients

Qiu Y, Wang HT, Zheng XF, Huang X, Meng JZ, Huang JP, Wen ZP, Yao J

3352 Identification of circ\_0000375 and circ\_0011536 as novel diagnostic biomarkers of colorectal cancer Yin TF, Du SY, Zhao DY, Sun XZ, Zhou YC, Wang QQ, Zhou GYJ, Yao SK

#### **Retrospective Study**

3369 Echocardiography in the diagnosis of Shone's complex and analysis of the causes for missed diagnosis and misdiagnosis

Li YD, Meng H, Pang KJ, Li MZ, Xu N, Wang H, Li SJ, Yan J

- Predictors and prognostic impact of post-operative atrial fibrillation in patients with hip fracture surgery 3379 Bae SJ, Kwon CH, Kim TY, Chang H, Kim BS, Kim SH, Kim HJ
- 3389 Added value of systemic inflammation markers for monitoring response to neoadjuvant chemotherapy in breast cancer patients

Ke ZR, Chen W, Li MX, Wu S, Jin LT, Wang TJ

3401 Washed microbiota transplantation reduces serum uric acid levels in patients with hyperuricaemia Cai JR, Chen XW, He YJ, Wu B, Zhang M, Wu LH

#### **Clinical Trials Study**

Concurrent chemoradiotherapy using gemcitabine and nedaplatin in recurrent or locally advanced head 3414 and neck squamous cell carcinoma

Huo RX, Jin YY, Zhuo YX, Ji XT, Cui Y, Wu XJ, Wang YJ, Zhang L, Zhang WH, Cai YM, Zheng CC, Cui RX, Wang QY, Sun Z, Wang FW

#### **META-ANALYSIS**

3426 Effect of enhanced recovery after surgery on inflammatory bowel disease surgery: A meta-analysis Peng D, Cheng YX, Tao W, Tang H, Ji GY

Accuracy of ultrasound elastography for predicting breast cancer response to neoadjuvant chemotherapy: 3436 A systematic review and meta-analysis

Chen W, Fang LX, Chen HL, Zheng JH



Camban	World Journal of Clinical Cases	
Contents Thrice Monthly Volume 10 Number 11 April 16, 2022		
3449	Association of chronic obstructive pulmonary disease with mild cognitive impairment and dementia risk: A systematic review and meta-analysis	
	Zhao LY, Zhou XL	
	CASE REPORT	
3461	Circulating tumor DNA genomic profiling reveals the complicated olaparib-resistance mechanism in prostate cancer salvage therapy: A case report	
	Yuan F, Liu N, Yang MZ, Zhang XT, Luo H, Zhou H	
3472	Difference and similarity between type A interrupted aortic arch and aortic coarctation in adults: Two case reports	
	Ren SX, Zhang Q, Li PP, Wang XD	
3478	Combination therapy (toripalimab and lenvatinib)-associated toxic epidermal necrolysis in a patient with metastatic liver cancer: A case report	
	Huang KK, Han SS, He LY, Yang LL, Liang BY, Zhen QY, Zhu ZB, Zhang CY, Li HY, Lin Y	
3485	Unusual glomus tumor of the lower leg: A case report	
	Wang HY, Duan P, Chen H, Pan ZY	
3490	Pulmonary <i>Cladosporium</i> infection coexisting with subcutaneous <i>Corynespora cassiicola</i> infection in a patient: A case report	
	Wang WY, Luo HB, Hu JQ, Hong HH	
3496	Preoperational diagnosis and management of breast ductal carcinoma <i>in situ</i> arising within fibroadenoma: Two case reports	
	Wu J, Sun KW, Mo QP, Yang ZR, Chen Y, Zhong MC	
3505	Reconstruction of complex chest wall defects: A case report	
	Huang SC, Chen CY, Qiu P, Yan ZM, Chen WZ, Liang ZZ, Luo KW, Li JW, Zhang YQ, Huang BY	
3511	Young children with multidrug-resistant epilepsy and vagus nerve stimulation responding to perampanel: A case report	
	Yang H, Yu D	
3518	Intramedullary nailing for pathological fractures of the proximal humerus caused by multiple myeloma: A case report and review of literature	
	Xu GQ, Wang G, Bai XD, Wang XJ	
3527	Double tracheal stents reduce side effects of progression of malignant tracheoesophageal fistula treated with immunotherapy: A case report	
	Li CA, Yu WX, Wang LY, Zou H, Ban CJ, Wang HW	
3533	Ankylosing spondylitis complicated with andersson lesion in the lower cervical spine: A case report	
	Peng YJ, Zhou Z, Wang QL, Liu XF, Yan J	
3541	Severe gastric insufflation and consequent atelectasis caused by gas leakage using AIR-Q laryngeal mask airway: A case report	
	Zhao Y. Li P. Li DW. Zhao GF. Li XY	



Contor	World Journal of Clinical Cases	
Conten	Thrice Monthly Volume 10 Number 11 April 16, 2022	
3547	Hypereosinophilic syndrome presenting as acute ischemic stroke, myocardial infarction, and arterial involvement: A case report	
	Sun RR, Chen TZ, Meng M	
3553	Cytochrome P450 family 17 subfamily A member 1 mutation causes severe pseudohermaphroditism: A case report	
	Gong Y, Qin F, Li WJ, Li LY, He P, Zhou XJ	
3561	Patellar dislocation following distal femoral replacement after extra-articular knee resection for bone sarcoma: A case report	
	Kubota Y, Tanaka K, Hirakawa M, Iwasaki T, Kawano M, Itonaga I, Tsumura H	
3573	Qingchang decoction retention enema may induce clinical and mucosal remission in left-sided ulcerative colitis: A case report	
	Li PH, Tang Y, Wen HZ	
3579	Anti-nuclear matrix protein 2+ juvenile dermatomyositis with severe skin ulcer and infection: A case report and literature review	
	Wang YT, Zhang Y, Tang T, Luo C, Liu MY, Xu L, Wang L, Tang XM	
3587	Ultrasound-guided local ethanol injection for fertility-preserving cervical pregnancy accompanied by fetal heartbeat: Two case reports	
	Kakinuma T, Kakinuma K, Matsuda Y, Ohwada M, Yanagida K, Kaijima H	
3593	Successful apatinib treatment for advanced clear cell renal carcinoma as a first-line palliative treatment: A case report	
	Wei HP, Mao J, Hu ZL	
3601	Del(5q) and inv(3) in myelodysplastic syndrome: A rare case report	
	Liang HP, Luo XC, Zhang YL, Liu B	
3609	Papillary thyroid microcarcinoma with contralateral lymphatic skip metastasis and breast cancer: A case report	
	Ding M, Kong YH, Gu JH, Xie RL, Fei J	
3615	Contrast-enhanced ultrasound manifestations of synchronous combined hepatocellular- cholangiocarcinoma and hepatocellular carcinoma: A case report	
	Gao L, Huang JY, Lu ZJ, Lu Q	
3624	Thyrotoxicosis after a massive levothyroxine ingestion: A case report	
	Du F, Liu SW, Yang H, Duan RX, Ren WX	
3630	Pleomorphic adenoma of the left lacrimal gland recurred and transformed into myoepithelial carcinoma after multiple operations: A case report	
	Huang WP, Li LM, Gao JB	



### Contents

Thrice Monthly Volume 10 Number 11 April 16, 2022

#### **ABOUT COVER**

Editorial Board Member of World Journal of Clinical Cases, Chi-Yuan Yeh, MD, PhD, Assistant Professor, Chief Doctor, radiation oncology, Tungs' Taichung MetroHarbor Hospital, Taichung 43503, Taiwan. peteryeh46@gmail.com

#### **AIMS AND SCOPE**

The primary aim of World Journal of Clinical Cases (WJCC, World J Clin Cases) is to provide scholars and readers from various fields of clinical medicine with a platform to publish high-quality clinical research articles and communicate their research findings online.

WJCC mainly publishes articles reporting research results and findings obtained in the field of clinical medicine and covering a wide range of topics, including case control studies, retrospective cohort studies, retrospective studies, clinical trials studies, observational studies, prospective studies, randomized controlled trials, randomized clinical trials, systematic reviews, meta-analysis, and case reports.

#### **INDEXING/ABSTRACTING**

The WJCC is now indexed in Science Citation Index Expanded (also known as SciSearch®), Journal Citation Reports/Science Edition, Scopus, PubMed, and PubMed Central. The 2021 Edition of Journal Citation Reports® cites the 2020 impact factor (IF) for WJCC as 1.337; IF without journal self cites: 1.301; 5-year IF: 1.742; Journal Citation Indicator: 0.33; Ranking: 119 among 169 journals in medicine, general and internal; and Quartile category: Q3. The WJCC's CiteScore for 2020 is 0.8 and Scopus CiteScore rank 2020: General Medicine is 493/793.

#### **RESPONSIBLE EDITORS FOR THIS ISSUE**

Production Editor: Hua-Ge Yn; Production Department Director: Xiang Li; Editorial Office Director: Jin-Lei Wang.

NAME OF JOURNAL	INSTRUCTIONS TO AUTHORS
World Journal of Clinical Cases	https://www.wjgnet.com/bpg/gerinfo/204
<b>ISSN</b>	GUIDELINES FOR ETHICS DOCUMENTS
ISSN 2307-8960 (online)	https://www.wignet.com/bpg/GerInfo/287
LAUNCH DATE	GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH
April 16, 2013	https://www.wjgnet.com/bpg/gerinfo/240
FREQUENCY	PUBLICATION ETHICS
Thrice Monthly	https://www.wjgnet.com/bpg/GerInfo/288
<b>EDITORS-IN-CHIEF</b> Bao-Gan Peng, Jerzy Tadeusz Chudek, George Kontogeorgos, Maurizio Serati, Ja Hyeon Ku	PUBLICATION MISCONDUCT https://www.wjgnet.com/bpg/gerinfo/208
EDITORIAL BOARD MEMBERS	ARTICLE PROCESSING CHARGE
https://www.wjgnet.com/2307-8960/editorialboard.htm	https://www.wignet.com/bpg/gerinfo/242
PUBLICATION DATE	STEPS FOR SUBMITTING MANUSCRIPTS
April 16, 2022	https://www.wignet.com/bpg/GerInfo/239
COPYRIGHT	ONLINE SUBMISSION
© 2022 Baishideng Publishing Group Inc	https://www.f6publishing.com

© 2022 Baishideng Publishing Group Inc. All rights reserved. 7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA E-mail: bpgoffice@wjgnet.com https://www.wjgnet.com



W J C C World Journal of Clinical Cases

Submit a Manuscript: https://www.f6publishing.com

World J Clin Cases 2022 April 16; 10(11): 3630-3638

DOI: 10.12998/wjcc.v10.i11.3630

ISSN 2307-8960 (online)

CASE REPORT

# Pleomorphic adenoma of the left lacrimal gland recurred and transformed into myoepithelial carcinoma after multiple operations: A case report

#### Wen-Peng Huang, Li-Ming Li, Jian-Bo Gao

Specialty type: Medicine, research and experimental

Provenance and peer review: Unsolicited article; Externally peer reviewed.

Peer-review model: Single blind

#### Peer-review report's scientific quality classification

Grade A (Excellent): 0 Grade B (Very good): 0 Grade C (Good): C, C Grade D (Fair): 0 Grade E (Poor): 0

P-Reviewer: Ding J, China; Mohey NM, Egypt

Received: December 14, 2021 Peer-review started: December 14, 2021 First decision: January 25, 2022 Revised: February 2, 2022 Accepted: February 27, 2022 Article in press: February 27, 2022 Published online: April 16, 2022



Wen-Peng Huang, Li-Ming Li, Jian-Bo Gao, Department of Radiology, The First Affiliated Hospital of Zhengzhou University, Zhengzhou 450052, Henan Province, China

Corresponding author: Jian-Bo Gao, PhD, Academic Research, Chairman, Chief Doctor, Instructor, Department of Radiology, The First Affiliated Hospital of Zhengzhou University, No. 1 East Jianshe Road, Zhengzhou 450052, Henan Province, China. jianbogaochina@163.com

## Abstract

#### BACKGROUND

Myoepithelial carcinoma (MC) is a clinically rare malignancy, there is controversy regarding its etiology and its biological behavior is not fully elucidated. Extensive surgical resection is the main treatment method. We describe a case of pleomorphic adenoma (PA) with multiple postoperative recurrences after malignant transformation, and the history of the disease in this patient was more than 20 years. Complete resection during the first surgery of PA and long-term postoperative follow-up is necessary.

#### CASE SUMMARY

A 34-year-old male with PA and a history of 5 postoperative recurrences over 21 years, each surgically removed, presented 15 d ago with headache, nasal congestion, protrusion of the right eyeball and loss of vision in the right eye, with progressively worsening symptoms. The patient underwent surgery, and MC was confirmed by pathology examination. A small PA component was locally visible under light microscope. The patient had a recurrence of the tumor 2 mo after surgery and underwent surgical resection.

#### CONCLUSION

During the first operation for PA, care should be taken not to rupture the envelope to prevent tumor cell implantation, and when complete resection is not possible due to the anatomical site, postoperative radiotherapy is necessary to control the lesion and prevent infiltration and malignant transformation of the tumor to MC. Computed tomography and magnetic resonance imaging is important for establishing diagnosis and developing a treatment plan.

Key Words: Myoepithelial carcinoma; Pleomorphic adenoma; Magnetic resonance



imaging; Treatment; X-ray computed tomography; Case report

©The Author(s) 2022. Published by Baishideng Publishing Group Inc. All rights reserved.

**Core Tip:** This case illustrates that pleomorphic adenoma should be completely removed during the first operation to prevent capsule rupture and tumor cell implantation. When complete resection is not possible due to the anatomical site, postoperative radiotherapy should be performed to control the lesion and prevent infiltration and malignant transformation to myoepithelial carcinoma. Postoperative follow-up, especially long-term follow-up for recurrent cases, is necessary and systemic examination should be undertaken to prevent distant metastasis.

**Citation**: Huang WP, Li LM, Gao JB. Pleomorphic adenoma of the left lacrimal gland recurred and transformed into myoepithelial carcinoma after multiple operations: A case report. *World J Clin Cases* 2022; 10(11): 3630-3638

**URL:** https://www.wjgnet.com/2307-8960/full/v10/i11/3630.htm **DOI:** https://dx.doi.org/10.12998/wjcc.v10.i11.3630

#### INTRODUCTION

Myoepithelial carcinoma (MC) is a rare malignant tumor with a low incidence[1]. The mechanism of myoepithelial carcinogenesis is still controversial and the biological behavior is not yet fully elucidated. Some scholars believe that the tumor may occur independently, while some believe that the tumor is a malignant transformation from pleomorphic adenoma (PA) or benign myoepithelial tumor[2]. Here, we present a rare case of PA which recurred many times after surgery and finally transformed into MC.

#### **CASE PRESENTATION**

#### Chief complaints

A 34-year-old male patient presented to our hospital 15 d ago with headache, nasal congestion, protrusion of the right eyeball and loss of vision in the right eye, with progressively worsening symptoms.

#### History of present illness

The patient had undergone resection of a left lacrimal gland tumor 21 years ago, and postoperative pathology showed PA. The tumor recurred at 19 years, 12 years, 9 years, and 2 years, respectively, and was surgically removed. He came to our hospital for medical treatment. He had normal spirit, appetite, sleep, defecation and urination, and no weight loss.

#### History of past illness

The patient had no previous history of hypertension or heart disease, no history of diabetes mellitus or cerebrovascular allergy, no history of infectious diseases or drug allergy.

#### Personal and family history

The patient had no family history of hereditary diseases.

#### **Physical examination**

On admission, he had old surgical incisions on the left frontotemporal and periorbital areas, partial depression of the surface, muscle atrophy, protrusion of the right eye, and normal movement of the right eye in multiple directions.

#### Laboratory examinations

Laboratory test results showed that the white blood cell count was  $13.11 \times 10^{9}/L$  (normal range,  $3.50-9.50 \times 10^{9}/L$ ), lymphocyte count was  $0.80 \times 10^{9}/L$  (normal range,  $1.10-3.20 \times 10^{9}/L$ ), red blood cell count was  $2.67 \times 10^{12}/L$  (normal values range,  $4.30-5.80 \times 10^{12}/L$ ), neutrophil count was  $11.84 \times 10^{9}/L$  (normal range,  $1.80-6.30 \times 10^{9}/L$ ), and hemoglobin was 82.1 g/L (normal range, 130-175 g/L).

Zaishidena® WJCC | https://www.wjgnet.com



DOI: 10.12998/wjcc.v10.i11.3630 Copyright © The Author(s) 2022.

**Figure 1 The first preoperative Magnetic resonance imaging findings.** A-I: Magnetic resonance imaging shows multiple masses with T1 weighted imaging (WI) inhomogeneous low signals (A-D) and T2WI inhomogeneous high signals (E-H) in the bilateral frontal, bilateral septal sinuses, bilateral frontal sinuses, left orbit, left nasal cavity, left pterygoid sinus and left maxillary sinus, the lesion showed equal or slightly lower signals on diffusion WI with poorly defined borders (I); J-L: Bilateral frontal lobe and right orbital compression resulted in a localized left shift of the frontal midline, and right shift of the nasal cavity. After enhancement, the lesion showed significant heterogeneous progressive enhancement.

#### Imaging examinations

Ultrasound showed multiple solid hypoechoic masses with poorly defined borders and uneven internal echogenicity in the bilateral orbits, and dotted line blood flow signals were detected on color Doppler flow imaging. Computed tomography (CT) revealed multiple round-like soft tissue masses in the left maxillary sinus, pterygoid sinus, bilateral septal sinuses, nasal tract and bilateral intraorbital, frontal sinus, and bilateral frontal lobes, with swelling, resorption destruction and compression displacement of the surrounding adjacent bones, and anterior convexity of the right eyeball with compression, with no obvious invasion of the right optic nerve. Magnetic resonance imaging (MRI) displayed multiple masses with inhomogeneous low signals on T1 weighted imaging (T1WI) and inhomogeneous high signals on T2WI in the bilateral frontal, bilateral septal sinuses, bilateral frontal sinuses, left orbit, left nasal cavity, left pterygoid sinus and left maxillary sinus, the lesion showed an equal or slightly lower signal on diffusion WI (DWI) with poorly defined borders. Bilateral frontal lobe and right orbital compression resulted in a localized left shift of the frontal midline, and right shift of the nasal cavity. After enhancement, the lesion showed significant heterogeneous progressive enhancement (Figure 1) of approximately 7.3 cm × 7.8 cm × 8.4 cm (AP × LR × SI). Bilateral frontal dura mater was visible as linear enhancement.

Baishideng® WJCC | https://www.wjgnet.com



DOI: 10.12998/wjcc.v10.i11.3630 Copyright © The Author(s) 2022.

Figure 2 The first postoperative immunohistochemistry results. A: Hematoxylin-eosin staining shows spindle-shaped tumor cells, mucinous mesenchyme and foci of necrosis, with nuclear schizophrenia and marked nuclear anisotropy, and a small localized pleomorphic adenomatous component in the tumor (magnification × 200); B-I: Immunohistochemistry showed that the tumor cells were positive for CK (B), CK7 (C), P63 (D), Calponin (E), S-100 (F), SMA (G), SOX-10 (H) and 5% of cells were positive for Ki-67 (I) [Envision (B-I) × 200].

#### FINAL DIAGNOSIS

Postoperative pathology combined with morphology and immunohistochemistry findings were consistent with PA transformation to MC.

#### TREATMENT

The patient underwent the fifth surgical resection of the tumor. Intraoperatively, the tumor tissue was located outside the dura mater, gravish yellow and gravish red in color, with a hard texture and rich blood supply, and the tumor had an envelope, which was lobulated and locally invaded the dura mater. A small amount of cerebrospinal fluid leaked out during resection of the tumor and its envelope. The fifth postoperative pathology showed spindle-shaped tumor cells, mucinous mesenchyme and foci of necrosis, with nuclear schizophrenia and marked nuclear anisotropy, and a small localized pleomorphic adenomatous component of the tumor (Figure 2). Immunohistochemistry findings were as follows: CK (+), EMA (-), S-100 (+), SSTR-2 (-), Ki-67 (5%+), CD34 (-), CD68 (-), SMA (+), SOX -10 (+), P63 (+), Calponin (+), and CK7 (+).



Baishideng® WJCC | https://www.wjgnet.com

Huang WP et al. Myoepithelial carcinoma after multiple operations



DOI: 10.12998/wjcc.v10.i11.3630 Copyright © The Author(s) 2022.

Figure 3 The second preoperative computed tomography findings. A: Computed tomography (CT) scan shows large lamellar soft tissue density masses in the left orbit, bilateral frontal areas, bilateral septal sinuses, pterygoid sinuses, left maxillary sinus, and anterior skull base, with CT attenuation of approximately 22 Hounsfield units (HU), peripheral multiple osteolytic destruction of bone; B: Arterial phase shows moderate enhancement of the lesion, with CT attenuation of approximately 58 HU.

> After surgery, the patient was given symptomatic supportive treatment such as anti-infection, dehydration, and pain relief. The patient did not receive adjuvant therapy.

#### OUTCOME AND FOLLOW-UP

Two months later the patient developed dizziness. Combined head and neck CT angiography showed posterior displacement of the left internal carotid artery due to compression, slightly dilated lumen of the cavernous sinus segment, slender right vertebral artery, bilateral internal carotid arteries with calcium shifts and mixed plaque shadow, and moderate lumen narrowing. A large lamellar soft tissue density shadow was seen in the left orbit, bilateral frontal areas, bilateral septal sinuses, pterygoid sinus, left maxillary sinus, and anterior skull base, with CT attenuation of about 37 Hounsfield units (HU) and multiple surrounding osteolytic destruction of bone, and CT attenuation of about 58 HU in the arterial phase (Figure 3). The patient underwent MRI, which showed multiple abnormal signals in the left orbit, bilateral frontal areas, bilateral septal sinuses, pterygoid sinuses, left maxillary sinus, and anterior skull base with significant enhancement (Figure 4), and tumor recurrence was considered. The patient underwent the sixth surgical resection again, and intraoperatively, grayish-white tumor tissue was seen in the anterior skull base epidural with a general blood supply, eroding the bone of the bilateral supraorbital wall, brow arch and anterior skull base, and growing into the anterior skull base, sieve sinus and nasal cavity on both sides. The bone of the bilateral supraorbital wall was eroded by the tumor, and the bone was thin, with localized worm-like changes of bone hyperplasia and partial defects in the eroded bone. The tumor had an envelope, with relative boundaries to the surrounding tissues but abnormally tight adhesions, and separation-like changes were seen within the tumor. The tumor, part of the tumor envelope, and the bone of the anterior skull base and supraorbital wall eroded by the tumor were removed microscopically in pieces. The sixth postoperative pathology showed diffuse distribution of spindle-shaped tumor cells, and immunohistochemistry findings were as follows: CK (+), CK7 (+), P63 (+), Calponin (+), CK5/6 (+), SOX-10 (+), S-100 (+), SMA (-), Des (-), CD99 (±), ERG (-), Ki-67 (8%+), and SSTR -2 (+). A postoperative pathological diagnosis of MC recurrence was made (Figure 5). The patient is still being followed up.

#### DISCUSSION

PA is one of the most common salivary gland tumors, and is mostly located in the parotid gland, followed by the submandibular gland and sublingual gland, with easy recurrence after surgery, histologically it originates from epithelial cells with multidirectional differentiation potential and complex composition, containing mucus, adeno-duct-like structures and cartilage-like tissue, molecular characters such as TERT promoter mutation, high PD-L1 expression, and t(3;8) chromosomal



WJCC https://www.wjgnet.com



DOI: 10.12998/wjcc.v10.i11.3630 Copyright © The Author(s) 2022.

Figure 4 The second preoperative magnetic resonance imaging findings. A-H: Magnetic resonance imaging shows multiple masses of T1 weighted imaging (WI) low signals (A-D) and T2WI inhomogeneous high signals (E-H) in the left orbit, bilateral frontal areas, bilateral septal sinuses, pterygoid sinuses, left maxillary sinus, and anterior skull base; I: Slightly lower signals on diffusion WI, and lesion borders were not well defined; J-L: After enhancement, the lesion showed significant heterogeneous enhancement.

> abnormality[3,4]. Because PA has a recurrent and infiltrative envelope, it is clinically classified as a junctional tumor, *i.e.*, a tumor that is intermediate between benign and malignant. The WHO has classified malignant PAs into three categories: (1) Carcinoma ex pleomorphic adenoma (CXPA), which refers to the presence of cancerous components in an existing PA, with high invasiveness and metastasis, the origin of which is still uncertain, and the cancerous components are diverse, mostly salivary gland ductal carcinoma, non-specific adenocarcinoma, a few are MC, epithelial-MC, undifferentiated carcinoma, and adenoid cystic carcinoma[5]; (2) Carcinosarcoma (CS), in which the epithelial component is often in the form of ductal carcinoma and the mesenchymal component is in the form of chondrosarcoma-like changes; and (3) Metastasizing pleomorphic adenoma, in which the risk of malignant transformation to CXPA is about 12% in patients who develop recurrence after surgical excision of PA[6]. In particular, if the mass is long-standing and appears to increase rapidly over a short period of time, the possibility of PA malignancy to CXPA should be highly suspected, and if the tumor infiltrates the nerves and surrounding tissues, it may be accompanied by facial palsy, skin infiltration, pain, and restricted mouth opening. The long course and rapid growth of the tumor are the important features in the present case. The patient had a typical clinical course of PA appearing malignant, and recurrence may have been due to incomplete surgical resection of tumor cells breaking through the envelope to infiltrate normal tissues, or to rupture of the tumor leading to implantation of tumor cells and recurrence. This patient has undergone five surgeries and has a disease history of more than 20



Baishideng® WJCC | https://www.wjgnet.com



DOI: 10.12998/wjcc.v10.i11.3630 Copyright © The Author(s) 2022.

Figure 5 The second postoperative immunohistochemistry results. A: Hematoxylin-eosin staining shows diffuse distribution of spindle-shaped tumor cells (magnification × 200); B-I: Immunohistochemistry showed that the tumor cells were positive for CK (B), CK5/6 (C), CK7 (D), P63 (E), Calponin (F), S-100 (G), SOX-10 (H) and SSTR2 (I) [Envision (B-I) × 200].

years, with associated malignancy. Early symptoms are atypical due to the hidden anatomical location of the orbital region and clinicians should be particularly alert to this disease. Depending on the degree of infiltration of cancer cells into surrounding tissues, CXPA can be divided into three subtypes: Non-invasive, minimally invasive (invading surrounding tissues to a depth of  $\leq 1.5$  mm), and invasive (invading surrounding tissues to a depth of  $\geq 1.5$  mm), and invasive (invading surrounding tissues an invasive CXPA with tight adhesion to the surrounding tissue, and hemorrhagic necrosis and cystic changes were seen on the cut surface.

Preoperative confirmation of the diagnosis of CXPA relies on highly accurate fine needle aspiration cytopathology. However, the large size of this patient's tumor, the high number of malignant components, and the small proportion of residual PA components made the cytocentrifugal diagnosis very difficult. In this patient, the combination of mass morphology, immunohistochemical findings, and medical history was consistent with a PA carcinoma with a malignant type of MC.

In this case, the lesion was located in the eye, was large in size, had swelling and invasive growth, involved many complex structures, and had an incomplete envelope, which is now generally considered not to be a true tissue envelope, but a fibrous package formed by the reaction of the surrounding tissues caused by the tumor growth process. The density on CT was similar to that of the adjacent muscles, and the border was poorly defined, and CT clearly showed the details of the bone changes. Due to the thin structure of the orbital area tissue, this caused compression, resorption, and osteolytic destruction of the surrounding bone. MRI showed the relationship between internal tumor structures and adjacent

Raishidena® WJCC | https://www.wjgnet.com

structures. In this case, the signal was heterogeneous on T1WI and T2WI, which was related to its diverse pathological composition. Some of the lesions showed equal or slightly lower signals on DWI, which was related to the sparseness of pathological structures and the amount of interstitial stroma, and some of the lesion centers showed more hypointense or obvious T1WI low signals and T2WI high signal cystic areas. Pathology mostly demonstrated a large number of proliferating tiny arteries and trophoblastic vessels, which showed marked enhancement, and the progressive enhancement pattern may have been related to the prolonged residence time of the contrast medium and slow contouring due to mucus and adenoid-like structures in CXPA. CT can show the location, size and density of the tumor, and is the preferred examination method for showing bony lesions. MRI has better resolution of soft tissues than CT, and can clearly show the histological features of the tumor and its relationship with surrounding tissues, and is currently considered the most valuable imaging method for diagnosing ocular lesions.

Current treatment is wide surgical excision to ensure adequate safety margins, with the surgical route and approach depending on the tumor site and size. Miccio et al[8] showed that adjuvant radiotherapy prolongs the survival of patients with grade III and IV MC. Polymorphic adenomas are prone to recurrence after surgery, and the cause of recurrence is related to the first inappropriate surgical approach. Avoidance of preoperative biopsy and intraoperative compression and clamping of the tumor, together with complete excision of the mass with the envelope may reduce the recurrence rate. When rupture of the tumor envelope occurs during surgery, tumor cells can easily spread into the adjacent tissues leading to recurrence. As the lymph node metastasis rate of this tumor is not high, selective cervical lymphatic dissection can be considered in principle. In the present case, recurrence and malignancy occurred 21 years after surgery and the disease was of long duration; therefore, postoperative follow-up, especially long-term follow-up in recurrent cases, is necessary and systemic examination may be performed to prevent distant metastases. CT and MRI can show features such as lesion morphology, size, blood supply, and growth characteristics, which are important for selection of the surgical modality and follow-up review.

#### CONCLUSION

In summary, we report a patient with PA that recurred multiple times after surgery and finally transformed to MC. During the first operation for PA, care should be taken not to rupture the envelope to prevent implantation of tumor cells, and when complete resection is not possible due to the anatomical site, postoperative radiotherapy should be performed to control the lesion and prevent infiltration and malignant transformation to MC. CT and MRI can provide accurate visualization and aid decision making for the surgical approach, which is important for clarifying clinical diagnosis and developing a treatment plan.

#### FOOTNOTES

Author contributions: Huang WP contributed to the acquisition and analysis of the work, drafted the manuscript, imaging data collection and analysis; Li LM contributed to the manuscript editing; Gao JB wrote the review and editing; all authors met the requirements for authorship for the submitted version and agreed to its submission.

Informed consent statement: Participant gave written consent to participate in the study.

Conflict-of-interest statement: The authors declare that they have no competing interests.

CARE Checklist (2016) statement: The authors have read the CARE Checklist (2016), and the manuscript was prepared and revised according to the CARE Checklist (2016).

**Open-Access:** This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is noncommercial. See: https://creativecommons.org/Licenses/by-nc/4.0/

#### Country/Territory of origin: China

**ORCID** number: Wen-Peng Huang 0000-0002-9104-1494; Li-Ming Li 0000-0002-2910-9742; Jian-Bo Gao 0000-0003-2621-3701.

S-Editor: Fan JR L-Editor: A P-Editor: Fan JR



#### REFERENCES

- 1 Xu B, Mneimneh W, Torrence DE, Higgins K, Klimstra D, Ghossein R, Katabi N. Misinterpreted Myoepithelial Carcinoma of Salivary Gland: A Challenging and Potentially Significant Pitfall. Am J Surg Pathol 2019; 43: 601-609 [PMID: 30789358 DOI: 10.1097/PAS.00000000001218]
- 2 Petersson F, Chao SS, Ng SB. Anaplastic myoepithelial carcinoma of the sinonasal tract: an underrecognized salivary-type tumor among the sinonasal small round blue cell malignancies? Head Neck Pathol 2011; 5: 144-153 [PMID: 21104210 DOI: 10.1007/s12105-010-0226-y]
- Erickson LA. Pleomorphic Adenoma in the Parotid Gland. Mayo Clin Proc 2017; 92: e55-e56 [PMID: 28259235 DOI: 3 10.1016/j.mayocp.2017.01.006]
- 4 Hernandez-Prera JC, Skálová A, Franchi A, Rinaldo A, Vander Poorten V, Zbären P, Ferlito A, Wenig BM. Pleomorphic adenoma: the great mimicker of malignancy. *Histopathology* 2021; 79: 279-290 [PMID: 33368685 DOI: 10.1111/his.14322]
- Lewis JE, Olsen KD, Sebo TJ. Carcinoma ex pleomorphic adenoma: pathologic analysis of 73 cases. Hum Pathol 2001; 32: 5 596-604 [PMID: 11431714 DOI: 10.1053/hupa.2001.25000]
- Yin LX, Van Abel KM, Rooker SA, Nagelschneider AA, Olsen KD, Price DL, Janus JR, Kasperbauer JL, Moore EJ. Risk 6 factors for carcinoma ex pleomorphic adenoma in patients presenting with recurrence after resection of pleomorphic adenoma. Head Neck 2021; 43: 419-427 [PMID: 33043539 DOI: 10.1002/hed.26489]
- 7 Hu Y, Xia L, Zhang C, Xia R, Tian Z, Li J. Clinicopathologic Features and Prognostic Factors of Widely Invasive Carcinoma Ex Pleomorphic Adenoma of Parotid Gland: A Clinicopathologic Analysis of 126 Cases in a Chinese Population. J Oral Maxillofac Surg 2020; 78: 2247-2257 [PMID: 32649893 DOI: 10.1016/j.joms.2020.06.013]
- 8 Miccio JA, Oladeru OT, Yang J, Xue Y, Hoda ST, Ryu S, Stessin AM, Parker RI. Myoepithelial Carcinoma: The Role of Radiation Therapy. A Case Report and Analysis of Data From the Surveillance, Epidemiology, and End Results (SEER) Registry. J Pediatr Hematol Oncol 2016; 38: 274-278 [PMID: 26886377 DOI: 10.1097/MPH.000000000000508]





## Published by Baishideng Publishing Group Inc 7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA Telephone: +1-925-3991568 E-mail: bpgoffice@wjgnet.com Help Desk: https://www.f6publishing.com/helpdesk https://www.wjgnet.com

