

World Journal of *Clinical Cases*

World J Clin Cases 2022 May 16; 10(14): 4327-4712



OPINION REVIEW

- 4327 Emerging role of biosimilars in the clinical care of inflammatory bowel disease patients
Najeeb H, Yasmin F, Surani S

MINIREVIEWS

- 4334 Practical insights into chronic management of hepatic Wilson's disease
Lynch EN, Campani C, Innocenti T, Dragoni G, Forte P, Galli A
- 4348 Adipose-derived stem cells in the treatment of hepatobiliary diseases and sepsis
Satilmis B, Cicek GS, Cicek E, Akbulut S, Sahin TT, Yilmaz S

ORIGINAL ARTICLE

Clinical and Translational Research

- 4357 Learning curve for a surgeon in robotic pancreaticoduodenectomy through a "G"-shaped approach: A cumulative sum analysis
Wei ZG, Liang CJ, Du Y, Zhang YP, Liu Y
- 4368 Clinical and prognostic significance of expression of phosphoglycerate mutase family member 5 and Parkin in advanced colorectal cancer
Wu C, Feng ML, Jiao TW, Sun MJ

Case Control Study

- 4380 Significance of preoperative peripheral blood neutrophil-lymphocyte ratio in predicting postoperative survival in patients with multiple myeloma bone disease
Xu ZY, Yao XC, Shi XJ, Du XR

Retrospective Study

- 4395 Association between depression and malnutrition in pulmonary tuberculosis patients: A cross-sectional study
Fang XE, Chen DP, Tang LL, Mao YJ
- 4404 Pancreatic cancer incidence and mortality patterns in 2006-2015 and prediction of the epidemiological trend to 2025 in China
Yin MY, Xi LT, Liu L, Zhu JZ, Qian LJ, Xu CF
- 4414 Evaluation of short- and medium-term efficacy and complications of ultrasound-guided ablation for small liver cancer
Zhong H, Hu R, Jiang YS

- 4425** Hematopoiesis reconstitution and anti-tumor effectiveness of Pai-Neng-Da capsule in acute leukemia patients with haploidentical hematopoietic stem cell transplantation

Yuan JJ, Lu Y, Cao JJ, Pei RZ, Gao RL

- 4436** Oral and maxillofacial pain as the first sign of metastasis of an occult primary tumour: A fifteen-year retrospective study

Shan S, Liu S, Yang ZY, Wang TM, Lin ZT, Feng YL, Pakezhati S, Huang XF, Zhang L, Sun GW

- 4446** Reduced serum high-density lipoprotein cholesterol levels and aberrantly expressed cholesterol metabolism genes in colorectal cancer

Tao JH, Wang XT, Yuan W, Chen JN, Wang ZJ, Ma YB, Zhao FQ, Zhang LY, Ma J, Liu Q

Observational Study

- 4460** Correlation of pressure gradient in three hepatic veins with portal pressure gradient

Wang HY, Song QK, Yue ZD, Wang L, Fan ZH, Wu YF, Dong CB, Zhang Y, Meng MM, Zhang K, Jiang L, Ding HG, Zhang YN, Yang YP, Liu FQ

- 4470** Multi-slice spiral computed tomography in diagnosing unstable pelvic fractures in elderly and effect of less invasive stabilization

Huang JG, Zhang ZY, Li L, Liu GB, Li X

SYSTEMATIC REVIEWS

- 4480** Distribution and changes in hepatitis C virus genotype in China from 2010 to 2020

Yang J, Liu HX, Su YY, Liang ZS, Rao HY

CASE REPORT

- 4494** Bow hunter's syndrome successfully treated with a posterior surgical decompression approach: A case report and review of literature

Orlandi N, Cavallieri F, Grisendi I, Romano A, Ghadirpour R, Napoli M, Moratti C, Zanichelli M, Pascarella R, Valzania F, Zedde M

- 4502** Histological remission of eosinophilic esophagitis under asthma therapy with IL-5 receptor monoclonal antibody: A case report

Huguenot M, Bruhm AC, Essig M

- 4509** Cutaneous mucosa-associated lymphoid tissue lymphoma complicating Sjögren's syndrome: A case report and review of literature

Liu Y, Zhu J, Huang YH, Zhang QR, Zhao LL, Yu RH

- 4519** Plexiform neurofibroma of the cauda equina with follow-up of 10 years: A case report

Chomanskis Z, Juskys R, Cepkus S, Dulko J, Hendrixson V, Ruksenas O, Rocka S

- 4528** Mixed porokeratosis with a novel mevalonate kinase gene mutation: A case report

Xu HJ, Wen GD

- 4535** Isolated pancreatic injury caused by abdominal massage: A case report

Sun BL, Zhang LL, Yu WM, Tuo HF

- 4541** Bronchiolar adenoma with unusual presentation: Two case reports
Du Y, Wang ZY, Zheng Z, Li YX, Wang XY, Du R
- 4550** Periodontal-orthodontic interdisciplinary management of a “periodontally hopeless” maxillary central incisor with severe mobility: A case report and review of literature
Jiang K, Jiang LS, Li HX, Lei L
- 4563** Anesthesia management for cesarean section in a pregnant woman with odontogenic infection: A case report
Ren YL, Ma YS
- 4569** Convulsive-like movements as the first symptom of basilar artery occlusive brainstem infarction: A case report
Wang TL, Wu G, Liu SZ
- 4574** Globe luxation may prevent myopia in a child: A case report
Li Q, Xu YX
- 4580** Computer tomography-guided negative pressure drainage treatment of intrathoracic esophagojejunal anastomotic leakage: A case report
Jiang ZY, Tao GQ, Zhu YF
- 4586** Primary or metastatic lung cancer? Sebaceous carcinoma of the thigh: A case report
Wei XL, Liu Q, Zeng QL, Zhou H
- 4594** Perianesthesia emergency repair of a cut endotracheal tube’s inflatable tube: A case report
Wang TT, Wang J, Sun TT, Hou YT, Lu Y, Chen SG
- 4601** Diagnosis of cytomegalovirus encephalitis using metagenomic next-generation sequencing of blood and cerebrospinal fluid: A case report
Xu CQ, Chen XL, Zhang DS, Wang JW, Yuan H, Chen WF, Xia H, Zhang ZY, Peng FH
- 4608** Primary sigmoid squamous cell carcinoma with liver metastasis: A case report
Li XY, Teng G, Zhao X, Zhu CM
- 4617** Acute recurrent cerebral infarction caused by moyamoya disease complicated with adenomyosis: A case report
Zhang S, Zhao LM, Xue BQ, Liang H, Guo GC, Liu Y, Wu RY, Li CY
- 4625** Serum-negative Sjogren's syndrome with minimal lesion nephropathy as the initial presentation: A case report
Li CY, Li YM, Tian M
- 4632** Successful individualized endodontic treatment of severely curved root canals in a mandibular second molar: A case report
Xu LJ, Zhang JY, Huang ZH, Wang XZ

- 4640** Successful treatment in one myelodysplastic syndrome patient with primary thrombocytopenia and secondary deep vein thrombosis: A case report
Liu WB, Ma JX, Tong HX
- 4648** Diagnosis of an extremely rare case of malignant adenomyoepithelioma in pleomorphic adenoma: A case report
Zhang WT, Wang YB, Ang Y, Wang HZ, Li YX
- 4654** Management about intravesical histological transformation of prostatic mucinous carcinoma after radical prostatectomy: A case report
Bai SJ, Ma L, Luo M, Xu H, Yang L
- 4661** Hepatopulmonary metastases from papillary thyroid microcarcinoma: A case report
Yang CY, Chen XW, Tang D, Yang WJ, Mi XX, Shi JP, Du WD
- 4669** PD-1 inhibitor in combination with fruquintinib therapy for initial unresectable colorectal cancer: A case report
Zhang HQ, Huang CZ, Wu JY, Wang ZL, Shao Y, Fu Z
- 4676** Cutaneous metastasis from esophageal squamous cell carcinoma: A case report
Zhang RY, Zhu SJ, Xue P, He SQ
- 4684** Rare pattern of Maisonneuve fracture: A case report
Zhao B, Li N, Cao HB, Wang GX, He JQ
- 4691** Suprasellar cistern tuberculoma presenting as unilateral ocular motility disorder and ptosis: A case report
Zhao BB, Tian C, Fu LJ, Zhang XB
- 4698** Development of plasma cell dyscrasias in a patient with chronic myeloid leukemia: A case report
Zhang N, Jiang TD, Yi SH
- 4704** Ovarian growing teratoma syndrome with multiple metastases in the abdominal cavity and liver: A case report
Hu X, Jia Z, Zhou LX, Kakongoma N

LETTER TO THE EDITOR

- 4709** Perfectionism and mental health problems: Limitations and directions for future research
Nazari N

ABOUT COVER

Editorial Board Member of *World Journal of Clinical Cases*, Jamir Pitton Rissardo, MD, Academic Research, Adjunct Associate Professor, Research Associate, Department of Medicine, Federal University of Santa Maria, Santa Maria 97105110, Brazil. jamirrissardo@gmail.com

AIMS AND SCOPE

The primary aim of *World Journal of Clinical Cases* (WJCC, *World J Clin Cases*) is to provide scholars and readers from various fields of clinical medicine with a platform to publish high-quality clinical research articles and communicate their research findings online.

WJCC mainly publishes articles reporting research results and findings obtained in the field of clinical medicine and covering a wide range of topics, including case control studies, retrospective cohort studies, retrospective studies, clinical trials studies, observational studies, prospective studies, randomized controlled trials, randomized clinical trials, systematic reviews, meta-analysis, and case reports.

INDEXING/ABSTRACTING

The WJCC is now indexed in Science Citation Index Expanded (also known as SciSearch®), Journal Citation Reports/Science Edition, Scopus, PubMed, and PubMed Central. The 2021 Edition of Journal Citation Reports® cites the 2020 impact factor (IF) for WJCC as 1.337; IF without journal self cites: 1.301; 5-year IF: 1.742; Journal Citation Indicator: 0.33; Ranking: 119 among 169 journals in medicine, general and internal; and Quartile category: Q3. The WJCC's CiteScore for 2020 is 0.8 and Scopus CiteScore rank 2020: General Medicine is 493/793.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: Hua-Ge Yin, Production Department Director: Xu Guo, Editorial Office Director: Jin-Lei Wang.

NAME OF JOURNAL

World Journal of Clinical Cases

ISSN

ISSN 2307-8960 (online)

LAUNCH DATE

April 16, 2013

FREQUENCY

Thrice Monthly

EDITORS-IN-CHIEF

Bao-Gan Peng, Jerzy Tadeusz Chudek, George Kontogeorgos, Maurizio Serati, Ja Hyeon Ku

EDITORIAL BOARD MEMBERS

<https://www.wjgnet.com/2307-8960/editorialboard.htm>

PUBLICATION DATE

May 16, 2022

COPYRIGHT

© 2022 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjgnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>

Case Control Study

Significance of preoperative peripheral blood neutrophil-lymphocyte ratio in predicting postoperative survival in patients with multiple myeloma bone disease

Zi-Yu Xu, Xing-Chen Yao, Xiang-Jun Shi, Xin-Ru Du

Specialty type: Orthopedics**Provenance and peer review:**

Unsolicited article; Externally peer reviewed.

Peer-review model: Single blind**Peer-review report's scientific quality classification**Grade A (Excellent): 0
Grade B (Very good): 0
Grade C (Good): C, C
Grade D (Fair): D
Grade E (Poor): 0**P-Reviewer:** Arigami T, Japan;
Fonteh Fru PN, South Africa**Received:** August 29, 2021**Peer-review started:** August 29, 2021**First decision:** December 17, 2021**Revised:** December 30, 2021**Accepted:** March 25, 2022**Article in press:** March 25, 2022**Published online:** May 16, 2022**Zi-Yu Xu, Xing-Chen Yao, Xin-Ru Du**, Department of Orthopedics, Beijing Chao-Yang Hospital, Beijing 100020, China**Xiang-Jun Shi**, Department of Hematology, Beijing Chao-Yang Hospital, Beijing 100020, China**Corresponding author:** Xin-Ru Du, PhD, Professor, Department of Orthopedics, Beijing Chao-Yang Hospital, No. 8 Gongren Tiyuguannan Road, Chaoyang District, Beijing 100020, China. duxinru@163.com

Abstract

BACKGROUND

The neutrophil-lymphocyte ratio (NLR) is often used to predict a poor prognosis in patients with tumors. This study investigated the preoperative peripheral blood NLR in predicting postoperative survival (POS) in patients with multiple myeloma bone disease (MMBD).

AIM

To evaluate whether NLR can be used to predict the prognosis of MMBD patients after surgery.

METHODS

The clinical data of 82 MMBD patients who underwent surgical treatments in Beijing Chao-yang Hospital were collected. The NLR was obtained from the absolute number of neutrophils and lymphocytes, calculated by the number of neutrophils and divided by the number of lymphocytes. The peripheral blood lymphocyte percentage was used as the major marker to analyze the change in characteristics of the immune statuses of multiple myeloma patients.

RESULTS

The NLR cut-off values of $NLR \geq 3$ patients and $NLR \geq 4$ patients were significantly correlated with POS. The 3- and 5-year cumulative survival rates of the high NLR group ($NLR \geq 3$ patients) were 19.1% and 0.0%, respectively, which were lower than those of the low NLR group ($NLR < 3$ patients) (67.2% and 48.3%) ($P = 0.000$). In the high NLR group, POS (14.86 ± 14.28) was significantly shorter than that in the low NLR group (32.68 ± 21.76). Univariate analysis showed that the lymphocyte percentage 1 wk after the operation (19.33 ± 9.08)

was significantly lower than that before the operation (25.72 ± 11.02). Survival analysis showed that postoperative chemotherapy, preoperative performance status and preoperative peripheral blood NLR ≥ 3 were independent risk factors for POS.

CONCLUSION

The preoperative peripheral blood NLR can predict POS in MMBD patients. MMBD patients with a high preoperative NLR (NLR ≥ 3) showed poor prognosis.

Key Words: Multiple myeloma bone disease; Neutrophil-lymphocyte ratio; Lymphocyte percentage; Postoperative survival

©The Author(s) 2022. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: The clinical data of 82 multiple myeloma bone disease (MMBD) patients who underwent operations in our hospital were collected. By observing the change characteristics of neutrophil-lymphocyte ratio (NLR) at different stages after the operation, to make a preliminary analysis of the effect of the operation on the immune status of multiple myeloma patients and study the prognostic value of peripheral blood NLR in predicting the treatment of MMBD. Data showed that NLR was significant for the prediction of the postoperative survival (POS) of MMBD patients treated with the operation, and MMBD patients with high preoperative NLR (NLR ≥ 3) had a poorer prognosis and shorter POS.

Citation: Xu ZY, Yao XC, Shi XJ, Du XR. Significance of preoperative peripheral blood neutrophil-lymphocyte ratio in predicting postoperative survival in patients with multiple myeloma bone disease. *World J Clin Cases* 2022; 10(14): 4380-4394

URL: <https://www.wjgnet.com/2307-8960/full/v10/i14/4380.htm>

DOI: <https://dx.doi.org/10.12998/wjcc.v10.i14.4380>

INTRODUCTION

Multiple myeloma (MM) is a malignant proliferative disease of plasma cells with the second highest incidence among blood tumor diseases. Due to the growth of the aging population the incidence of MM has increased and there are many studies on prognosis, especially on prognostic factors and relevant stratification data. Such studies are based on the characteristics of patients, tumor load and cytogenetics [1], and mainly target the selection of chemotherapy regimens and other medical treatments. However, to date, only a few studies have reported MM surgical treatment prognosis stratification[2-4].

As a marker of systematic inflammation, the neutrophil-lymphocyte ratio (NLR) has been used to diagnose infectious diseases. The NLR has also been used as an immune status evaluation indicator to assess the prognosis of solid tumors, such as hepatocellular carcinoma, gastric cancer, renal cell carcinoma, colorectal cancer, ovarian cancer and prostate cancer. These studies suggested that a high NLR may predict a poor prognosis[5-7]. Although some retrospective studies have initially explored the influence of NLR on the prognosis of MM in recent years, these studies have been inadequate due to a small number of cases included in the study and shorter follow-up times[8,9]. These studies did not report on whether there was any change in the immune status of MM patients before and after the operation. Can NLR be used for the prognostic evaluation of surgically treated MM bone disease (MMBD)? Thus, the effect of the NLR on the prognosis of MM patients remains to be studied further.

This study reviewed the general conditions, indicators, surgical information and follow-up of MMBD patients from 2007 to 2017 to analyze the value of the preoperative peripheral blood NLR in predicting the prognosis of patients and its impact on postoperative survival (POS). According to the changes in the absolute neutrophil count, absolute lymphocyte count and lymphocyte percentage at different time points before and after the surgical treatments, the effect of the surgical treatments on the immune status of patients was discussed.

MATERIALS AND METHODS

General data

The institutional review board of our hospital approved this retrospective study. A total of 82 MMBD patients from the Beijing Chao-Yang Hospital were enrolled between 2007 and 2017. The sites and number of their lesions were determined according to computed tomography and magnetic resonance

imaging findings. General information of patients, including age, sex, Durie/Salmon (DS) stage, international staging system (ISS) stage, isotype, lesion site, preoperative/postoperative treatment, chemotherapy regimens, radiotherapy, operation scheme and stem cell transplantation therapy were collected.

The general physical condition of patients before surgical treatments including nutrition status, body mass index (BMI), performance status and MM activity were collected. The classification criteria of nutrition status are as follows: (1) "Well", the mucous membrane is ruddy, the skin is shiny and elastic, the subcutaneous fat is plump and elastic, the skinfold thickness is normal or enlarged, the muscle is strong, the nails and hair are moist, the intercostal space and supraclavicular fossa are moderate and the scapular and abdominal muscles are plump. Body weight and body mass index are in the normal range or slightly above normal; (2) "Poorly", the skin and mucous membrane is dry, the elasticity is reduced, the subcutaneous fat is thin, the skinfold thickness is lower than normal, the muscle is loose and weak, the nails are rough, the hair is sparse and dull, the intercostal space and supraclavicular fossa are sunken and the shoulder, ribs and iliac bones are prominent. Body weight and body mass index were significantly lower than normal; and (3) "Fairly", between the above two classification criteria.

The evaluation of the patient's performance status is based on the Zubrod-ECOG-WHO score (ZPS score, 5-point method), as follows: Grade 0 normal activity; Grade 1 symptoms but nearly ambulatory; Grade 2 some bed time but needs to be in bed less than 50% of the normal daytime; Grade 3 needs to be in bed more than 50% of the normal daytime; Grade 4 unable to get out of bed; Grade 5 death.

According to routine blood examination, the NLR was obtained using the absolute number of neutrophils and lymphocytes, calculated by the number of neutrophils divided by the number of lymphocytes. According to their NLR values, patients were grouped into a high NLR group ($\text{NLR} \geq 3$ patients) and a low NLR group ($\text{NLR} < 3$ patients). Moreover, their cumulative survival rates were calculated according to their survival 1 year, 3 years and 5 years after the operation.

Statistical analysis

Patient clinical data were analyzed by the Chi-square test and independent sample *t*-test. The risk factors that may affect the POS in patients were analyzed by Cox model analysis. The survival time was assessed by the Kaplan-Meier method and examined by the log-rank test. The chi-square test was used to analyze the 1-year, 3-year and 5-year cumulative survival rates of patients with different preoperative peripheral blood NLRs to calculate the cut-off value and select the NLR cut-off value with the most significant difference for grouping. Peripheral blood lymphocyte percentage was used as the evaluation indicator to preliminarily assess the immune status of 82 MMBD patients before and after the operation and to perform a univariate analysis on the lymphocyte percentages of the 82 postoperative patients at different stages and preliminarily assess the effect of operations on the immune status of MMBD patients. One-way ANOVA and the Mann-Whitney *U* test for further comparisons between specific group pairs were used. All data were analyzed by SPSS 19.0 statistical software (IBM SPSS Statistics 19.0). The measurement data are expressed as the mean $\pm$ SD, and $P < 0.05$ was considered as a significant difference.

RESULTS

Among the 82 MMBD patients with an average age of 60.33 ± 9.62 years, 44 patients were male (53.7%) and 38 patients were female (46.3%). The spine was involved in 56 patients (68.3%), long bone of the limb in 19 (23.2%) patients and the soft tissue in 7 (8.5%) patients. The average time from diagnosis to treatment of 82 patients was 19.48 mo (19.48 ± 26.69). Nine patients (11%) underwent complete excision, 72 (87.8%) underwent partial excision/intra-lesion curettage and 1 patient (1.2%) underwent tissue biopsy. Twenty patients (24.4%) received autologous stem cell transplantation while sixty-two patients (75.6%) did not. The average follow-up period of 82 patients was 27.03 ± 21.31 mo (range, 0.25-84 mo) (Table 1).

General physical condition before surgical treatment

The results showed that the preoperative BMI of 82 patients was 23.14 ± 3.36 . The preoperative nutrition status classification showed "Well" in 68 patients, "Fairly" in 6 patients and "Poorly" in 8 patients. The results of the preoperative performance status of 82 patients showed that 16 patients (19.5%) were grade 0-2, and 66 patients (80.5%) were grade 3-4. The data on MM activity at the time of surgery of 82 patients showed that there were 2 cases (2.4%) of partial remission, 8 cases (9.8%) of stable disease and 72 cases (87.8%) of progressive disease (Table 2).

Effect of preoperative peripheral blood NLR on the POS of MMBD patients

After statistically analyzing the correlation between the preoperative peripheral blood NLR and POS, it was observed that the NLR cut-off values of the preoperative peripheral blood $\text{NLR} \geq 3$ and $\text{NLR} \geq 4$ were significantly correlated with the postoperative cumulative survival rate (Table 3). In particular, when the NLR cut-off value was set to 3, the difference was most significant. The POS of $\text{NLR} \geq 3$

Table 1 General information of the 82 multiple myeloma bone disease patients who underwent surgical treatments

Characteristic	Value <i>n</i> (%) or <i>n</i> [SD]
Age (yr)	60.33 [9.620]
Gender	
Male	44 (53.7)
Female	38 (46.3)
ISS stage	
ISS-I	15 (18.3)
ISS-II	37 (45.1)
ISS-III	30 (36.6)
Number of lesions	
1	53 (64.6)
2	18 (22.0)
≥ 3	11 (13.4)
Site of lesion	
Spine	56 (68.3)
Long bone of limb	19 (23.2)
Soft tissue	7 (8.5)
Isotype	
IgA	23 (28)
IgD	3 (3.7)
IgG	45 (54.9)
Light chain κ/λ	11 (13.4)
Preoperative chemotherapy	
Yes	48 (58.5)
No	34 (41.5)
Postoperative chemotherapy	
Yes	71 (86.6)
No	11 (13.4)
Preoperative radiotherapy	
Yes	6 (7.3)
No	76 (92.7)
Time from a diagnosis to surgery (mo)	19.48 [26.69]
Operation scheme	
Complete excision ¹	9 (11)
Partial excision/intramedullary curettage ²	72 (87.8)
Tissue biopsy ³	1 (1.2)
Autologous stem cell transplantation	
Yes	20 (24.4)
No	62 (75.6)
POS (mo)	27.03 [21.31]
Total survival (mo)	46.63 [31.04]

¹The soft-tissue mass extended resection or partial rib resection.

²Intralesional resection and filling with polymethylmethacrylate bone cement in long bone of limb metastases or incomplete mastectomy in spinal metastases.

³Percutaneous biopsy.

SD: Standard deviation; ISS: International staging system; POS: Postoperative survival.

Table 2 General physical condition of 82 multiple myeloma bone disease patients before surgical treatments

Characteristic	Value <i>n</i> (%) or <i>n</i> [SD]
Preoperative nutrition	
Well	68 (82.9)
Fairly	6 (7.3)
Poorly	8 (9.8)
Preoperative BMI (kg/m ²)	23.14 [3.36]
Preoperative performance status	
Grade 0-2	16 (19.5)
Grade 3-4	66 (80.5)
Preoperative MM activity	
Partial remission	2 (2.4)
Stable disease	8 (9.8)
Progression disease	72 (87.8)

SD: Standard deviation; BMI: Body mass index; MM: Multiple myeloma.

Table 3 Comparison of 1-year, 3-year and 5-year cumulative survival rates between the high and the low neutrophil-lymphocyte ratio group

Cutoff value of NLR	1-yr CSR ^a	3-yr CSR ^a	5-yr CSR ^a	Chi-square	<i>P</i> value
NLR ≥ 4	41.7% vs 82.5%	10.4% vs 60.9%	0.0% vs 41.7%	15.969	0.000a
NLR ≥ 3 (<i>n</i> = 26)	52.0% vs 87.3%	19.1% vs 67.2%	0.0% vs 48.3%	22.192	0.000a
NLR ≥ 2	65.8% vs 88.9%	46.5% vs 61.1%	31.0% vs 34.0%	2.015	0.156
NLR ≥ 1	75.3% vs 85.7%	52.6% vs -	30.6% vs -	0.459	0.498

^a*P* < 0.05.

NLR: Neutrophil-lymphocyte ratio; CSR: Cumulative survival rates.

patients (*n* = 26) (14.86 ± 14.28 mo) was significantly lower than that of NLR < 3 patients (*n* = 56) (32.68 ± 21.76 mo) (*P* = 0.000) (Table 4, Figure 1). Among the 82 patients, 48 patients have received chemotherapy before surgical treatments. There were 11 (22.9%) patients in the immunomodulator-based treatment group, 20 (41.7%) patients in the proteasome inhibitor-based treatment group and 17 (35.4%) patients in the immunomodulator and proteasome inhibitor treatment groups. Seventy-one patients have received chemotherapy after surgical treatments (Table 4).

Univariate analysis showed that the preoperative BMI in the high NLR group was significantly lower than that in the low NLR group (21.74 ± 3.13 vs 23.79 ± 3.29, *P* = 0.009). There was no significant difference in preoperative chemotherapy regimens (*P* = 0.216), the time from diagnosis of MM to surgery (*P* = 0.321), the performance status (*P* = 0.066), MM activity at the time of surgery (*P* = 0.590) and postoperative chemotherapy regimens (*P* = 0.159) between the high NLR group and the low NLR group (Table 4).

The general condition of patients on preoperative peripheral blood NLR

Univariate analysis showed that the number of lesions affected NLR to some extent (*P* = 0.033), while sex, ISS stage, lesion sites, chemotherapy and autologous stem cell transplantation did not affect the


Table 4 Univariate analysis of the general information of patients in the high and the low neutrophil-lymphocyte ratio group

Characteristic	NLR < 3 (n = 56)	NLR ≥ 3 (n = 26)	P value
Age (yr)	60.93 ± 9.88	59.04 ± 9.10	0.411
Gender, n (%)			0.981
Male	30 (53.6)	14 (53.8)	
Female	26 (46.4)	12 (46.2)	
ISS stage, n (%)			0.693
ISS-I	10 (17.9)	5 (19.2)	
ISS-II	27 (48.2)	10 (38.5)	
ISS-III	19 (33.9)	11 (42.3)	
Number of lesions, n (%)			0.033 ^a
1	41 (73.2)	12 (46.2)	
2	8 (14.3)	10 (38.5)	
≥ 3	7 (12.5)	4 (15.4)	
Site of lesion, n (%)			0.982
Spine	38 (67.9)	18 (69.2)	
Long bone of limb	13 (23.2)	6 (23.1)	
Soft tissue	5 (8.9)	2 (7.7)	
Time from a diagnosis to surgery (mo)	21.18 ± 29.75	15.81 ± 18.44	0.321
Preoperative nutrition, n (%)			0.071
Well	50 (89.3)	18 (69.2)	
Fairly	3 (5.4)	3 (11.5)	
Poorly	3 (5.4)	5 (19.2)	
Preoperative BMI (kg/m ²)	23.80 ± 3.29	21.74 ± 3.13	0.009 ^a
Preoperative performance status, n (%)			0.066
Grade 0-2	14 (25.0)	2 (7.7)	
Grade 3-4	42 (75.0)	24 (92.3)	
MM activity at the time of surgery, n (%)			0.590
Partial response	2 (3.6)	0 (0.0)	
Stable disease	5 (8.9)	3 (11.5)	
Progressive disease	49 (87.5)	23 (88.5)	
Preoperative chemotherapy, n (%)			0.707
Yes	32 (57.1)	16 (61.5)	
No	24 (42.9)	10 (38.5)	
Preoperative chemotherapy regimens (n = 48), n (%)			0.216
Immunomodulator-based	6 (37.5)	5 (15.6)	
Proteasome inhibitor-based	6 (37.5)	14 (43.8)	
Both immunomodulator and proteasome inhibitor	4 (25.0)	13 (40.6)	
Postoperative chemotherapy, n (%)			0.080
Yes	51 (91.1)	20 (76.9)	
No	5 (8.9)	6 (23.1)	
Postoperative chemotherapy regimens (n = 71), n (%)			0.159
Immunomodulator-based	3 (6.0)	4 (19.0)	

Proteasome inhibitor-based	17 (34.0)	4 (19.0)	
Both immunomodulator and proteasome inhibitor	30 (60.0)	13 (61.9)	
Autologous stem cell transplantation, <i>n</i> (%)			0.458
Yes	15 (26.8)	5 (19.2)	
No	41 (73.2)	21 (80.8)	
POS (mo)	32.68 ± 21.76	14.86 ± 14.28	0.001

^a*P* < 0.05.

NLR: Neutrophil-lymphocyte ratio; POS: Postoperative survival; ISS: International staging system; BMI: Body mass index; MM: Multiple myeloma.



DOI: 10.12998/wjcc.v10.i14.4380 Copyright ©The Author(s) 2022.

Figure 1 Comparison of survival time of different neutrophil-lymphocyte ratio groups. A: Comparison of postoperative survival (POS) in patients within the neutrophil-lymphocyte ratio (NLR ≥ 1) group and NLR < 1 group, among which the POS of patients in the NLR ≥ 1 group and the NLR < 1 group showed no significant difference (*P* > 0.05); B: The POS of patients in the preoperative peripheral blood NLR ≥ 2 group and the NLR < 2 group showed no significant difference (*P* > 0.05); C: The POS of patients in the preoperative peripheral blood NLR < 3 group was significantly longer than that in the NLR ≥ 3 group (*P* < 0.05); D: The POS of patients in the preoperative peripheral blood NLR < 4 group was significantly longer than that in the NLR ≥ 4 group (*P* < 0.05). NLR: Neutrophil-lymphocyte ratio.

NLR (Table 4). Differences in preoperative Hb and aspartate aminotransferase (AST) were found between the high NLR group and the low NLR group (*P* = 0.047, *P* = 0.007), while no difference was found for PLT and pre/postoperative C-reactive protein (CRP) (Table 5).

Changes in peripheral blood absolute lymphocyte count and absolute neutrophil count before and after the operation

The absolute neutrophil count 1 wk after the operation significantly increased ($4.78 \pm 2.37 \times 10^9/L$). One month after the operation ($3.72 \pm 1.96 \times 10^9/L$), the levels returned to those before the operation (Table 6, Figure 2A and B). The peripheral blood NLR after the operation significantly increased (2.78 ± 1.97 vs 4.56 ± 2.33 , *P* = 0.002) and then declined 1 mo after the operation (3.14 ± 2.19). The levels increased again at the last follow-up visit (5.03 ± 6.73) (Table 6, Figure 2C).

When preliminarily analyzing the immune status of the 82 MMBD patients before and after the operation with the peripheral lymphocyte percentage as the evaluation indicator, the results showed

Table 5 Univariate analysis of the preoperative laboratory examination of patients in the high and the low neutrophil-lymphocyte ratio group

Preoperative laboratory examination	NLR < 3 (<i>n</i> = 56)	NLR ≥ 3 (<i>n</i> = 26)	<i>P</i> value
Hb (g/L)	112.66 ± 24.88	100.58 ± 25.91	0.047 ^a
PLT (10 ⁹ /L) ^a	208.38 ± 77.39	196.89 ± 103.64	0.577
ALb (g/L)	32.88 ± 6.02	31.16 ± 6.29	0.237
AST (U/L)	22.89 ± 8.39	33.96 ± 18.79	0.007 ^a
ALT (U/L)	19.89 ± 10.28	28.46 ± 24.17	0.093
Γ-GT (U/L)	34.11 ± 27.52	47.58 ± 38.19	0.073
Blood β2-MG (mg/L)	5.43 ± 9.08	4.10 ± 2.85	0.475
Cr (umol/L)	84.54 ± 73.73	89.38 ± 85.84	0.794
Preoperative CRP (mg/L)	2.60 ± 3.84	3.67 ± 3.93	0.411
Post-operative CRP (mg/L)	1.98 ± 2.79	1.83 ± 2.06	0.877

^a*P* < 0.05.

NLR: Neutrophil-lymphocyte ratio; CPR: C-reactive protein; AST: Aspartate aminotransferase; Hb: Hemoglobin; Alb: Albumin; ALT: Alanine transaminase; Γ-GT: Glutamyl transpeptidase; Cr: Creatinine.

Table 6 One-way ANOVA results of peripheral blood absolute lymphocyte counts, absolute neutrophil counts and neutrophil-lymphocyte ratio of the 82 multiple myeloma bone disease patients at different time points (mean ± SD)

Characteristic	Before operation	1 wk after operation	1 mo after operation	Last follow-up visit	<i>F</i>	<i>P</i> value
Absolute blood lymphocyte counts	1.34 ± 0.75	1.19 ± 0.55	1.45 ± 0.76	1.15 ± 0.82 ^{1,2}	2.382	0.070
Absolute neutrophil counts	3.87 ± 2.07	4.79 ± 2.37 ^{3,5}	3.72 ± 1.96	3.36 ± 2.04	4.376	0.005
NLR	2.78 ± 1.97 ^{6,7}	4.56 ± 2.33	3.14 ± 2.19 ⁸	5.03 ± 6.73	4.998	0.002
Lymphocyte percentages (%)	25.09 ± 9.96	18.19 ± 7.19 ^{3,4}	26.66 ± 10.70	22.83 ± 12.68	5.770	0.001

¹When that at the last follow-up visit was compared to that 1 wk after operation, *P* = 0.055.²When that at the last follow-up visit was compared to that 1 wk after operation, *P* < 0.05.³When that 1 wk after operation was compared to that before operation, *P* < 0.05.⁴When that 1 wk after operation was compared to that 1 mo after operation, *P* < 0.05.⁵When that 1 wk after operation was compared to that at the last follow-up visit, *P* < 0.05.⁶When that before operation was compared to that 1 wk after operation, *P* < 0.05.⁷When that before operation was compared to that at the last follow-up visit, *P* < 0.05.⁸When that 1 mo after operation was compared to that at the last follow-up visit, *P* < 0.05.

NLR: Neutrophil-lymphocyte ratio.

that the peripheral lymphocyte percentage of MMBD patients before operation (25.09% ± 9.96%) was at a normal (low) level; 1 wk after operation, it (18.19% ± 7.19%) significantly declined; 1 mo after operation, it (26.66% ± 10.70%) returned to that before operation; at the last follow-up visit, it (22.83% ± 12.68%) returned to a low level again (Table 6, Figure 2D).

Factors affecting the POS of MMBD patients

Kaplan-Meier analysis showed that the number of lesions, site of lesion, postoperative chemotherapy, preoperative performance status and preoperative peripheral blood NLR ≥ 3 were risk factors affecting POS (Table 7). Multivariate survival analysis showed that postoperative chemotherapy (*P* = 0.000), preoperative performance status (*P* = 0.002), and preoperative peripheral blood NLR ≥ 3 (*P* = 0.000) were independent risk factors affecting POS (Table 8).

Table 7 Kaplan–Meier analysis of postoperative survival of patients with multiple myeloma bone disease (univariate survival analysis)

Characteristic	Chi-square	P value
Gender	1.703	0.192
ISS stage	1.966	0.374
Number of lesions	7.055	0.029 ^a
Site of lesion	8.160	0.017 ^a
Isotype	1.526	0.676
Preoperative chemotherapy	0.387	0.534
Postoperative chemotherapy	35.837	0.000 ^a
Preoperative radiotherapy	0.908	0.341
Time from a diagnosis to surgery	1.841	0.606
Operation scheme	3.583	0.167
Stem cell transplantation	2.836	0.092
Preoperative peripheral blood NLR ≥ 3	22.192	0.000 ^a
Preoperative nutrition	0.782	0.676
Preoperative performance status	20.775	0.000 ^a
Preoperative MM activity	1.634	0.652

^a $P < 0.05$.

NLR: Neutrophil-lymphocyte ratio; ISS: International staging system; MM: Multiple myeloma.

Table 8 Multivariate Cox regression analysis of postoperative survival of patients with multiple myeloma bone disease

Characteristic	P value	HR	95% CI
Site of lesion	0.185	-	-
Number of lesions	0.462	-	-
Postoperative chemotherapy	0.000 ^a	0.142	0.057-0.357
Preoperative peripheral blood NLR ≥ 3	0.000 ^a	3.503	1.815-6760
Preoperative performance status	0.002 ^a	0.041	0.005-0.318


^a $P < 0.05$.

NLR: Neutrophil-lymphocyte ratio.

DISCUSSION

There is clinical value in using peripheral blood NLR in predicting the prognosis of MMBD patients treated with an operation. Multiple myeloma is a malignant proliferative disease of plasma cells, where its occurrence and development are strongly dependent on the microenvironment. The interaction between MM cells and the microenvironment has a crucial role in the pathogenesis of MM[10-12]. The MM microenvironment is mainly composed of inflammatory cells (including macrophages, dendritic cells, mast cells and myeloid-derived suppressor cells). Inflammatory cells are the main source of MM-infiltrating bone marrow cytokines and also mediate immune suppression of MM[13-16]. Inflammation is one of the characteristics of malignant tumors and tumor-related inflammation has an important role in promoting tumorigenesis by inducing tumor cell growth, angiogenesis and genomic instability[17]. As a marker of systematic inflammation, NLR has also been used as an immune status indicator for the prognosis of solid tumors[18].

Our study showed that compared with MMBD patients in the low NLR group, those in the high NLR group had a different number of lesions, preoperative hemoglobin levels, preoperative glutamate transaminase and preoperative absolute lymphocyte count. Multivariate survival analysis showed that postoperative chemotherapy and preoperative peripheral blood NLR were independent risk factors affecting the POS of MMBD patients.



DOI: 10.12998/wjcc.v10.i14.4380 Copyright ©The Author(s) 2022.

Figure 2 Comparison of white blood cell count or ratio between preoperation, 1 wk after the operation, 1 mo after the operation and the last follow-up visit in multiple myeloma bone disease patients. A: The absolute neutrophil count increased significantly 1 wk after the operation and returned to that before the operation 1 mo after the operation; B: The peripheral blood absolute lymphocyte count declined after the operation, returned to that before the operation 1 mo after the operation, and returned to a low level at the last follow-up visit; C: Peripheral blood neutrophil-lymphocyte ratio; D: Lymphocyte percentage. NLR: Neutrophil-lymphocyte ratio.

Some studies have also reported on NLR prognosis prediction in patients with hematologic tumors [19-23]. Zhou *et al*[24] found that an elevated NLR is associated with decreased overall survival in MM patients receiving bortezomib induction therapy. Wongrakpanich *et al*[25] reported that an elevated NLR and a decreased platelet-lymphocyte ratio are independent prognostic factors for progression-free survival in MM patients. Shi *et al*[26] indicated that an elevated NLR, a decreased monocyte-lymphocyte ratio and a decreased platelet-lymphocyte ratio predict adverse clinical outcomes in MM patients and may serve as cost-effective and readily available prognostic biomarkers. Lee *et al*[27] analyzed 179 MM patients treated with the VMP regimen and found that a high NLR was an independent poor prognostic factor. Solmaz *et al*[28] reviewed 150 MM patients who underwent autologous stem cell transplantation (ASCT) and found that a high NLR on the 100th d post transplantation was associated with inferior overall survival (OS) and Progression-free survival (PFS). Romano *et al*[29] showed that the median PFS was significantly shorter for MM patients with NLR ≥ 2 when considering the clinical outcomes of ASCT. In summary, MM patients have shortened OS with a high NLR.

To date, there are no reports on the prognostic value of NLR in MM patients who undergo an operation. We think that elevated NLR may be used as a prognostic biomarker for the prediction of survival and help predict clinical outcome in patients with MMBD. Before it can be used in clinical practice, the time to detect NLR and the cut-off value should be further standardized.

Change laws of peripheral blood NLR before and after the operation

Due to the great range of tumor tissue removal during surgery, the tumor load significantly decreased, and the stimulating effect of tumor cells on the immune and inflammatory reaction was reduced resulting in great changes in the NLR before and after the operation. A previous study found that the NLR of a non-small cell lung cancer patient at 1 mo after operation increased by over 0.27, and the DFS and OS of the same patient were significantly shorter than those whose NLR declined or increased by less than 0.27 thus suggesting that NLR changes before and after the operation could be used for the early evaluation of the effectiveness of tumor therapy[30].

In this study, we found that MM patients who had a high NLR before the operation, had a significantly increased NLR 1 wk after the operation. In the 2nd wk after the operation, the absolute neutrophil count significantly increased while the absolute lymphocyte count declined; in the 2nd mo, the NLR declined while the absolute neutrophil count and absolute lymphocyte count returned to levels before the operation. At the last follow-up visit, the NLR returned to a high level while the absolute neutrophil count significantly increased and the absolute lymphocyte count significantly declined. This change indicates that a high preoperative NLR, which may be the result of the imbalance of inflammatory reactions between antitumor and protumor effects, may be used to predict poor prognosis in MM patients[26]. An elevated NLR indicates that the neutrophils in an MM patient's blood increased while the lymphocytes decreased, resulting in an imbalance between the antitumor effect of lymphocytes and the protumor effect of neutrophils, thus affecting the prognosis of tumor patients.

Changes in the immune status of MMBD patients before and after the operation and their effect on prognosis

Cellular immunity and humoral immunity play a key role in the occurrence and progression of tumors. The function and status of cellular immunity largely reflects the progression of tumors and the prognosis of patients. At present, antitumor cell immunotherapy based on the correlation between cellular immunologic function and status and tumors is commonly applied in clinical practice.

In this study, we found normal levels of neutrophils in MM patients before surgery. These levels significantly increased 1 wk after the operation and then returned to normal levels 1 mo after the operation; at the last follow-up visit, the levels were lower than before the operation. We believe that 1 wk after the operation, patients were in the postoperative inflammatory reaction stage, so their neutrophils were high. However, with time, the immune status recovered, which led to a decrease in neutrophils. This may be a normal body reaction; however, whether such a reaction affects tumors needs to be further explored. At the last follow-up visit, the tumor status of patients was improved, and the level of their neutrophils declined over that before operation due to the application of treatment measures such as operations and chemotherapy. Peripheral blood NLR before and 1 mo after the operation may reflect the immune status of patients from one side and can be used as one of the indicators for predicting operative prognoses.

Correct judgment and evaluation of the cellular immunologic function of patients with malignant tumors have an important role in evaluating disease development and guiding clinical treatment. Multiple myeloma is characterized by immunologic dysfunction. MM immune injury has been considered as another tool to predict prognosis[30]. Long-term survival in MM patients may result from unique immunologic characteristics[31]. Lymphocytes are an important part of the immune system. They contain subsets of different functions, which can be roughly divided into T cells, B cells and NK cells. Domestic and foreign studies have found various defects in the immune system of MM patients, including abnormal quantities and functional defects in B cells, T cells, NK cells and DC cells, as well as abnormal regulatory Tregs[32,33]. Bernal *et al*[34] found that the number of NK cells in the peripheral blood of MM patients was increased, while their activity declined. Furthermore, Jurisic *et al*[35] proved that the decline in NK cell activity in patients was related to disease progression and the degree of plasma cell infiltration.

T cells have a major role in the antitumor immune reaction of tumor patients and regulate the immune reaction and maintain immune stability. T cell subset expression anomalies have an important role in the pathogenesis of MM. Koike *et al*[36] and Oken *et al*[37] have found a decrease in CD4(+) T/CD8(+) T cells in MM patients, which disrupts the immune balance of MM patients and weakens their existing antitumor functions. Tregs are a subset of T cells that control autoimmunity and have a negative immunoregulatory function, which maintains the immune system stability of the body together with Th1 and Th2 cells. It is related to the immune tolerance and tumor immune escape mechanisms[38,39]. Recent studies have found a lower proportion of Th1 and Th2 cells in MM patients than in healthy controls, suggesting that both humoral and cellular immunity of MM patients are impaired to varying degrees[40]. Therefore, lymphocyte subsets and Th cells are of great significance in the prognostic evaluation of MM patients. Lymphocytes have a key role in cell-mediated antitumor immune reactions and their amount reflects the degree of reaction of the host immune system.

Whether operations affect the immune status of MM patients has not yet been clarified. Our data showed that the lymphocyte percentage of MM patients was within normal (low) levels before the operation. These levels declined 1 mo after surgery and returned to normal levels later on (at the last follow-up visit), which suggests that operations affect the immune status of patients with MM. After the operation, the immune balance of MM patients is disrupted, which in turn affects the immune status. However, with time, immune status is recovered. Therefore, for MM patients who undergo surgery, adjuvant therapy to enhance their immunity may be given when necessary to allow patients to build up their immune status, which may be helpful for the overall treatment of MM patients. The correlation between such results and the timing of chemotherapy after surgery needs further study.

Neutrophils have a multivariate role in tumor diseases. However, their role in cancer has not been fully understood[41]. Neutrophils reflect the inflammatory state of tumor patients and have different roles in different stages of tumor diseases[42]. Neutrophils release reactive nitrogen species, reactive

oxygen species and proteases that promote the occurrence of tumors[43]. Nitric oxide synthase or arginase 1 released by neutrophils can inhibit the antitumor reaction of CD8(+) T lymphocytes[44] by weakening the immune system, promoting tumor proliferation, and invasion and stimulating TGF- β [45]. Moreover, some researchers believe that an increase in neutrophil count is one of the adverse prognostic factors of head and neck cancer progression[46].

Nonetheless, this study had some limitations. Due to the length of diagnosis and treatment and numerous follow-up visits, the lymphocyte subset information at different time points was not available for some patients. Second, due to different treatment regimens of patients during the treatment period, the detection of peripheral blood-related cells alone cannot fully reflect the body's immune status.

Related risk factors in MMBD patients

The lesion site, number of lesions, postoperative chemotherapy, preoperative peripheral blood NLR and preoperative performance status are the risk factors affecting POS in MMBD patients. Postoperative chemotherapy, preoperative peripheral blood NLR and preoperative performance status are independent risk factors.

Performance status, as a score to evaluate the physical activity status of patients, can reveal their general health status and tolerance to treatment based on their physical strength. For patients with poor general health before surgery, their tolerance to surgery is worse, recovery time is slower, postoperative complications are more likely to occur and the survival time is often shorter. However, as a subjective scoring system, some results do not fully respond to the true condition of patients. Comprehensive evaluation of other indicators such as BMI, muscle fat ratio and bone density is required to determine this.

Chemotherapy is another independent risk factor affecting the POS of patients with MMBD. Chemotherapy is the most important treatment method for multiple myeloma and helps to control tumors and extend the survival period. However, there is often a large disparity in treatment tolerance that affects the statistical analysis. If some patients are insensitive to chemotherapy or change chemotherapy regimens frequently, this may affect the accuracy of the statistical results. To a certain extent, NLR is an indicator of systemic inflammatory reactions which can reflect the balance between inflammation and immunity in tumor patients. We can obtain relevant values directly from the peripheral blood. However, NLR easily interferes with other factors. For patients with other infectious diseases or those who are using certain drugs, the values from the peripheral blood are not completely accurate. If the numerical difference is found in the measurement, it should be rerecorded in time.

CONCLUSION

MMBD patients with a high preoperative NLR had a poorer prognosis and shorter POS. The immune status of MMBD patients was at a normal low level and then significantly declined after surgical treatments. The NLR can predict the prognosis of MM patients undergoing surgical treatments.

ARTICLE HIGHLIGHTS

Research background

This study investigated the preoperative peripheral blood neutrophil-lymphocyte ratio (NLR) in predicting postoperative survival (POS) in patients with multiple myeloma bone disease (MMBD).

Research motivation

As a marker of systematic inflammation, NLR has been used to diagnose infectious diseases. Although some retrospective studies have initially explored the influence of NLR on the prognosis of multiple myeloma (MM) in recent years, these studies didn't report on whether there is any change in the immune status of MM patients before and after the operation.

Research objectives

To analyze the effect of preoperative peripheral blood NLR on the prognosis of MMBD patients and evaluate the immune status of MMBD patients in different time periods.

Research methods

The clinical data of 82 MMBD patients who underwent surgical treatments in Beijing Chaoyang hospital were collected.

Research results

Data showed that the NLR cut-off values of the $NLR \geq 3$ group and $NLR \geq 4$ group were significantly

correlated with POS. The POS of NLR ≥ 3 patients (14.86 ± 14.28) was significantly shorter than that of NLR < 3 patients (32.68 ± 21.76). The lymphocyte percentage 1 wk after operation (19.329 ± 9.083) was significantly lower than that before operation (25.723 ± 11.016). Survival analysis showed that postoperative chemotherapy and preoperative peripheral blood NLR were independent risk factors for POS.

Research conclusions

MMBD patients with high preoperative NLR had a poorer prognosis and shorter POS. NLR can predict the prognosis of MM patients undergoing surgical treatments.

Research perspectives

Preoperative peripheral blood NLR can predict POS in MMBD patients. The immune status of MMBD patients was at a normal low level and then it significantly declined after surgical treatments.

FOOTNOTES

Author contributions: Xu ZY drafted and revised the manuscript, and contributed to the data collection and drafting of the manuscript; Yao XC and Shi XJ contributed to the data analysis and interpretation; Du XR and Xu ZY contributed to the conception of the article and manuscript revision; all authors read and approved the final manuscript.

Supported by the Municipal Natural Science Foundation of Beijing of China, No. 7192071.

Institutional review board statement: The study was reviewed and approved by Beijing Chao-Yang Hospital Human Research Ethics Committee.

Informed consent statement: All study participants, or their legal guardian, provided informed written consent prior to study enrollment.

Conflict-of-interest statement: There are no conflicts of interest to report.

Data sharing statement: No additional data are available.

STROBE statement: The authors have read the STROBE Statement—checklist of items, and the manuscript was prepared and revised according to the STROBE Statement—checklist of items.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Country/Territory of origin: China

ORCID number: Zi-Yu Xu 0000-0002-7551-8889; Xing-Chen Yao 0000-0002-1317-3782; Xiang-Jun Shi 0000-0003-3647-1675; Xin-Ru Du 0000-0002-9694-4377.

S-Editor: Chen YL

L-Editor: Filipodia

P-Editor: Chen YL

REFERENCES

- 1 **Rajkumar SV.** Multiple myeloma: 2016 update on diagnosis, risk-stratification, and management. *Am J Hematol* 2016; **91**: 719-734 [PMID: 27291302 DOI: 10.1002/ajh.24402]
- 2 **Miller JA, Balagamwala EH, Chao ST, Emch T, Suh JH, Djemil T, Angelov L.** Spine stereotactic radiosurgery for the treatment of multiple myeloma. *J Neurosurg Spine* 2017; **26**: 282-290 [PMID: 27791828 DOI: 10.3171/2016.8.SPINE16412]
- 3 **Surgeon's Committee of the Chinese Myeloma Working Group of the International Myeloma Foundation.** Consensus on Surgical Management of Myeloma Bone Disease. *Orthop Surg* 2016; **8**: 263-269 [PMID: 27627707 DOI: 10.1111/os.12267]
- 4 **Yao X, Xu Z, Du X.** PKP/PVP combine chemotherapy in the treatment of multiple myeloma patients with vertebral pathological fractures: minimum 3-year follow-up of 108 cases. *J Orthop Surg Res* 2019; **14**: 42 [PMID: 30744644 DOI: 10.1186/s13018-019-1078-0]

- 5 **Diem S**, Schmid S, Krapf M, Flatz L, Born D, Jochum W, Templeton AJ, Früh M. Neutrophil-to-Lymphocyte ratio (NLR) and Platelet-to-Lymphocyte ratio (PLR) as prognostic markers in patients with non-small cell lung cancer (NSCLC) treated with nivolumab. *Lung Cancer* 2017; **111**: 176-181 [PMID: 28838390 DOI: 10.1016/j.lungcan.2017.07.024]
- 6 **Sun S**, Wang X, Chen J. Using Pre-Treatment Neutrophil-to-Lymphocyte Ratio to Predict the Prognosis of Young Patients with Hepatocellular Carcinoma Implemented Minimally Invasive Treatment. *J Adolesc Young Adult Oncol* 2020; **9**: 85-89 [PMID: 31621472 DOI: 10.1089/jayao.2019.0046]
- 7 **Luvira V**, Kamsa-Ard S, Pugkhem A, Luvira V, Srisuk T, Titapun A, Silsirivanit A, Wongkham S, Khuntikeo N, Pairojkul C, Bhudhisawasdi V. Predictive utility of neutrophil-to-lymphocyte ratio and platelet-to-lymphocyte ratio in intraductal papillary neoplasm of the bile duct. *Clin Exp Hepatol* 2019; **5**: 250-255 [PMID: 31598563 DOI: 10.5114/ceh.2019.87641]
- 8 **Zeng Q**, Liu Z, Li Q, Liu T. Prognostic value of neutrophil to lymphocyte ratio and clinicopathological characteristics for multiple myeloma: A meta-analysis. *Medicine (Baltimore)* 2018; **97**: e12678 [PMID: 30313061 DOI: 10.1097/MD.00000000000012678]
- 9 **Liu S**, Shi J, Guo H, Xu F, Wei M, Sun K, Chen Y. Prognostic Significance Of The Inflammatory Index-Based Scoring System In Patients Preliminarily Diagnosed With Multiple Myeloma In The Bortezomib-Based Chemotherapy Era. *Cancer Manag Res* 2019; **11**: 9409-9420 [PMID: 31807070 DOI: 10.2147/CMAR.S227671]
- 10 **Kumar S**. Multiple myeloma - current issues and controversies. *Cancer Treat Rev* 2010; **36** Suppl 2: S3-11 [PMID: 20472186 DOI: 10.1016/S0305-7372(10)70006-2]
- 11 **Spicka I**. Advances in multiple myeloma therapy during two past decades. *Comput Struct Biotechnol J* 2014; **10**: 38-40 [PMID: 25210597 DOI: 10.1016/j.csbj.2014.05.005]
- 12 **Palumbo A**, Anderson K. Multiple myeloma. *N Engl J Med* 2011; **364**: 1046-1060 [PMID: 21410373 DOI: 10.1056/NEJMra1011442]
- 13 **Sun Y**, Pan J, Mao S, Jin J. IL-17/miR-192/IL-17Rs regulatory feedback loop facilitates multiple myeloma progression. *PLoS One* 2014; **9**: e114647 [PMID: 25489847 DOI: 10.1371/journal.pone.0114647]
- 14 **Chauhan D**, Singh AV, Brahmandam M, Carrasco R, Bandi M, Hideshima T, Bianchi G, Podar K, Tai YT, Mitsiades C, Raje N, Jaye DL, Kumar SK, Richardson P, Munshi N, Anderson KC. Functional interaction of plasmacytoid dendritic cells with multiple myeloma cells: a therapeutic target. *Cancer Cell* 2009; **16**: 309-323 [PMID: 19800576 DOI: 10.1016/j.ccr.2009.08.019]
- 15 **Tai YT**, Anderson KC. B cell maturation antigen (BCMA)-based immunotherapy for multiple myeloma. *Expert Opin Biol Ther* 2019; **19**: 1143-1156 [PMID: 31277554 DOI: 10.1080/14712598.2019.1641196]
- 16 **Görgün GT**, Whitehill G, Anderson JL, Hideshima T, Maguire C, Laubach J, Raje N, Munshi NC, Richardson PG, Anderson KC. Tumor-promoting immune-suppressive myeloid-derived suppressor cells in the multiple myeloma microenvironment in humans. *Blood* 2013; **121**: 2975-2987 [PMID: 23321256 DOI: 10.1182/blood-2012-08-448548]
- 17 **Musolino C**, Allegra A, Innaro V, Allegra AG, Pioggia G, Gangemi S. Inflammatory and Anti-Inflammatory Equilibrium, Proliferative and Antiproliferative Balance: The Role of Cytokines in Multiple Myeloma. *Mediators Inflamm* 2017; **2017**: 1852517 [PMID: 29089667 DOI: 10.1155/2017/1852517]
- 18 **Cummings M**, Merone L, Keeble C, Burland L, Grzelinski M, Sutton K, Begum N, Thacoar A, Green B, Sarveswaran J, Hutson R, Orsi NM. Preoperative neutrophil: lymphocyte and platelet: lymphocyte ratios predict endometrial cancer survival. *Br J Cancer* 2015; **113**: 311-320 [PMID: 26079303 DOI: 10.1038/bjc.2015.200]
- 19 **Luo G**, Guo M, Liu Z, Xiao Z, Jin K, Long J, Liu L, Liu C, Xu J, Ni Q, Yu X. Blood neutrophil-lymphocyte ratio predicts survival in patients with advanced pancreatic cancer treated with chemotherapy. *Ann Surg Oncol* 2015; **22**: 670-676 [PMID: 25155401 DOI: 10.1245/s10434-014-4021-y]
- 20 **Guthrie GJ**, Charles KA, Roxburgh CS, Horgan PG, McMillan DC, Clarke SJ. The systemic inflammation-based neutrophil-lymphocyte ratio: experience in patients with cancer. *Crit Rev Oncol Hematol* 2013; **88**: 218-230 [PMID: 23602134 DOI: 10.1016/j.critrevonc.2013.03.010]
- 21 **Feng JF**, Liu JS, Huang Y. Lymphopenia predicts poor prognosis in patients with esophageal squamous cell carcinoma. *Medicine (Baltimore)* 2014; **93**: e257 [PMID: 25501097 DOI: 10.1097/MD.0000000000000257]
- 22 **Kokcu A**, Kurtoglu E, Celik H, Tosun M, Malatyalioglu E, Ozdemir AZ. May the platelet to lymphocyte ratio be a prognostic factor for epithelial ovarian cancer? *Asian Pac J Cancer Prev* 2014; **15**: 9781-9784 [PMID: 25520104 DOI: 10.7314/apjcp.2014.15.22.9781]
- 23 **Gungorduk K**, Ertas IE, Ozdemir A, Akkaya E, Telli E, Taskin S, Gokcu M, Guzel AB, Oge T, Akman L, Toptas T, Solmaz U, Dogan A, Terek MC, Sancı M, Ozsaran A, Simsek T, Vardar MA, Yalcin OT, Ozalp S, Yildirim Y, Ortac F. Prognostic Significance of Retroperitoneal Lymphadenectomy, Preoperative Neutrophil Lymphocyte Ratio and Platelet Lymphocyte Ratio in Primary Fallopian Tube Carcinoma: A Multicenter Study. *Cancer Res Treat* 2015; **47**: 480-488 [PMID: 25622588 DOI: 10.4143/crt.2014.058]
- 24 **Zhou X**, Wang J, Xia J, Cheng F, Mao J, Zhu J, Guo H. Evaluation of neutrophil-to-lymphocyte ratio in newly diagnosed patients receiving borte- zomib-based therapy for multiple myeloma. *Cancer Biomark* 2018; **22**: 43-48 [PMID: 29562497 DOI: 10.3233/CBM-170795]
- 25 **Wongrakpanich S**, George G, Chaiwatcharayut W, Bisio S, Candelario N, Mittal V, Pomerantz S, Varadi G. The Prognostic Significance of Neutrophil-to-Lymphocyte and Platelet-to-Lymphocyte Ratios in Patients With Multiple Myeloma. *J Clin Lab Anal* 2016; **30**: 1208-1213 [PMID: 27239981 DOI: 10.1002/jcla.22004]
- 26 **Shi L**, Qin X, Wang H, Xia Y, Li Y, Chen X, Shang L, Tai YT, Feng X, Acharya P, Acharya C, Xu Y, Deng S, Hao M, Zou D, Zhao Y, Ru K, Qiu L, An G. Elevated neutrophil-to-lymphocyte ratio and monocyte-to-lymphocyte ratio and decreased platelet-to-lymphocyte ratio are associated with poor prognosis in multiple myeloma. *Oncotarget* 2017; **8**: 18792-18801 [PMID: 27852046 DOI: 10.18632/oncotarget.13320]
- 27 **Uz B**. The Prognostic Value of the Derived Neutrophil-to-Lymphocyte Ratio in Transplantation-Ineligible Patients with Multiple Myeloma. *Acta Haematol* 2018; **140**: 157-158 [PMID: 30253407 DOI: 10.1159/000491988]
- 28 **Solmaz Medeni S**, Acar C, Olgun A, Acar A, Seyhanlı A, Taskiran E, Sevinçik OG, Alacacioglu I, Piskin O, Ozcan MA, Demirkan F, Undar B, Ozsan GH. Can Neutrophil-to-Lymphocyte Ratio, Monocyte-to-Lymphocyte Ratio, and Platelet-to-Lymphocyte Ratio at Day +100 be used as a prognostic marker in Multiple Myeloma patients with autologous

- transplantation? *Clin Transplant* 2018; **32**: e13359 [PMID: 30053318 DOI: 10.1111/ctr.13359]
- 29 **Romano A**, Parrinello NL, Consoli ML, Marchionni L, Forte S, Conticello C, Pompa A, Corso A, Milone G, Di Raimondo F, Borrello I. Neutrophil to lymphocyte ratio (NLR) improves the risk assessment of ISS staging in newly diagnosed MM patients treated upfront with novel agents. *Ann Hematol* 2015; **94**: 1875-1883 [PMID: 26223359 DOI: 10.1007/s00277-015-2462-4]
- 30 **Pessoa de Magalhães RJ**, Vidriales MB, Paiva B, Fernandez-Gimenez C, García-Sanz R, Mateos MV, Gutierrez NC, Lecrevisse Q, Blanco JF, Hernández J, de las Heras N, Martínez-López J, Roig M, Costa ES, Ocio EM, Perez-Andres M, Maiolino A, Nucci M, De La Rubia J, Lahuerta JJ, San-Miguel JF, Orfao A; Spanish Myeloma Group (GEM); Grupo Castellano-Leones de Gammopatías Monoclonales, cooperative study groups. Analysis of the immune system of multiple myeloma patients achieving long-term disease control by multidimensional flow cytometry. *Haematologica* 2013; **98**: 79-86 [PMID: 22773604 DOI: 10.3324/haematol.2012.067272]
- 31 **Bryant C**, Suen H, Brown R, Yang S, Favalaro J, Aklilu E, Gibson J, Ho PJ, Iland H, Fromm P, Woodland N, Nassif N, Hart D, Joshua DE. Long-term survival in multiple myeloma is associated with a distinct immunological profile, which includes proliferative cytotoxic T-cell clones and a favourable Treg/Th17 balance. *Blood Cancer J* 2013; **3**: e148 [PMID: 24036947 DOI: 10.1038/bcj.2013.34]
- 32 **Feyler S**, von Lilienfeld-Toal M, Jarmin S, Marles L, Rawstron A, Ashcroft AJ, Owen RG, Selby PJ, Cook G. CD4(+)CD25(+)FoxP3(+) regulatory T cells are increased whilst CD3(+)CD4(-)CD8(-)alpha-betaTCR(+) Double Negative T cells are decreased in the peripheral blood of patients with multiple myeloma which correlates with disease burden. *Br J Haematol* 2009; **144**: 686-695 [PMID: 19133978 DOI: 10.1111/j.1365-2141.2008.07530.x]
- 33 **Pratt G**, Goodyear O, Moss P. Immunodeficiency and immunotherapy in multiple myeloma. *Br J Haematol* 2007; **138**: 563-579 [PMID: 17686051 DOI: 10.1111/j.1365-2141.2007.06705.x]
- 34 **Bernal M**, Garrido P, Jiménez P, Carretero R, Almagro M, López P, Navarro P, Garrido F, Ruiz-Cabello F. Changes in activatory and inhibitory natural killer (NK) receptors may induce progression to multiple myeloma: implications for tumor evasion of T and NK cells. *Hum Immunol* 2009; **70**: 854-857 [PMID: 19580833 DOI: 10.1016/j.humimm.2009.07.004]
- 35 **Jurisić V**, Srdić T, Konjević G, Marković O, Colović M. Clinical stage-depending decrease of NK cell activity in multiple myeloma patients. *Med Oncol* 2007; **24**: 312-317 [PMID: 17873307 DOI: 10.1007/s12032-007-0007-y]
- 36 **Koike M**, Sekigawa I, Okada M, Matsumoto M, Iida N, Hashimoto H, Oshimi K. Relationship between CD4(+)CD8(+) T cell ratio and T cell activation in multiple myeloma: reference to IL-16. *Leuk Res* 2002; **26**: 705-711 [PMID: 12191564 DOI: 10.1016/s0145-2126(01)00192-8]
- 37 **Oken MM**, Kay NE. T-cell subpopulations in multiple myeloma: correlation with clinical disease status. *Br J Haematol* 1981; **49**: 629-634 [PMID: 6976183 DOI: 10.1111/j.1365-2141.1981.tb07273.x]
- 38 **Tu JF**, Ding YH, Ying XH, Wu FZ, Zhou XM, Zhang DK, Zou H, Ji JS. Regulatory T cells, especially ICOS⁺FOXP3⁺ regulatory T cells, are increased in the hepatocellular carcinoma microenvironment and predict reduced survival. *Sci Rep* 2016; **6**: 35056 [PMID: 27725696 DOI: 10.1038/srep35056]
- 39 **Fan L**, Strasser-Weippl K, Li JJ, St Louis J, Finkelstein DM, Yu KD, Chen WQ, Shao ZM, Goss PE. Breast cancer in China. *Lancet Oncol* 2014; **15**: e279-e289 [PMID: 24872111 DOI: 10.1016/S1470-2045(13)70567-9]
- 40 **Beyer M**, Kochanek M, Giese T, Endl E, Weihrach MR, Knolle PA, Classen S, Schultze JL. In vivo peripheral expansion of naive CD4⁺CD25^{high} FoxP3⁺ regulatory T cells in patients with multiple myeloma. *Blood* 2006; **107**: 3940-3949 [PMID: 16410445 DOI: 10.1182/blood-2005-09-3671]
- 41 **Grecian R**, Whyte MKB, Walmsley SR. The role of neutrophils in cancer. *Br Med Bull* 2018; **128**: 5-14 [PMID: 30137312 DOI: 10.1093/bmb/ldy029]
- 42 **Coffelt SB**, Wellenstein MD, de Visser KE. Neutrophils in cancer: neutral no more. *Nat Rev Cancer* 2016; **16**: 431-446 [PMID: 27282249 DOI: 10.1038/nrc.2016.52]
- 43 **Jamieson T**, Clarke M, Steele CW, Samuel MS, Neumann J, Jung A, Huels D, Olson MF, Das S, Nibbs RJ, Sansom OJ. Inhibition of CXCR2 profoundly suppresses inflammation-driven and spontaneous tumorigenesis. *J Clin Invest* 2012; **122**: 3127-3144 [PMID: 22922255 DOI: 10.1172/JCI61067]
- 44 **Bodogai M**, Moritoh K, Lee-Chang C, Hollander CM, Sherman-Baust CA, Wersto RP, Araki Y, Miyoshi I, Yang L, Trinchieri G, Biragyn A. Immunosuppressive and Prometastatic Functions of Myeloid-Derived Suppressive Cells Rely upon Education from Tumor-Associated B Cells. *Cancer Res* 2015; **75**: 3456-3465 [PMID: 26183924 DOI: 10.1158/0008-5472.CAN-14-3077]
- 45 **Spiegel A**, Brooks MW, Houshyar S, Reinhardt F, Ardolino M, Fessler E, Chen MB, Krall JA, DeCock J, Zervantonakis IK, Iannello A, Iwamoto Y, Cortez-Retamozo V, Kamm RD, Pittet MJ, Raulet DH, Weinberg RA. Neutrophils Suppress Intraluminal NK Cell-Mediated Tumor Cell Clearance and Enhance Extravasation of Disseminated Carcinoma Cells. *Cancer Discov* 2016; **6**: 630-649 [PMID: 27072748 DOI: 10.1158/2159-8290.CD-15-1157]
- 46 **Valero C**, Pardo L, López M, García J, Camacho M, Quer M, León X. Pretreatment count of peripheral neutrophils, monocytes, and lymphocytes as independent prognostic factor in patients with head and neck cancer. *Head Neck* 2017; **39**: 219-226 [PMID: 27534525 DOI: 10.1002/hed.24561]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

Telephone: +1-925-3991568

E-mail: bpgoffice@wjgnet.com

Help Desk: <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

