

World Journal of *Clinical Cases*

World J Clin Cases 2022 May 26; 10(15): 4713-5123



EDITORIAL

- 4713 Diet and intestinal bacterial overgrowth: Is there evidence?
Souza C, Rocha R, Cotrim HP

MINIREVIEWS

- 4717 Definition and classification of acute-on-chronic liver diseases
Zhang YY, Meng ZJ
- 4726 Management of neurosurgical patients during coronavirus disease 2019 pandemics: The Ljubljana, Slovenia experience
Velnar T, Bosnjak R

ORIGINAL ARTICLE**Clinical and Translational Research**

- 4737 Glycolytic and fatty acid oxidation genes affect the treatment and prognosis of liver cancer
Zou JY, Huang YJ, He J, Tang ZX, Qin L
- 4761 Detection of a novel panel of 24 genes with high frequencies of mutation in gastric cancer based on next-generation sequencing
Zeng HH, Yang Z, Qiu YB, Bashir S, Li Y, Xu M

Case Control Study

- 4776 Outcomes of cervical degenerative disc disease treated by anterior cervical discectomy and fusion with self-locking fusion cage
Zhang B, Jiang YZ, Song QP, An Y
- 4785 Impact of COVID-19 pandemic on clinicopathological features of transplant recipients with hepatocellular carcinoma: A case-control study
Akbulut S, Sahin TT, Ince V, Yilmaz S

Retrospective Study

- 4799 Risk factors and optimal predictive scoring system of mortality for children with acute paraquat poisoning
Song Y, Wang H, Tao YH
- 4810 Application effect of thoracoscopic tricuspid valvuloplasty in geriatric patients with tricuspid valve disease
Jiang W, Long XM, Wei KQ, Li SC, Zhang Z, He BF, Li H
- 4818 Endoscopic ultrasonography in the evaluation of condition and prognosis of ulcerative colitis
Jin RF, Chen YM, Chen RP, Ye HJ

- 4827 Dynamic interaction nursing intervention on functional rehabilitation and self-care ability of patients after aneurysm surgery

Xie YE, Huang WC, Li YP, Deng JH, Huang JT

Clinical Trials Study

- 4836 Validations of new cut-offs for surgical drains management and use of computerized tomography scan after pancreatoduodenectomy: The DALCUT trial

Caputo D, Coppola A, La Vaccara V, Passa R, Carbone L, Ciccozzi M, Angeletti S, Coppola R

Observational Study

- 4843 Psychosocial adaptation and influencing factors among patients with chemotherapy-induced peripheral neuropathy

Zhou X, Wang DY, Ding CY, Liu H, Sun ZQ

META-ANALYSIS

- 4856 Outcome of the efficacy of Chinese herbal medicine for functional constipation: A systematic review and meta-analysis

Lyu Z, Fan Y, Bai Y, Liu T, Zhong LL, Liang HF

CASE REPORT

- 4878 Familial gastrointestinal stromal tumors with *KIT* germline mutation in a Chinese family: A case report

Yuan W, Huang W, Ren L, Xu C, Luan LJ, Huang J, Xue AW, Fang Y, Gao XD, Shen KT, Lv JH, Hou YY

- 4886 Nonfunctional pancreatic neuroendocrine tumours misdiagnosed as autoimmune pancreatitis: A case report and review of literature

Lin ZQ, Li X, Yang Y, Wang Y, Zhang XY, Zhang XX, Guo J

- 4895 Sudden deafness as a prodrome of cerebellar artery infarction: Three case reports

Li BL, Xu JY, Lin S

- 4904 Importance of abdominal X-ray to confirm the position of levonorgestrel-releasing intrauterine system: A case report

Maebayashi A, Kato K, Hayashi N, Nagaishi M, Kawana K

- 4911 Bedside ultrasonic localization of the nasogastric tube in a patient with severe COVID-19: A case report

Zhu XJ, Liu SX, Li QT, Jiang YJ

- 4917 Paradoxical herniation after decompressive craniectomy provoked by mannitol: A case report

Du C, Tang HJ, Fan SM

- 4923 Targeted next-generation sequencing identifies a novel nonsense mutation in ANK1 for hereditary spherocytosis: A case report

Fu P, Jiao YY, Chen K, Shao JB, Liao XL, Yang JW, Jiang SY

- 4929 Nonfunctional bladder paraganglioma misdiagnosed as hemangioma: A case report

Chen J, Yang HF

- 4935** Special type of Wernekink syndrome in midbrain infarction: Four case reports
Yang YZ, Hu WX, Zhai HJ
- 4942** Primary extraskeletal Ewing's sarcoma of the lumbar nerve root: A case report
Lei LH, Li F, Wu T
- 4949** Yellow nail syndrome accompanied by minimal-change nephrotic syndrome: A case report
Zhang YN, Wang MH, Yu WC, Cheng W, Cong JP, Huang XP, Wang FF
- 4957** Total femur replacement with 18 years of follow-up: A case report
Yang YH, Chen JX, Chen QY, Wang Y, Zhou YB, Wang HW, Yuan T, Sun HP, Xie L, Yao ZH, Yang ZZ
- 4964** Male metaplastic breast cancer with poor prognosis: A case report
Kim HY, Lee S, Kim DI, Jung CS, Kim JY, Nam KJ, Choo KS, Jung YJ
- 4971** CD8-positive indolent T-Cell lymphoproliferative disorder of the gastrointestinal tract: A case report and review of literature
Weng CY, Ye C, Fan YH, Lv B, Zhang CL, Li M
- 4985** Bone flare after initiation of novel hormonal therapy in patients with metastatic hormone-sensitive prostate cancer: A case report
Li KH, Du YC, Yang DY, Yu XY, Zhang XP, Li YX, Qiao L
- 4991** Postoperative infection of the skull base surgical site due to suppurative parotitis: A case report
Zhao Y, Zhao Y, Zhang LQ, Feng GD
- 4998** Blunt aortic injury-traumatic aortic isthmus pseudoaneurysm with right iliac artery dissection aneurysm: A case report
Fang XX, Wu XH, Chen XF
- 5005** Extensive complex thoracoabdominal aortic aneurysm salvaged by surgical graft providing landing zone for endovascular graft: A case report
Jang AY, Oh PC, Kang JM, Park CH, Kang WC
- 5012** Gastric heterotopia of colon found cancer workup in liver abscess: A case report
Park JG, Suh JI, Kim YU
- 5018** Clinical manifestations and gene analysis of Hutchinson-Gilford progeria syndrome: A case report
Zhang SL, Lin SZ, Zhou YQ, Wang WQ, Li JY, Wang C, Pang QM
- 5025** Neurocutaneous melanosis with an intracranial cystic-solid meningeal melanoma in an adult: A case report and review of literature
Liu BC, Wang YB, Liu Z, Jiao Y, Zhang XF
- 5036** Metastasis of liver cancer to the thyroid after surgery: A case report
Zhong HC, Sun ZW, Cao GH, Zhao W, Ma K, Zhang BY, Feng YJ

- 5042** Spontaneous liver rupture following SARS-CoV-2 infection in late pregnancy: A case report
Ambrož R, Stašek M, Molnár J, Špička P, Klos D, Hambálek J, Skanderová D
- 5051** Carotid blowout syndrome caused by chronic infection: A case report
Xie TH, Zhao WJ, Li XL, Hou Y, Wang X, Zhang J, An XH, Liu LT
- 5057** Is repeat wide excision plus radiotherapy of localized rectal melanoma another choice before abdominoperineal resection? A case report
Chiu HT, Pu TW, Yen H, Liu T, Wen CC
- 5064** Metaplastic breast cancer with chondrosarcomatous differentiation combined with concurrent bilateral breast cancer: A case report
Yang SY, Li Y, Nie JY, Yang ST, Yang XJ, Wang MH, Zhang J
- 5072** Rare solitary splenic metastasis from a thymic carcinoma detected on fluorodeoxyglucose-positron emission tomography: A case report
Tsai YH, Lin KH, Huang TW
- 5077** Type A aortic dissection following heart transplantation: A case report
Zeng Z, Yang LJ, Zhang C, Xu F
- 5082** Catheter-related infections caused by *Mycobacterium abscessus* in a patient with motor neurone disease: A case report
Pan SF, Zhang YY, Wang XZ, Sun JJ, Song SL, Tang YR, Wang JL
- 5088** Clear aligner treatment for a four-year-old patient with anterior cross-bite and facial asymmetry: A case report
Zou YR, Gan ZQ, Zhao LX
- 5097** Knot impingement after arthroscopic rotator cuff repair mimicking infection: A case report
Kim DH, Jeon JH, Choi BC, Cho CH
- 5103** Solitary primary pulmonary synovial sarcoma: A case report
He WW, Huang ZX, Wang WJ, Li YL, Xia QY, Qiu YB, Shi Y, Sun HM
- 5111** Anesthetic management for intraoperative acute pulmonary embolism during inferior vena cava tumor thrombus surgery: A case report
Hsu PY, Wu EB
- 5119** Delayed diagnosis of arytenoid cartilage dislocation after tracheal intubation in the intensive care unit: A case report
Yan WQ, Li C, Chen Z

ABOUT COVER

Editorial Board Member of *World Journal of Clinical Cases*, Jing Yang, MD, Associate Professor, Department of the First General Surgery, Gansu Provincial Hospital, Lanzhou 730000, Gansu Province, China. 21634604@qq.com

AIMS AND SCOPE

The primary aim of *World Journal of Clinical Cases* (*WJCC*, *World J Clin Cases*) is to provide scholars and readers from various fields of clinical medicine with a platform to publish high-quality clinical research articles and communicate their research findings online.

WJCC mainly publishes articles reporting research results and findings obtained in the field of clinical medicine and covering a wide range of topics, including case control studies, retrospective cohort studies, retrospective studies, clinical trials studies, observational studies, prospective studies, randomized controlled trials, randomized clinical trials, systematic reviews, meta-analysis, and case reports.

INDEXING/ABSTRACTING

The *WJCC* is now indexed in Science Citation Index Expanded (also known as SciSearch®), Journal Citation Reports/Science Edition, Scopus, PubMed, and PubMed Central. The 2021 Edition of Journal Citation Reports® cites the 2020 impact factor (IF) for *WJCC* as 1.337; IF without journal self cites: 1.301; 5-year IF: 1.742; Journal Citation Indicator: 0.33; Ranking: 119 among 169 journals in medicine, general and internal; and Quartile category: Q3. The *WJCC*'s CiteScore for 2020 is 0.8 and Scopus CiteScore rank 2020: General Medicine is 493/793.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: *Ying-Yi Yuan*; Production Department Director: *Xiang Li*; Editorial Office Director: *Jin-Lei Wang*.

NAME OF JOURNAL

World Journal of Clinical Cases

ISSN

ISSN 2307-8960 (online)

LAUNCH DATE

April 16, 2013

FREQUENCY

Thrice Monthly

EDITORS-IN-CHIEF

Bao-Gan Peng, Jerzy Tadeusz Chudek, George Kontogeorgos, Maurizio Serati, Ja Hyeon Ku

EDITORIAL BOARD MEMBERS

<https://www.wjgnet.com/2307-8960/editorialboard.htm>

PUBLICATION DATE

May 26, 2022

COPYRIGHT

© 2022 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gcrinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gcrinfo/240>

PUBLICATION ETHICS

<https://www.wjgnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gcrinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gcrinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>

Anesthetic management for intraoperative acute pulmonary embolism during inferior vena cava tumor thrombus surgery: A case report

Pei-Yu Hsu, En-Bo Wu

Specialty type: Medicine, research and experimental

Provenance and peer review: Unsolicited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's scientific quality classification

Grade A (Excellent): 0
Grade B (Very good): B
Grade C (Good): C
Grade D (Fair): D
Grade E (Poor): 0

P-Reviewer: Gao L, China; Hui P, China; Oley MH, Indonesia

Received: January 6, 2022

Peer-review started: January 6, 2022

First decision: February 21, 2022

Revised: March 1, 2022

Accepted: March 27, 2022

Article in press: March 27, 2022

Published online: May 26, 2022



Pei-Yu Hsu, Department of Anesthesiology, China Medical University Hospital, China Medical University, Taichung 404, Taiwan

En-Bo Wu, Department of Anesthesiology, Kaohsiung Chang Gung Memorial Hospital and Chang Gung University College of Medicine, Kaohsiung 833, Taiwan

Corresponding author: En-Bo Wu, MD, Attending Doctor, Department of Anesthesiology, Kaohsiung Chang Gung Memorial Hospital and Chang Gung University College of Medicine, No. 123 Ta-Pei Road, Niao-Song District, Kaohsiung 833, Taiwan. enbofive@gmail.com

Abstract

BACKGROUND

Acute pulmonary embolism (APE) is a rare and potentially life-threatening condition, even with early detection and prompt management. Intraoperative APE required specific ways for detecting since classic symptoms of APE in the awake patient could not be observed or self-reported by the patient under general anesthesia.

CASE SUMMARY

A 44-year-old man with a history of hepatic cell carcinoma was admitted for radical nephrectomy and tumor thrombectomy due to a newly found kidney tumor with inferior vena cava (IVC) tumor thrombus. APE that occurred during tumor thrombectomy with hypercapnia and desaturation. The capnography combined with the transesophageal echocardiography (TEE) provided a crucial differential diagnosis during the operation. The patient was continuously managed with aggressive intravenous fluid resuscitation and blood transfusion under continuous cardiac output monitoring to maintain hemodynamic stability. He completed the surgery under stable hemodynamics and was extubated after percutaneous mechanical thrombectomy by a certified cardiologist. There were no significant symptoms and signs or obvious discomfort in the patient's self-report during visits to the general ward.

CONCLUSION

Under general anesthesia for IVC tumor thrombus surgery, a sudden decrease in end-tidal carbon dioxide is the initial indicator of APE, which occurs before hemodynamic changes. When intraoperative APE is suspected, TEE is useful in the diagnosis and monitoring before computer tomography pulmonary

angiogram. Timely clinical impression and supportive treatment and intervention should be conducted to obtain a better prognosis.

Key Words: Acute pulmonary embolism; Anesthesia; End-tidal carbon dioxide; Inferior vena cava tumor thrombus; Transesophageal echocardiography; Case report

©The Author(s) 2022. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: Intraoperative acute pulmonary embolism (APE) is a potentially life-threatening condition, while early detection with prompt management may help improve prognosis. An acute decrease in end-tidal carbon dioxide is an early sign indicating APE. When intraoperative APE is suspected, transesophageal echocardiography is useful in the diagnosis and monitoring before computer tomography pulmonary angiogram. Besides rapid diagnosis, further management and supportive treatment should also be considered to save the patient's life.

Citation: Hsu PY, Wu EB. Anesthetic management for intraoperative acute pulmonary embolism during inferior vena cava tumor thrombus surgery: A case report. *World J Clin Cases* 2022; 10(15): 5111-5118

URL: <https://www.wjgnet.com/2307-8960/full/v10/i15/5111.htm>

DOI: <https://dx.doi.org/10.12998/wjcc.v10.i15.5111>

INTRODUCTION

Acute pulmonary embolism (APE) is a rare and potentially life-threatening condition and causes more than 10000 deaths annually worldwide[1]. The mortality is approximately 25% with massive pulmonary embolism without cardiopulmonary arrest, much higher with the presence of cardiopulmonary arrest [2]. The severity of pulmonary embolism is related to the degree of obstruction of the pulmonary artery's (PA) flow by the emboli. Rapid progression of hemodynamic instability due to PA occlusion requires immediate management. Early and precise detection is the most important strategy when APE occurs. Classic symptoms of APE in the awake patient, such as dyspnea and pleuritic chest pain, could not be observed or self-reported by the patient under general anesthesia[3]. Thus, there should be some specific ways for detecting intraoperative APE, different from the general conditions. Although there are guidelines and management suggestions for APE, few of them discuss APE that occurred intraoperatively under general anesthesia[1,3]. This report describes the early signs and anesthetic management of intraoperative APE.


CASE PRESENTATION

Chief complaints

A 44-year-old male presented to the department of urology with a 4 mo history of mild flank pain.

History of present illness

The patient had received abdominal magnetic resonance imaging, which revealed a focal lesion with infiltrative margin involving the left renal parenchyma, renal pelvis, and renal vein and extending to the inferior vena cava (IVC) (Figure 1). According to his history of stage IVB hepatic cell carcinoma (HCC), metastasis from previous HCC could not be excluded. Therefore, the origin of the tumor may come from clear cell type of renal cell carcinoma (RCC) or metastasize from previous HCC. There were two reasons which made us consider the tumor may be derived from clear cell type of RCC. First, HCC and RCC share similar MRI image features. Both of them have intracellular lipid that results in decreased signal on out-of-phase image as compared with in-phase images through the MRI. Second, the kidney is uncommon as a site for distant metastases of HCC[4]. However, there was no pathology report before surgery. Because of the abundant blood flow to the kidneys, preoperative biopsy may be complicated by massive bleeding. After discussing the benefits and complications with the patient, the patient declined preoperative biopsy and opted for direct surgery. Preoperative electrocardiography and chest radiography revealed no obvious dysfunction or lesion. Spirometry demonstrated early obstructive ventilatory defects with FEV1/FVC of 76% and FEV1 of 3.63 L (FEV1, forced expiratory volume in 1 s; FVC, forced vital capacity). The functional capacity is quantified in five metabolic equivalents of task. A sudden decrease in end-tidal carbon dioxide (ETCO₂) was found during surgery.



DOI: 10.12998/wjcc.v10.i15.5111 Copyright ©The Author(s) 2022.

Figure 1 Abdominal magnetic resonance imaging for preoperative evaluation. A: The inferior vena cava (IVC) tumor thrombus was at intrahepatic level, which was graded as level II (extends into the IVC, > 2 cm above the renal vein but below the hepatic veins; indicated by yellow arrow); B: Left renal tumor with renal vein invasion (indicated by yellow arrow).

History of past illness

The patient had hypertension, which was poorly controlled due to irregular medication use. He had a history of hepatectomy and lung resection under general anesthesia due to HCC with lung metastasis. The TNM stage of the patient's HCC was T₁N₀M₁, stage IVB, and the Child-Pugh classification was A. The prothrombin time (PT) was 10.0 s with an INR of 0.98, and the activated partial thromboplastin time (aPTT) was 26.8 s. Both PT and aPTT were within the normal limits.

Personal and family history

There were no allergy history and no smoking or drinking history. No relevant family medical history was present.

Physical examination

The airway was examined by a certified anesthesiologist and evaluated as Mallampati class II, thyromental distance < 6 cm, inter-incisor gap of 5 cm, and full neck extension. Clear and equal bilateral breath sounds was confirmed by direct auscultation with a stethoscope.

Laboratory examinations

Preoperative blood and biochemical tests showed no abnormal data. During the surgery, however, an increasing gradient of ETCO₂ to arterial partial pressure of carbon dioxide (PaCO₂) was found by intraoperative arterial gas analysis. The results were as follows: PaCO₂, 47.5 mmHg; arterial partial pressure of oxygen (PaO₂), 133 mmHg; and arterial oxygen saturation, 100%. In addition, his preoperative laboratory test values were within normal range.

Imaging examinations

A sudden decrease in ETCO₂ from 33 mmHg to 18 mmHg was found. In addition to the decrease in ETCO₂, oscillation of the plateau phase (phase III) of the capnography was also observed (Figure 2). After ETCO₂ decrease, the SpO₂ suddenly decreased from 98% to 92%. There was a tumor thrombus found at the IVC in the middle-esophageal bicaval view of transesophageal echocardiography (TEE) (Figure 3).

FINAL DIAGNOSIS

After 6.5 h, the surgery was completed under acceptable hemodynamic status. The patient underwent emergent chest computer tomography pulmonary angiogram (CTPA), which showed filling defects in bilateral pulmonary arteries during the arterial phase, and diagnosis of PE was confirmed (Figure 4).


TREATMENT

Pulmonary angiography performed by the cardiologist revealed bilateral PA occlusive thrombus with impaired flow. Thus, the patient received catheter-directed thrombolysis and systemic anticoagulation for treatment of PE. Thrombolytic agent with recombinant tissue plasminogen activator, also known as r-tPA, and urokinase under EkoS™ Endovascular System for directed thrombolysis. Systemic antico-




DOI: 10.12998/wjcc.v10.i15.5111 Copyright ©The Author(s) 2022.

Figure 2 Intraoperative capnography when acute pulmonary embolism was suspected. Oscillating of the plateau phase (phase III, indicated by yellow arrow) of capnography was observed in each breath.



DOI: 10.12998/wjcc.v10.i15.5111 Copyright ©The Author(s) 2022.

Figure 3 Intraoperative transesophageal echocardiography. Middle-esophageal bicaval view of transesophageal echocardiography showed tumor thrombus in inferior vena cava (indicated by yellow arrow).



DOI: 10.12998/wjcc.v10.i15.5111 Copyright ©The Author(s) 2022.

Figure 4 Postoperative chest computer tomography pulmonary angiogram. Filling defects was observed in bilateral pulmonary arteries (indicated by yellow arrow).

agulation with heparin was administered, and aPTT was maintained more than 1.5-2.0 times the normal value. However, increased wound drainage and unstable vital signs with hypotension, tachycardia, and decreased urine output noted after heparinization. Thrombolytic therapy and anticoagulant therapy should be discontinued due to active bleeding. Then, the patient received emergent percutaneous

mechanical thrombectomy (PMT). Right PA thrombus resolution was noted from pulmonary angiography before recanalization of the left PA after PMT. The IVC filter was also placed at infrarenal IVC for prevention of further deep venous thrombosis thereafter, and the anticoagulant was not suitable due to active bleeding.

OUTCOME AND FOLLOW-UP

The patient was weaned from the ventilator on postoperative day 4, and the endotracheal tube was removed. Under stable conditions, the patient was transferred to the general ward smoothly. In the general ward, a smooth respiratory pattern was observed, and there was no obvious discomfort in the patient's self-report. After recovery, the patient was discharged on postoperative day 13 uneventfully. The patient regularly followed at the oncology and cardiology outpatient department, and the IVC filter was removed after 2 mo.

DISCUSSION

The IVC tumor thrombus occurs in approximately 10% of patients who were diagnosed with RCC[5]. It also could be noted in the patient with advanced stage of HCC[6,7]. Patients with RCC with IVC tumor thrombus underwent radical nephrectomy and thrombectomy as a mainstay treatment strategy[8]. However, HCC and extrahepatic spread were considered as extremely advanced diseases with poor prognosis. Thus, conservative treatment with target therapy was often recommended[9]. Judging from the abovementioned evidence, there are indeed two different interventions to the same clinical finding before the pathological sample is determined.

In our case, the origin of the renal tumor from metastatic HCC was proved by the certified pathologist postoperatively, but could not be judged by the preoperative image study. Looking back, the patient should receive conservative treatment instead of surgical intervention according to the current guideline[9]. While the success of the surgery had extended the patient's life expectancy since he still regularly follows up at our outpatient department for HCC, it had been almost two years after the surgery. A recent study has reported that surgical resection with IVC tumor thrombectomy in a patient with HCC showed benefits in long-term survival[10]. Our patient seems to be one of the proofs, although further investigations are still needed.

Patients with IVC tumor thrombus are under high risks of pulmonary embolism caused by the debris of tumor thrombus, and tumor thrombus also impairs the patency of IVC, leading to venous stasis, resulting in increased thromboembolic risks[5]. According to Virchow's triad, endothelial injury and venous stasis increase the risk of thrombosis. During the course of treatment in our case, the emboli located in the right PA disappeared after thrombolysis, suggesting a thrombosis rather than tumor emboli. But left PA emboli still exist after thrombolysis, requiring PMT, suggesting that it should be tumor emboli. Based on the above reasons, IVC thrombus of this case seems to be composed of both tumor tissue and thrombus. IVC filters are often used as a prophylactic treatment for patients with venous thrombosis to prevent pulmonary embolism[11]. It is not suitable for our patient to undergo implantation preoperatively as prophylaxis. In the process of IVC filter placement, the wire needs to bypass the tumor thrombus, which highly increases the risk of tumor thrombus rupture, leading to APE and distant metastasis from the tumor debrides[12].

PE is a fatal pathology that is difficult to diagnose without expensive imaging techniques. The nonspecific clinical manifestations of PE challenge its identification, and an erroneous diagnostic approach is detrimental and related to recurrent thromboembolism, bleeding, or even death[13]. The gold standard for diagnosis of PE is a CTPA[14]. It still remains unclear whether the intraoperative capnography could provide a correct diagnosis before CT. In patients with IVC tumor thrombus undergoing surgery, there is a high risk of intraoperative pulmonary embolism. Although vital sign changes are the most common signs of APE, a decline in $ETCO_2$ could be an early sign of APE[15].

Occlusion of the flow of pulmonary arteries by emboli leads to alveolar dead space and ventilation-perfusion mismatch. Dysfunction of gas exchange in the alveoli ends up with increasing pressure gradient from $ETCO_2$ to $PaCO_2$. Phase I of capnography is air from the airway, and phase II includes the transition to alveolar air. Phase III of capnography is composed of alveolar air. PE would flatten the slope of phase III[13], and the oscillation of phase III of capnography was also observed in our case (Figure 2). Early detection and intervention with early resuscitation of vital signs could help improve the prognosis[16]. This phenomenon was also observed in another case, which was diagnosed with air and fat embolism[13,17]. Moreover, Wiegand *et al* demonstrated that capnography can be effectively applied to the noninvasive monitoring of thrombolytic therapy of APE, as a tool to monitor the improvement of lung perfusion[18].

Intraoperative TEE is crucial in the decision-making process in both surgical and anesthetic management. TEE also evaluates mobility of tumor thrombus, whether it is fragile or adherent to the IVC[19]. Direct visualization of the thrombus from TEE confirms the diagnosis. Other findings include

dilatation of right ventricle (RV) with impaired systolic function, bowing of the interventricular septum, and small and hyperdynamic left ventricle[20]. Real-time TEE monitoring can be proceeded by the anesthesiologists without suspension of the surgical interventions and exhibit the sudden disappearance of the tumor, presence of the thrombus from the IVC to the pulmonary arteries, or any sign of right heart dysfunction[21]. Moreover, intraoperative TEE assessment provides real-time information and promptly rules out other emergent pathologies, including aortic dissection, aortic aneurysm, cardiac tamponade, and regional wall motion abnormality when intraoperative hemodynamic instability occurs.

Once APE is detected, cardiopulmonary support is important to stabilize the patient's condition. Supplemental oxygenation should be provided to prevent hypoxemia. Ventilation strategy should avoid high airway pressure and high positive end-expiratory pressure in order not to increase RV loading[22]. Ventilator settings should be adjusted according to blood gas analysis to prevent hypercapnia and respiratory acidosis. Severe acidosis may lead to electrolyte imbalance, such as hyperkalemia, which may increase the risk of fatal arrhythmia and hemodynamic instability. Mechanical obstruction of PA blood flow caused vasoconstrictor release, resulting in both pulmonary vascular resistance and PA pressure increase, both of which increases the right ventricular afterload[23]. Inotropic agents, such as dobutamine, could be used for RV support to prevent RV failure, but it also causes peripheral vessel dilatation. Norepinephrine can also be administered when hypotension occurs since it has both vasoconstrictive and β 1-inotropic effects[22]. Furthermore, elevated blood pressure could enhance perfusion to the RV. Besides pharmacological support, fluid resuscitation may also be helpful in maintaining hemodynamic stability. However, volume status should be evaluated delicately, and high blood volume increases the right heart burden, which may result in RV failure[24]. In this case, combined with intraoperative TEE monitoring, evaluation of continuous cardiac output and stroke volume variation by FloTrac/Vigileo™ system and central venous pressure provided excellent evaluation of intravascular fluid status. If the patient failed to maintain relative stable hemodynamics under inotropic agents, veno-arterial ECMO (V-A ECMO) should be initiated to provide both hemodynamic and respiratory support as a rescue or bridge therapy before PA reperfusion[25].

After the diagnosis was confirmed by CTPA, the target of treatment is to restore the PA blood flow. Noninvasive treatment with systemic anticoagulants is presently viewed as current standard therapy and should be initiated immediately by the American College of Chest Physicians (ACCP) recommendations[26]. Intravenous unfractionated heparin or low-molecular-weight heparins were often used. If unfractionated heparin was used, aPTT should be checked every 6 h to evaluate the efficacy, and the value should be targeted about two times to three times of normal level[27]. A select group of direct oral anticoagulants was recently approved by the US Food and Drug Administration as another treatment for APE[27]. For patients with massive or submassive pulmonary embolism, systemic thrombolysis should be considered for reperfusion of the major pulmonary arteries. Apart from systemic thrombolysis, thrombolytic agent administered directly from a catheter to the thrombus is called catheter-directed thrombolysis[22]. Localized administration of thrombolytic agent reduced total systemic thrombolytic dose, which minimizes the side effect of bleeding and also improves the efficacy. This is presently recommended in patients with high risks of bleeding or patients who had failed from systemic thrombolysis by ACCP guidelines. However, for patients who could not tolerate thrombolytic therapy or those with hemodynamic instability who need rapid removal of the embolism, direct thrombectomy or embolectomy would be more useful[23].

PMT is a device system combined with a catheter and complex suction device that enables extraction of the large emboli. Although the advantage of PMT is preventing the risk of bleeding, it is more invasive and required more technical threshold. The last means of reperfusion is surgical pulmonary embolectomy, which should be performed under cardiopulmonary bypass. Direct emboli are removed after PA incision, which is often performed in patients with great size thrombi, proximal thrombi, intracardiac thrombi, or paradoxical emboli[28]. In our patient, reperfusion therapy was started with catheter-directed thrombolysis since a recent major surgery is contraindicated to systemic thrombolysis. However, bleeding with hemodynamic instability still occurred after catheter-directed thrombolysis. Thus, treatment for intraoperative APE should prevent bleeding.

Our study has two limitations. First, this is an observational case report. Our patient underwent surgery for HCC with distant metastases, which is not recommended in the current protocol. Prognosis of this case appears to show better outcomes compared to current life expectancy in advanced HCC. However, more evidence is needed to demonstrate the effectiveness of the treatment. Second, there is no standardized anesthesia protocol for IVC tumor thrombus surgery. APE during surgery in high-risk patients can lead to life-threatening conditions. We hope to see a comprehensive approach to anesthesia management in the future.

CONCLUSION

Intraoperative APE is a potentially life-threatening condition, while early detection with prompt management may help improve prognosis. An acute decrease in $ETCO_2$ is an early sign indicating APE.

When intraoperative APE is suspected, TEE is useful in the diagnosis and monitoring before CTPA. Besides rapid diagnosis, further management and supportive treatment should also be considered to save the patient's life.

ACKNOWLEDGEMENTS

We thank Ms. Yuan WC, a certified registered nurse anesthetist, for her help in the clinical work. The authors would like to thank the anonymous reviewers and editor for their comments. Finally, we are grateful to the patient, who gave his informed consent for publication.

FOOTNOTES

Author contributions: Hsu PY collected clinical data, discussed the details, contributed to the draft writing; Wu EB modified the manuscript and was a major contributor in writing the manuscript. Both authors read and approved the final manuscript.

Informed consent statement: Informed written consent was obtained from the patient for publication of this report and any accompanying images.

Conflict-of-interest statement: The authors have no conflicts of interest to declare.

CARE Checklist (2016) statement: The authors have read the CARE Checklist (2016), and the manuscript was prepared and revised according to the CARE Checklist (2016).

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Country/Territory of origin: Taiwan

ORCID number: Pei-Yu Hsu 0000-0003-0401-8892; En-Bo Wu 0000-0001-9581-571X.

S-Editor: Wang LL

L-Editor: A

P-Editor: Wang LL

REFERENCES

- Martinez Licha CR, McCurdy CM, Maldonado SM, Lee LS. Current Management of Acute Pulmonary Embolism. *Ann Thorac Cardiovasc Surg* 2020; **26**: 65-71 [PMID: 31588070 DOI: 10.5761/ates.ra.19-00158]
- Choi JH, O'Malley TJ, Maynes EJ, Weber MP, D'Antonio ND, Mellado M, West FM, Galanis T, Gonsalves CF, Marhefka GD, Awsare BK, Merli GJ, Tchanchaleishvili V. Surgical Pulmonary Embolectomy Outcomes for Acute Pulmonary Embolism. *Ann Thorac Surg* 2020; **110**: 1072-1080 [PMID: 32151576 DOI: 10.1016/j.athoracsur.2020.01.075]
- Cormican D, Morkos MS, Winter D, Rodrigue MF, Wendel J, Ramakrishna H. Acute Perioperative Pulmonary Embolism-Management Strategies and Outcomes. *J Cardiothorac Vasc Anesth* 2020; **34**: 1972-1984 [PMID: 31883768 DOI: 10.1053/j.jvca.2019.11.018]
- D'Antonio A, Caleo A, Caleo O, Adesso M, Boscaino A. Hepatocellular carcinoma metastatic to the kidney mimicking renal oncocytoma. *Hepatobiliary Pancreat Dis Int* 2010; **9**: 550-552 [PMID: 20943467]
- Fukazawa K, Fong CT, Gologorsky E. Inferior Vena Cava Tumor Thrombus Dynamics and Perioperative Pulmonary Embolism: A Single-Center Experience. *J Cardiothorac Vasc Anesth* 2019; **33**: 2728-2734 [PMID: 31072702 DOI: 10.1053/j.jvca.2019.03.011]
- Ariizumi SI, Kikuchi C, Tokitou F, Yamashita S, Kotera Y, Omori A, Kato T, Nemoto S, Niinami H, Yamamoto M. Cavoatrial thrombectomy prior to hepatectomy for hepatocellular carcinoma with tumor thrombus in the right atrium: a case report. *Surg Case Rep* 2019; **5**: 57 [PMID: 30972512 DOI: 10.1186/s40792-019-0620-y]
- Qiu W, Wang C, Zhang R, Wei F, Shi X, Sun X, Ma D, Lv G, Wang G. Surgical treatment of a rare case of hepatocellular carcinoma with right atrial metastasis: A case report. *Medicine (Baltimore)* 2020; **99**: e21630 [PMID: 32769926 DOI: 10.1097/MD.00000000000021630]
- Yazici S, Inci K, Bilen CY, Gudeloglu A, Akdogan B, Ertoy D, Kaynaroglu V, Demircin M, Ozen H. Renal cell carcinoma with inferior vena cava thrombus: the Hacettepe experience. *Urol Oncol* 2010; **28**: 603-9 [PMID: 19171489 DOI: 10.1016/j.urolonc.2008.11.017]
- Vogel A, Martinelli E; ESMO Guidelines Committee. Electronic address: clinicalguidelines@esmo.org; ESMO Guidelines

- Committee. Updated treatment recommendations for hepatocellular carcinoma (HCC) from the ESMO Clinical Practice Guidelines. *Ann Oncol* 2021; **32**: 801-805 [PMID: 33716105 DOI: 10.1016/j.annonc.2021.02.014]
- 10 **Bai Y**, Wu J, Zeng Y, Chen J, Wang S, Chen S, Qiu F, Zhou S, You S, Tian Y, Wang Y, Yan M. Nomogram for Predicting Long-Term Survival after Synchronous Resection for Hepatocellular Carcinoma and Inferior Vena Cava Tumor Thrombosis: A Multicenter Retrospective Study. *J Oncol* 2020; **2020**: 3264079 [PMID: 32322268 DOI: 10.1155/2020/3264079]
 - 11 **Piazza G**. Advanced Management of Intermediate- and High-Risk Pulmonary Embolism: JACC Focus Seminar. *J Am Coll Cardiol* 2020; **76**: 2117-2127 [PMID: 33121720 DOI: 10.1016/j.jacc.2020.05.028]
 - 12 **Marron RM**, Rali P, Hountras P, Bull TM. Inferior Vena Cava Filters: Past, Present, and Future. *Chest* 2020; **158**: 2579-2589 [PMID: 32795479 DOI: 10.1016/j.chest.2020.08.002]
 - 13 **Manara A**, D'hoore W, Thys F. Capnography as a diagnostic tool for pulmonary embolism: a meta-analysis. *Ann Emerg Med* 2013; **62**: 584-91 [PMID: 23769645 DOI: 10.1016/j.annemergmed.2013.04.010]
 - 14 **Enea I**, Bongarzoni A, Vedovati MC, Vatrano M, Misuraca L, Picariello C, Roncon L, Gabrielli D, Gulizia MM. Linee guida ESC 2019 per la diagnosi e la gestione dell'embolia polmonare: quali sono le novità? *G Ital Cardiol (Rome)* 2020; **21**: 175-178 [PMID: 32100729 DOI: 10.1714/3306.32765]
 - 15 **Lupei MI**, Kloesel B, Trillos L, Apostolidou I. Survival of intraoperative massive pulmonary embolism using alteplase and VA-ECMO. *J Clin Anesth* 2019; **57**: 112 [PMID: 30965272 DOI: 10.1016/j.jclinane.2019.03.007]
 - 16 **Cox JC**, Jablons DM. Operative and Perioperative Pulmonary Emboli. *Thorac Surg Clin* 2015; **25**: 289-99 [PMID: 26210925 DOI: 10.1016/j.thorsurg.2015.04.010]
 - 17 **Peña W**, Cárdenas-Camarena L, Bayter-Marin JE, McCormick M, Durán H, Ramos-Gallardo G, Robles-Cervantes JA, Macías AA. Macro Fat Embolism After Gluteal Augmentation With Fat: First Survival Case Report. *Aesthet Surg J* 2019; **39**: NP380-NP383 [PMID: 31102410 DOI: 10.1093/asj/sjz151]
 - 18 **Wiegand UK**, Kurowski V, Giannitsis E, Katus HA, Djonlagic H. Effectiveness of end-tidal carbon dioxide tension for monitoring thrombolytic therapy in acute pulmonary embolism. *Crit Care Med* 2000; **28**: 3588-92 [PMID: 11098958 DOI: 10.1097/00003246-200011000-00003]
 - 19 **Koide Y**, Mizoguchi T, Ishii K, Okumura F. Intraoperative management for removal of tumor thrombus in the inferior vena cava or the right atrium with multiplane transesophageal echocardiography. *J Cardiovasc Surg (Torino)* 1998; **39**: 641-7 [PMID: 9833725]
 - 20 **Jelic T**, Baimel M, Chenkin J. Bedside Identification of Massive Pulmonary Embolism with Point-of-Care Transesophageal Echocardiography. *J Emerg Med* 2017; **53**: 722-725 [PMID: 29128035 DOI: 10.1016/j.jemermed.2017.08.011]
 - 21 **Oikawa T**, Shimazui T, Johraku A, Kihara S, Tsukamoto S, Miyanaga N, Hattori K, Kawai K, Uchida K, Takeshima H, Saito S, Toyooka H, Akaza H. Intraoperative transesophageal echocardiography for inferior vena caval tumor thrombus in renal cell carcinoma. *Int J Urol* 2004; **11**: 189-92 [PMID: 15028094 DOI: 10.1111/j.1442-2042.2003.00780.x]
 - 22 **Konstantinides SV**, Meyer G, Becattini C, Bueno H, Geersing GJ, Harjola VP, Huisman MV, Humbert M, Jennings CS, Jiménez D, Kucher N, Lang IM, Lankeit M, Lorusso R, Mazzolai L, Meneveau N, Ni Ainle F, Prandoni P, Pruszczyk P, Righini M, Torbicki A, Van Belle E, Zamorano JL; ESC Scientific Document Group. 2019 ESC Guidelines for the diagnosis and management of acute pulmonary embolism developed in collaboration with the European Respiratory Society (ERS). *Eur Heart J* 2020; **41**: 543-603 [PMID: 31504429 DOI: 10.1093/eurheartj/ehz405]
 - 23 **Duffett L**, Castellucci LA, Forgie MA. Pulmonary embolism: update on management and controversies. *BMJ* 2020; **370**: m2177 [PMID: 32759284 DOI: 10.1136/bmj.m2177]
 - 24 **Desciak MC**, Martin DE. Perioperative pulmonary embolism: diagnosis and anesthetic management. *J Clin Anesth* 2011; **23**: 153-65 [PMID: 21377083 DOI: 10.1016/j.jclinane.2010.06.011]
 - 25 **Oh YN**, Oh DK, Koh Y, Lim CM, Huh JW, Lee JS, Jung SH, Kang PJ, Hong SB. Use of extracorporeal membrane oxygenation in patients with acute high-risk pulmonary embolism: a case series with literature review. *Acute Crit Care* 2019; **34**: 148-154 [PMID: 31723920 DOI: 10.4266/acc.2019.00500]
 - 26 **Stevens SM**, Woller SC, Kreuziger LB, Bounameaux H, Doerschug K, Geersing GJ, Huisman MV, Kearon C, King CS, Knighton AJ, Lake E, Murin S, Vintch JRE, Wells PS, Moores LK. Antithrombotic Therapy for VTE Disease: Second Update of the CHEST Guideline and Expert Panel Report. *Chest* 2021; **160**: e545-e608 [PMID: 34352278 DOI: 10.1016/j.chest.2021.07.055]
 - 27 **Stevens SM**, Woller SC, Baumann Kreuziger L, Bounameaux H, Doerschug K, Geersing GJ, Huisman MV, Kearon C, King CS, Knighton AJ, Lake E, Murin S, Vintch JRE, Wells PS, Moores LK. Executive Summary: Antithrombotic Therapy for VTE Disease: Second Update of the CHEST Guideline and Expert Panel Report. *Chest* 2021; **160**: 2247-2259 [PMID: 34352279 DOI: 10.1016/j.chest.2021.07.056]
 - 28 **Deas DS**, Keeling B. Surgical Pulmonary Embolectomy. *Crit Care Clin* 2020; **36**: 497-504 [PMID: 32473694 DOI: 10.1016/j.ccc.2020.02.009]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA
Telephone: +1-925-3991568
E-mail: bpgoffice@wjgnet.com
Help Desk: <https://www.f6publishing.com/helpdesk>
<https://www.wjgnet.com>

