

World Journal of *Clinical Cases*

World J Clin Cases 2022 July 16; 10(20): 6759-7186



Contents

Thrice Monthly Volume 10 Number 20 July 16, 2022

OPINION REVIEW

- 6759 Semaglutide might be a key for breaking the vicious cycle of metabolically associated fatty liver disease spectrum?

Cigrovski Berkovic M, Rezić T, Bilic-Curcic I, Mrzljak A

MINIREVIEWS

- 6769 Drainage of pancreatic fluid collections in acute pancreatitis: A comprehensive overview
Bansal A, Gupta P, Singh AK, Shah J, Samanta J, Mandavdhare HS, Sharma V, Sinha SK, Dutta U, Sandhu MS, Kochhar R

- 6784 Frontiers of COVID-19-related myocarditis as assessed by cardiovascular magnetic resonance

Luo Y, Liu BT, Yuan WF, Zhao CX

ORIGINAL ARTICLE

Case Control Study

- 6794 Urinary and sexual function changes in benign prostatic hyperplasia patients before and after transurethral columnar balloon dilatation of the prostate

Zhang DP, Pan ZB, Zhang HT

- 6803 Effects of the information-knowledge-attitude-practice nursing model combined with predictability intervention on patients with cerebrovascular disease

Huo HL, Gui YY, Xu CM, Zhang Y, Li Q

Retrospective Cohort Study

- 6811 Effects of Kampo medicine hangebyakujutsutemmato on persistent postural-perceptual dizziness: A retrospective pilot study

Miwa T, Kanemaru SI

Retrospective Study

- 6825 Longitudinal changes in personalized platelet count metrics are good indicators of initial 3-year outcome in colorectal cancer

Herold Z, Herold M, Lohinszky J, Szasz AM, Dank M, Somogyi A

- 6845 Efficacy of Kegel exercises in preventing incontinence after partial division of internal anal sphincter during anal fistula surgery

Garg P, Yagnik VD, Kaur B, Menon GR, Dawka S

Observational Study

- 6855 Influence of the water jet system vs cavitron ultrasonic surgical aspirator for liver resection on the remnant liver

Hanaki T, Tsuda A, Sunaguchi T, Goto K, Morimoto M, Murakami Y, Kihara K, Matsunaga T, Yamamoto M, Tokuyasu N, Sakamoto T, Hasegawa T, Fujiwara Y

- 6865** Critical values of monitoring indexes for perioperative major adverse cardiac events in elderly patients with biliary diseases

Zhang ZM, Xie XY, Zhao Y, Zhang C, Liu Z, Liu LM, Zhu MW, Wan BJ, Deng H, Tian K, Guo ZT, Zhao XZ

- 6876** Comparative study of surface electromyography of masticatory muscles in patients with different types of bruxism

Lan KW, Jiang LL, Yan Y

Randomized Controlled Trial

- 6890** Dural puncture epidural technique provides better anesthesia quality in repeat cesarean delivery than epidural technique: Randomized controlled study

Wang SY, He Y, Zhu HJ, Han B

SYSTEMATIC REVIEWS

- 6900** Network pharmacology-based strategy for predicting therapy targets of Sanqi and Huangjing in diabetes mellitus

Cui XY, Wu X, Lu D, Wang D

META-ANALYSIS

- 6915** Endoscopic submucosal dissection for early signet ring cell gastric cancer: A systematic review and meta-analysis

Weng CY, Sun SP, Cai C, Xu JL, Lv B

- 6927** Prognostic value of computed tomography derived skeletal muscle mass index in lung cancer: A meta-analysis

Pan XL, Li HJ, Li Z, Li ZL

CASE REPORT

- 6936** Autosomal dominant osteopetrosis type II resulting from a *de novo* mutation in the *CLCN7* gene: A case report

Song XL, Peng LY, Wang DW, Wang H

- 6944** Clinical expression and mitochondrial deoxyribonucleic acid study in twins with 14484 Leber's hereditary optic neuropathy: A case report

Chuenkongkaew WL, Chinkulkitnivat B, Lertrit P, Chirapapaisan N, Kaewsutthi S, Suktitipat B, Mitrpant C

- 6954** Management of the enteroatmospheric fistula: A case report

Cho J, Sung K, Lee D

- 6960** Lower lip recurrent keratoacanthoma: A case report

Liu XG, Liu XG, Wang CJ, Wang HX, Wang XX

- 6966** Optic disc coloboma associated with macular retinoschisis: A case report

Zhang W, Peng XY

- 6974** A 7-year-old boy with recurrent cyanosis and tachypnea: A case report
Li S, Chen LN, Zhong L
- 6981** Schwannomatosis patient who was followed up for fifteen years: A case report
Li K, Liu SJ, Wang HB, Yin CY, Huang YS, Guo WT
- 6991** Intentional replantation combined root resection therapy for the treatment of type III radicular groove with two roots: A case report
Tan D, Li ST, Feng H, Wang ZC, Wen C, Nie MH
- 6999** Clinical features and genetic variations of severe neonatal hyperbilirubinemia: Five case reports
Lin F, Xu JX, Wu YH, Ma YB, Yang LY
- 7006** Percutaneous transhepatic access for catheter ablation of a patient with heterotaxy syndrome complicated with atrial fibrillation: A case report
Wang HX, Li N, An J, Han XB
- 7013** Secondary positioning of rotationally asymmetric refractive multifocal intraocular lens in a patient with glaucoma: A case report
Fan C, Zhou Y, Jiang J
- 7020** Laparoscopic repair of diaphragmatic hernia associating with radiofrequency ablation for hepatocellular carcinoma: A case report
Tsunoda J, Nishi T, Ito T, Inaguma G, Matsuzaki T, Seki H, Yasui N, Sakata M, Shimada A, Matsumoto H
- 7029** Hypopituitary syndrome with pituitary crisis in a patient with traumatic shock: A case report
Zhang XC, Sun Y
- 7037** Solitary plasmacytoma of the left rib misdiagnosed as angina pectoris: A case report
Yao J, He X, Wang CY, Hao L, Tan LL, Shen CJ, Hou MX
- 7045** Secondary coronary artery ostial lesions: Three case reports
Liu XP, Wang HJ, Gao JL, Ma GL, Xu XY, Ji LN, He RX, Qi BYE, Wang LC, Li CQ, Zhang YJ, Feng YB
- 7054** Bladder perforation injury after percutaneous peritoneal dialysis catheterization: A case report
Shi CX, Li ZX, Sun HT, Sun WQ, Ji Y, Jia SJ
- 7060** Myotonic dystrophy type 1 presenting with dyspnea: A case report
Jia YX, Dong CL, Xue JW, Duan XQ, Xu MY, Su XM, Li P
- 7068** Novel mutation in the *SALL1* gene in a four-generation Chinese family with uraemia: A case report
Fang JX, Zhang JS, Wang MM, Liu L
- 7076** Malignant transformation of primary mature teratoma of colon: A case report
Liu J

- 7082** Treatment of pyogenic liver abscess by surgical incision and drainage combined with platelet-rich plasma: A case report
Wang JH, Gao ZH, Qian HL, Li JS, Ji HM, Da MX
- 7090** Left bundle branch pacing in a ventricular pacing dependent patient with heart failure: A case report
Song BX, Wang XX, An Y, Zhang YY
- 7097** Solitary fibrous tumor of the liver: A case report and review of the literature
Xie GY, Zhu HB, Jin Y, Li BZ, Yu YQ, Li JT
- 7105** MutL homolog 1 germline mutation c.(453+1_454-1)_(545+1_546-1)del identified in lynch syndrome: A case report and review of literature
Zhang XW, Jia ZH, Zhao LP, Wu YS, Cui MH, Jia Y, Xu TM
- 7116** Malignant histiocytosis associated with mediastinal germ cell tumor: A case report
Yang PY, Ma XL, Zhao W, Fu LB, Zhang R, Zeng Q, Qin H, Yu T, Su Y
- 7124** Immunoglobulin G4 associated autoimmune cholangitis and pancreatitis following the administration of nivolumab: A case report
Agrawal R, Guzman G, Karimi S, Giulianotti PC, Lora AJM, Jain S, Khan M, Boulay BR, Chen Y
- 7130** Portal vein thrombosis in a noncirrhotic patient after hemihepatectomy: A case report and review of literature
Zhang SB, Hu ZX, Xing ZQ, Li A, Zhou XB, Liu JH
- 7138** Microvascular decompression for a patient with oculomotor palsy caused by posterior cerebral artery compression: A case report and literature review
Zhang J, Wei ZJ, Wang H, Yu YB, Sun HT
- 7147** Topical halometasone cream combined with fire needle pre-treatment for treatment of primary cutaneous amyloidosis: Two case reports
Su YQ, Liu ZY, Wei G, Zhang CM
- 7153** Simultaneous robot-assisted approach in a super-elderly patient with urothelial carcinoma and synchronous contralateral renal cell carcinoma: A case report
Yun JK, Kim SH, Kim WB, Kim HK, Lee SW
- 7163** Nursing a patient with latent autoimmune diabetes in adults with insulin-related lipodystrophy, allergy, and exogenous insulin autoimmune syndrome: A case report
He F, Xu LL, Li YX, Dong YX
- 7171** Incidental diagnosis of medullary thyroid carcinoma due to persistently elevated procalcitonin in a patient with COVID-19 pneumonia: A case report
Saha A, Mukhopadhyay M, Paul S, Bera A, Bandyopadhyay T
- 7178** Macular hole following phakic intraocular lens implantation: A case report
Li XJ, Duan JL, Ma JX, Shang QL

LETTER TO THE EDITOR

- 7184** Is every microorganism detected in the intensive care unit a nosocomial infection? Isn't prevention more important than detection?

Yildirim F, Karaman I, Yildirim M

ABOUT COVER

Editorial Board Member of *World Journal of Clinical Cases*, Jie-Feng Huang, PhD, Associate Chief Physician, Associate Professor, Department of Orthopaedics and Traumatology, The First Affiliated Hospital of Zhejiang Chinese Medical University, Hangzhou 310006, Zhejiang Province, China. 40983285@qq.com

AIMS AND SCOPE

The primary aim of *World Journal of Clinical Cases* (WJCC, *World J Clin Cases*) is to provide scholars and readers from various fields of clinical medicine with a platform to publish high-quality clinical research articles and communicate their research findings online.

WJCC mainly publishes articles reporting research results and findings obtained in the field of clinical medicine and covering a wide range of topics, including case control studies, retrospective cohort studies, retrospective studies, clinical trials studies, observational studies, prospective studies, randomized controlled trials, randomized clinical trials, systematic reviews, meta-analysis, and case reports.

INDEXING/ABSTRACTING

The WJCC is now abstracted and indexed in Science Citation Index Expanded (SCIE, also known as SciSearch®), Journal Citation Reports/Science Edition, Current Contents®/Clinical Medicine, PubMed, PubMed Central, Scopus, Reference Citation Analysis, China National Knowledge Infrastructure, China Science and Technology Journal Database, and Superstar Journals Database. The 2022 Edition of Journal Citation Reports® cites the 2021 impact factor (IF) for WJCC as 1.534; IF without journal self cites: 1.491; 5-year IF: 1.599; Journal Citation Indicator: 0.28; Ranking: 135 among 172 journals in medicine, general and internal; and Quartile category: Q4. The WJCC's CiteScore for 2021 is 1.2 and Scopus CiteScore rank 2021: General Medicine is 443/826.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: Hua-Ge Yin; Production Department Director: Xu Guo; Editorial Office Director: Jin-Lei Wang.

NAME OF JOURNAL

World Journal of Clinical Cases

ISSN

ISSN 2307-8960 (online)

LAUNCH DATE

April 16, 2013

FREQUENCY

Thrice Monthly

EDITORS-IN-CHIEF

Bao-Gan Peng, Jerzy Tadeusz Chudek, George Kontogeorgos, Maurizio Serati, Ja Hyeon Ku

EDITORIAL BOARD MEMBERS

<https://www.wjgnet.com/2307-8960/editorialboard.htm>

PUBLICATION DATE

July 16, 2022

COPYRIGHT

© 2022 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjgnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>



Case Control Study

Effects of the information–knowledge–attitude–practice nursing model combined with predictability intervention on patients with cerebrovascular disease

Hong-Liang Huo, Yuan-Yuan Gui, Chun-Miao Xu, Yan Zhang, Qiang Li

Specialty type: Nursing

Provenance and peer review:

Unsolicited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's scientific quality classification

Grade A (Excellent): 0
Grade B (Very good): B
Grade C (Good): C
Grade D (Fair): 0
Grade E (Poor): 0

P-Reviewer: Gillies RJ, United States; Gronberg A, Sweden

A-Editor: Zhu JQ, China

Received: March 8, 2022

Peer-review started: March 8, 2022

First decision: April 13, 2022

Revised: April 27, 2022

Accepted: May 28, 2022

Article in press: May 28, 2022

Published online: July 16, 2022



Hong-Liang Huo, Department of Nursing, The Fourth Affiliated Hospital of Qiqihar Medical University, Qiqihar 161000, Heilongjiang Province, China

Yuan-Yuan Gui, Chun-Miao Xu, Yan Zhang, Qiang Li, Nursing School, Qiqihar Medical University, Qiqihar 161006, Heilongjiang Province, China

Corresponding author: Chun-Miao Xu, MM, Nurse, Nursing School, Qiqihar Medical University, No. 333 Bukui North Street, Jianhua District, Qiqihar 161006, Heilongjiang Province, China. xcmiao02@163.com

Abstract

BACKGROUND

Cerebrovascular disease (CVD) poses a serious threat to human health and safety. Thus, developing a reasonable exercise program plays an important role in the long-term recovery and prognosis for patients with CVD. Studies have shown that predictive nursing can improve the quality of care and that the information–knowledge–attitude–practice (IKAP) nursing model has a positive impact on patients who suffered a stroke. Few studies have combined these two nursing models to treat CVD.

AIM

To explore the effect of the IKAP nursing model combined with predictive nursing on the Fugl–Meyer motor function (FMA) score, Barthel index score, and disease knowledge mastery rate in patients with CVD.

METHODS

A total of 140 patients with CVD treated at our hospital between December 2019 and September 2021 were randomly divided into two groups, with 70 patients in each. The control group received routine nursing, while the observation group received the IKAP nursing model combined with predictive nursing. Both groups were observed for self-care ability, motor function, and disease knowledge mastery rate after one month of nursing.

RESULTS

There was no clear difference between the Barthel index and FMA scores of the two groups before nursing ($P > 0.05$); however, their scores increased after

nursing. This increase was more apparent in the observation group, and the difference was statistically significant ($P < 0.05$). The rates of disease knowledge mastery, timely medication, appropriate exercise, and reasonable diet were significantly higher in the observation group than in the control group ($P < 0.05$). The satisfaction rate in the observation group (97.14%) was significantly higher than that in the control group (81.43%; $P < 0.05$).

CONCLUSION

The IKAP nursing model, combined with predictive nursing, is more effective than routine nursing in the care of patients with CVD, and it can significantly improve the Barthel index and FMA scores with better knowledge acquisition, as well as produce high satisfaction in patients. Moreover, they can be widely used in the clinical setting.

Key Words: Information–knowledge–attitude–practice nursing model; Predictive nursing; Cerebrovascular disease; Barthel index; Fugl–Meyer motor function score; Disease knowledge mastery rate

©The Author(s) 2022. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: The information–knowledge–attitude–practice nursing model combined with predictable intervention enhances the nursing effect of cerebrovascular disease cases, significantly improves the Barthel index and Fugl–Meyer motor function scores, and leaves patients highly satisfied and can be widely used in clinical practice.

Citation: Huo HL, Gui YY, Xu CM, Zhang Y, Li Q. Effects of the information–knowledge–attitude–practice nursing model combined with predictability intervention on patients with cerebrovascular disease. *World J Clin Cases* 2022; 10(20): 6803-6810

URL: <https://www.wjgnet.com/2307-8960/full/v10/i20/6803.htm>

DOI: <https://dx.doi.org/10.12998/wjcc.v10.i20.6803>

INTRODUCTION

Cerebrovascular disease (CVD) is common in clinical settings; its characteristics include acute onset, severe condition, and poor prognosis, which pose a serious threat to human health and safety[1]. In addition to symptomatic treatment, patients with CVD require functional exercises for long-term rehabilitation, which play an important role in long-term recovery and prognosis[2,3]. However, it is difficult for patients to adhere to the process of rehabilitation training, so nursing staff should formulate reasonable plans to improve patients' enthusiasm and compliance with exercise to improve their prognosis[4,5]. Predictive nursing refers to forecasting outcomes before patients have obvious symptoms, anticipating symptoms, and implementing targeted and effective nursing measures. Research has shown that predictive nursing can improve the quality of nursing[6-8]. Recently, the information–knowledge–attitude–practice (IKAP) nursing model has shown a positive effect on stroke patients, which entails the formulation of individualized, humanized, and professional nursing measures according to a patient's unique circumstances[9]. Few studies have investigated the combination of these two nursing modes for CVD. This study explored the effect of the IKAP nursing mode combined with predictive nursing on the Fugl–Meyer motor function (FMA) score, Barthel index score, and disease knowledge mastery rate in patients with CVD, and it described an intervention for clinical nursing.

MATERIALS AND METHODS

General information

A total of 140 patients with CVD treated at our hospital between December 2019 and September 2021 were randomly divided into two groups of 70 individuals each. The observation group comprised 39 men and 31 women. They ranged in age from 43 to 77 years, with an average age of 53.4 ± 4.7 years. Disease types included cerebral hemorrhage (21 cases), cerebral infarction (30 cases), subarachnoid hemorrhage (10 cases), and transient ischemic attack (9 cases). The control group included 33 women and 37 men. They ranged in age from 42 to 75 years, with an average age of 53.9 ± 4.3 years. The disease types included cerebral hemorrhage ($n = 23$), cerebral infarction ($n = 28$), subarachnoid hemorrhage ($n = 14$), and transient ischemic attack ($n = 5$). There was no clear difference between the two groups ($P >$

0.05), indicating comparability.

Inclusion and exclusion criteria

The inclusion criteria were as follows: meeting the CVD diagnostic criteria; the patient was clearly diagnosed with a head computed tomography scan or magnetic resonance imaging; and all patients were included in the first or experienced onset.

The exclusion criteria were as follows: severe liver, heart, kidney, or other organ dysfunction; previous CVD; traumatic brain dysfunction; other brain diseases; and other neurological and mental disorders.

Methods

The control group received routine nursing care, including monitoring vital signs, oxygen inhalation, and other care activities. The observation group received IKAP nursing combined with predictive nursing.

The IKAP nursing model is implemented as follows. To understand the patient's information, design a CVD-related knowledge questionnaire to communicate with patients. Offer individualized psychological guidance to understand the patient's psychological status; provide the corresponding psychological care; alleviate the patient's anxiety, depression, and other negative emotions; provide health education; develop a health education manual; and assist patients in establishing a reasonable diet. Health education is provided to patients and their families through seminars or exchanges, including guidance for those with speech disorders, whereby the type and extent of speech disorders can be assessed. Furthermore, patients are guided through rehabilitation, plural, and stimulation training according to their specific circumstances to maintain maximum communication skills. Patients receive limb function training and early guidance for functional exercises, thereby achieving a better prognosis. Corresponding training is carried out according to the patients' unique circumstances, from passive to active, to help them with autonomous eating, dressing, as well as removing their shoes. Equipment use training was simultaneously provided. Patients are assisted in changing their bad habits, promoting healthy behavior, and improving their self-care abilities.

Predictive nursing was conducted as follows. In terms of dietary nursing, patients with CVD may have swallowing disorders and be prone to choking or coughing due to degeneration of nerve reflex activity. Therefore, diverse paste-like foods should be provided at an early stage. If the patient is bedridden for a long time, intestinal peristalsis can slow down, and digestive function can become reduced; therefore, it is necessary for the patient to chew slowly, eat smaller portions, and consume more meals. To prevent pulmonary infection, one should recognize that patients with longer hospital stays are at an increased risk of pulmonary infection and therefore require intensive nursing. Patients should be encouraged to cough and excrete sputum independently. For those who cannot easily cough sputum, back patting and nebulized inhalation are feasible for excretion. For comatose patients, the respiratory tract should remain unobstructed, and the head should be placed on one side. To actively treat the underlying disease of the lungs, stress ulcers should be examined. Gastric and duodenal ulcers can emerge in patients with CVD. The more severe the condition, the higher the incidence. Therefore, timely measures should be implemented.

First, we determined whether each patient had a history of ulcers or erratic and abdominal distension to rule out stress factors. We then monitored changes in vital signs and observed the patient's state of consciousness. We noted the condition of the stools, checked occult blood regularly, and observed the nature of vomitus.

To prevent a decubitus ulcer, the color of the skin in areas prone to pressure sores should be checked. For pressure sores, the site, size, and depth should be assessed, and a nursing plan should be developed accordingly. The clothes should be changed regularly, the skin should be kept clean and dry, and soft pads should be used to reduce the pressure on the contact surface. In addition, patients should be turned over regularly (their position should be adjusted once every two hours), and the areas of pressure sores should be massaged often. The patients in both groups were nursed for one month.

The Barthel index scores of patients in both groups

The Barthel index was used to evaluate patients' ability to live before and after nursing.

Comparison of motor functions

Motor function was evaluated before and after nursing using the FMA score. The highest score was 100, with lower scores indicating more pronounced motor impairment.

Comparison of compliance behavior between the two groups

The disease knowledge mastery rate, timely medication administration, reasonable diet, and appropriate exercise were compared between the two groups after nursing.

Satisfaction rate

Satisfaction questionnaires were designed to observe satisfaction among patients in both groups; the satisfaction rates were classified as “very satisfied,” “satisfied,” “average,” or “unsatisfactory.”

Statistical analysis

Statistical data were analyzed using the SPSS software version 19.0. The count data were represented as a percentage. A chi-square test was used to compare the groups, and measurement data were represented by mean \pm SD. Independent samples and paired *t*-tests were used, with $P < 0.05$, indicating that the difference was statistically significant.

RESULTS**Comparison of the living ability of the two groups of patients**

The difference between the Barthel index scores of the two groups before nursing was not statistically significant ($P > 0.05$), but that of both groups increased after nursing. The increase was greater in the observation group, and the difference was statistically significant ($P < 0.05$; Table 1).

Comparison of motor function between the two groups of patients

The difference between the FMA scores of the patients in the two groups before nursing was not statistically significant ($P > 0.05$), but increased for both groups after nursing. The increase was greater in the observation group, and the difference was statistically significant ($P < 0.05$; Table 2).

Comparison of compliance behavior between the two groups

The disease knowledge mastery rate, timely medication, appropriate exercise, and reasonable diet were significantly higher in the observation group than in the control group ($P < 0.05$; Table 3).

Comparison of satisfaction between the two groups

The satisfaction rate was 97.14% in the observation group and 81.43% in the control group, which was significantly higher than that in the control group ($P < 0.05$; Table 4).

DISCUSSION

CVD is a common illness that manifests mainly as cerebral hemorrhage and infarction, and stroke is the leading cause of adult deaths[10,11]. CVD has a high morbidity and mortality rate; it often starts suddenly and is serious, leaving patients with sequelae such as speech impairment and hemiplegia, which bring great inconvenience to their lives. Appropriate nursing can effectively improve prognosis and quality of life[12,13]. This study explored the effect of IKAP nursing model combined with predictive nursing on patients with CVD. The IKAP nursing model is a systematic, comprehensive nursing model that requires individualized education with a focus on communication and interactions among nursing staff, families, patients, and society. The model entails adopting appropriate methods to communicate with patients in a targeted manner to improve their knowledge related to CVD, boost their self-management ability, change their mindset, and motivate them to exercise[14,15]. The IKAP nursing model lays the foundation for nursing by collecting information and designing a knowledge questionnaire on CVD so that patients can understand the disease. The results of the full survey and the cognitive ability of the patients were used to develop an educational program on the prevention and treatment of CVD so that the patients could understand the importance of rehabilitation. An individualized psychological nursing plan should be developed, psychological health counseling should be provided, and patients' belief in recovery should be reinforced. Rehabilitation training, including speech and muscle function training, is performed according to the patient's specific situation to facilitate behavioral changes[16,17]. Predictive nursing involves a comprehensive, integrated analysis of a patient during the nursing procedure to anticipate risks in advance so that effective measures can be taken early to avoid or reduce complications. The IKAP model centers on the prevention and control of possible dangers or problems in all aspects of treatment based on the experience of previous work and interventions through individualized assessments to make nursing safer, more effective, and more cooperative with patients. Simultaneously, predictive nursing can help nurses cultivate their thinking and judgment abilities, improve work motivation, and increase work efficiency[18-20]. In this study, there was no statistically significant difference between the FMA and Barthel index scores of the two groups before nursing ($P > 0.05$). The Barthel index and FMA scores in both groups increased after nursing, with the increase being more significant in the observation group. The differences were statistically significant ($P < 0.05$). The rates of disease knowledge mastery, timely medication, appropriate exercise, and reasonable diet were significantly higher in the observation group than in the control

Table 1 Comparison of living ability of the two groups of patients (mean \pm SD)

Group	Number of cases	Barthel index			
		Before nursing	After nursing	<i>t</i> value	<i>P</i> value
Observation group	70	28.44 \pm 5.01	68.94 \pm 10.52	32.292	0.000
Control group	70	28.26 \pm 4.68	45.63 \pm 9.63	13.472	0.000
<i>t</i> value		0.227	13.673		
<i>P</i> value		0.821	0.000		

Table 2 Comparison of motor function between the two groups (mean \pm SD)

Group	Number of cases	FMA scores			
		Before nursing	After nursing	<i>t</i> value	<i>P</i> value
Observation group	70	27.33 \pm 4.52	56.73 \pm 7.47	30.814	0.000
Control group	70	27.29 \pm 4.85	40.39 \pm 6.84	12.849	0.000
<i>t</i> value		0.054	13.504		
<i>P</i> value		0.957	0.000		

FMA: Fugl-Meyer motor function score.

Table 3 Comparisons of compliance behavior between the two groups, *n* (%)

Group	Number of cases	Disease knowledge acquisition	Timely medication	Appropriate exercise	Reasonable diet
Observation group	70	66 (94.29)	70 (100.00)	64 (91.43)	67 (95.71)
Control group	70	52 (74.29)	56 (80.00)	48 (68.57)	50 (71.43)
χ^2		10.570	15.556	11.429	15.035
<i>P</i> value		0.001	0.000	0.001	0.000

Table 4 Comparison of satisfaction between the two groups of invalids, *n* (%)

Group	Very satisfied	Satisfied	Average	Unsatisfied	Satisfaction rate
Observation group	45 (64.29)	18 (25.71)	5 (7.14)	2 (2.86)	68 (97.14)
Control group	24 (34.29)	23 (32.86)	10 (14.29)	13 (18.57)	57 (81.43)
χ^2					9.035
<i>P</i> value					0.003

group ($P < 0.05$). The satisfaction rate in the observation group (97.14%) was higher than that in the control group (81.43%), and the difference was statistically significant ($P < 0.05$), indicating that the IKAP nursing model combined with predictive nursing was more effective in the care of patients with CVD. It allows nursing staff to fully understand a patient's specific situation, to predict problems in advance, to think of appropriate nursing measures, and to provide timely and appropriate care for problems that have already occurred, thereby significantly improving a patient's ability to live, along with improving their motor function, disease knowledge mastery, timely medication, and satisfaction levels.

CONCLUSION

The IKAP nursing model, combined with predictive nursing, is more effective than routine nursing in the care of patients with CVD; it can significantly improve the Barthel index and FMA scores with better

knowledge acquisition and produce high satisfaction in patients. Moreover, it can be widely used in the clinical setting.

ARTICLE HIGHLIGHTS

Research background

Studies have shown that predictive nursing can improve the quality of care and that the information-knowledge-attitude-practice (IKAP) nursing model has a positive impact on stroke patients. Establishing a reasonable exercise plan plays an important role in long-term rehabilitation and prognosis for patients with cerebrovascular disease (CVD).

Research motivation

There are no reports of predictive nursing combined with the IKAP nursing model for CVD.

Research objectives

This study aimed to explore the effects of IKAP nursing model combined with predictive nursing on the Fugl-Meyer motor function (FMA) score, Barthel index score, and disease knowledge mastery rate in patients with CVD.

Research methods

A total of 140 patients with CVD were randomly divided into control and observation groups. The control group received routine nursing, while the observation group received IKAP combined with predictive nursing. Both groups were observed for self-care ability, motor function, and disease knowledge mastery rate after one month of nursing.

Research results

There was no clear difference between the Barthel index and FMA scores of the two groups before nursing, but their scores increased after nursing. This increase was more apparent in the observation group, and the difference was statistically significant. The rates of disease knowledge mastery, timely medication, appropriate exercise, and reasonable diet were significantly higher in the observation group than in the control group. The satisfaction rate in the observation group was significantly higher than that in the control group.

Research conclusions

The IKAP nursing model, combined with predictive nursing, is more effective than routine nursing in the care of patients with CVD; it can significantly improve the Barthel index and FMA scores with better knowledge acquisition and produce high satisfaction in patients. Moreover, it can be widely used in the clinical setting.

Research perspectives

The IKAP model of care, combined with predictive care, can be widely used in clinical settings to significantly improve long-term care and outcomes in patients with CVD.

FOOTNOTES

Author contributions: Huo HL designed and performed the research, collected and analyzed the data, and wrote the article; Gui YY and Zhang Y collected the data and samples; Xu CM analyzed the data; and Xu CM and Li Q edited, reviewed, and approved the final article.

Supported by Basic scientific research industry of Heilongjiang Provincial undergraduate universities in 2019, No. 2019-KYYWF-1213.

Institutional review board statement: The study protocol conformed to the ethical guidelines of the Declaration of Helsinki and was approved by the Ethics Committee of Qiqihar Medical University.

Informed consent statement: All study participants or their legal guardians provided written informed consent prior to study enrollment.

Conflict-of-interest statement: The authors declare no conflicts of interest.

Data sharing statement: No additional data are available.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Country/Territory of origin: China

ORCID number: Hong-Liang Huo 0000-0002-4652-7487; Yuan-Yuan Gui 0000-0001-6763-7725; Chun-Miao Xu 0000-0002-2112-0146; Yan Zhang 0000-0002-3024-7923; Qiang Li 0000-0002-1864-2437.

S-Editor: Wang JL

L-Editor: A

P-Editor: Wang JL

REFERENCES

- 1 Liu L, Chen W, Zhou H, Duan W, Li S, Huo X, Xu W, Huang L, Zheng H, Liu J, Liu H, Wei Y, Xu J, Wang Y; Chinese Stroke Association Stroke Council Guideline Writing Committee. Chinese Stroke Association guidelines for clinical management of cerebrovascular disorders: executive summary and 2019 update of clinical management of ischaemic cerebrovascular diseases. *Stroke Vasc Neurol* 2020; **5**: 159-176 [PMID: 32561535 DOI: 10.1136/svn-2020-000378]
- 2 Kim Y, Lai B, Mehta T, Thirumalai M, Padalabalanarayanan S, Rimmer JH, Motl RW. Exercise Training Guidelines for Multiple Sclerosis, Stroke, and Parkinson Disease: Rapid Review and Synthesis. *Am J Phys Med Rehabil* 2019; **98**: 613-621 [PMID: 30844920 DOI: 10.1097/PHM.0000000000001174]
- 3 Rose DK, Nadeau SE, Wu SS, Tilson JK, Dobkin BH, Pei Q, Duncan PW. Locomotor Training and Strength and Balance Exercises for Walking Recovery After Stroke: Response to Number of Training Sessions. *Phys Ther* 2017; **97**: 1066-1074 [PMID: 29077960 DOI: 10.1093/ptj/pzx079]
- 4 Cai WX, Su D, Zhang JL, Dian HJ. [Evaluation of clinical teaching model for nursing practice in intensive care unit for cerebrovascular disease: An individualized instruction model]. *Zhongguo Cuzhong Zazhi* 2015; **5**: 619-621 [DOI: 10.3969/j.issn.1673-5765.2015.07.015]
- 5 Gouya G, Arrich J, Wolzt M, Huber K, Verheugt FW, Gurbel PA, Pirker-Kees A, Siller-Matula JM. Antiplatelet treatment for prevention of cerebrovascular events in patients with vascular diseases: a systematic review and meta-analysis. *Stroke* 2014; **45**: 492-503 [PMID: 24368560 DOI: 10.1161/STROKEAHA.113.002590]
- 6 Li XX, Du XW, Song W, Lu C, Hao WN. Effect of continuous nursing care based on the IKAP theory on the quality of life of patients with chronic obstructive pulmonary disease: A randomized controlled study. *Medicine (Baltimore)* 2020; **99**: e19543 [PMID: 32176107 DOI: 10.1097/MD.00000000000019543]
- 7 Guo X, Men F, Han X, Wang Z. The efficacy of continuous nursing care for patients with chronic obstructive pulmonary disease: A randomized controlled trial protocol. *Medicine (Baltimore)* 2021; **100**: e23974 [PMID: 33466137 DOI: 10.1097/MD.00000000000023974]
- 8 Zhao DD, Bi J. [Observation on application effect of predictive nursing for patients undergoing ICU operation in department of neurosurgery]. *Huli Yanjiu* 2015; **5**: 4286-4288 [DOI: 10.3969/j.issn.1009-6493.2015.34.020]
- 9 Du AP. [Impact of IKAP care model on quality of life and treatment adherence of stroke patients]. *Guizhou Yiyao* 2017; **41**: 556-557 [DOI: 10.3969/j.issn.1000-744X.2017.05.051]
- 10 Qu JF, Chen YK, Zhong HH, Li W, Lu ZH. Preexisting Cerebral Abnormalities and Functional Outcomes After Acute Ischemic Stroke. *J Geriatr Psychiatry Neurol* 2019; **32**: 327-335 [PMID: 31480988 DOI: 10.1177/0891988719862631]
- 11 Kim J, Thayabaranathan T, Donnan GA, Howard G, Howard VJ, Rothwell PM, Feigin V, Norrving B, Owolabi M, Pandian J, Liu L, Cadilhac DA, Thrift AG. Global Stroke Statistics 2019. *Int J Stroke* 2020; **15**: 819-838 [PMID: 32146867 DOI: 10.1177/1747493020909545]
- 12 Zhou HS, Chen WL, Li MY. Analysis on related risk factors of nosocomial infections in patients with acute cerebrovascular diseases. *Zhonghua Yiyuan Ganranxue Zazhi* 2015; **5**: 1576-1578
- 13 Jiang L, Shi R, Li DD, Wang AH. Research progress of salidroside and its analogs in the treatment of ischemic cerebrovascular diseases. *Shandong Yiyao* 2016; **56**: 106-109 [DOI: 10.3969/j.issn.1002-266X.2016.36.038]
- 14 Xie XH. Application of continuous nursing based on IKAP theory in patients with cerebrovascular intervention. *Zhonghua Xiandai Huli Zazhi* 2020; **26**: 631-635 [DOI: 10.3760/cma.j.issn.1674-2907.2020.05.015]
- 15 Zeng MM, Zhang MJ. Effect of IKAP nursing mode combined with low frequency repetitive transcranial electrical stimulation on the curative effect of convalescent patients with stroke. *Guizhou Yiyao* 2020; **44**: 488-489 [DOI: 10.3969/j.issn.1000-744X.2020.03.060]
- 16 He SL. Effect of IKAP care model combined with low-frequency repetitive transcranial electrical stimulation on the efficacy of patients recovering from stroke. *Heilongjiang Yiyao Kexue* 2018; **4**: 32-24 [DOI: 10.3969/j.issn.1008-0104.2018.01.047]
- 17 Wang QL, Liu WQ, Jiao F. Health education for patients with hypertensive cerebral hemorrhage using IKAP theory. *Zhongguo Yiyao Daobao* 2010; **9**: 132-133 [DOI: 10.3969/j.issn.1673-7210.2010.09.085]
- 18 Pi Y, Tang LJ, Zhou SJ, Liu L. Predictive nursing care in treatment of the severe neurosurgical patients. *Xibu Yixue* 2014; **26**: 1728-1729; 1732 [DOI: 10.3969/j.issn.1672-3511.2014.12.055]
- 19 Dai JP, Chen YJ, Wang LX. Evaluation of the effect of anticipatory care in preventing constipation after minimally

- invasive surgery in patients with hypertensive cerebral hemorrhage. *Jianyan Yixue Yu Linchuang* 2016; **13**: 2370-2372 [DOI: [10.3969/j.issn.1672-9455.2016.16.055](https://doi.org/10.3969/j.issn.1672-9455.2016.16.055)]
- 20 **Xu DX.** Application of anticipatory care in patients with cerebrovascular disease interventions. *Quanke Huli* 2018; **5**: 578-580 [DOI: [10.3969/j.issn.1674-4748.2018.05.027](https://doi.org/10.3969/j.issn.1674-4748.2018.05.027)]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

Telephone: +1-925-3991568

E-mail: bpgoffice@wjgnet.com

Help Desk: <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

