

World Journal of *Clinical Cases*

World J Clin Cases 2022 July 16; 10(20): 6759-7186



OPINION REVIEW

- 6759 Semaglutide might be a key for breaking the vicious cycle of metabolically associated fatty liver disease spectrum?
Cigrovski Berkovic M, Rezic T, Bilic-Curcic I, Mrzljak A

MINIREVIEWS

- 6769 Drainage of pancreatic fluid collections in acute pancreatitis: A comprehensive overview
Bansal A, Gupta P, Singh AK, Shah J, Samanta J, Mandavdhare HS, Sharma V, Sinha SK, Dutta U, Sandhu MS, Kochhar R
- 6784 Frontiers of COVID-19-related myocarditis as assessed by cardiovascular magnetic resonance
Luo Y, Liu BT, Yuan WF, Zhao CX

ORIGINAL ARTICLE**Case Control Study**

- 6794 Urinary and sexual function changes in benign prostatic hyperplasia patients before and after transurethral columnar balloon dilatation of the prostate
Zhang DP, Pan ZB, Zhang HT
- 6803 Effects of the information-knowledge-attitude-practice nursing model combined with predictability intervention on patients with cerebrovascular disease
Huo HL, Gui YY, Xu CM, Zhang Y, Li Q

Retrospective Cohort Study

- 6811 Effects of Kampo medicine hangebyakujutsutemmato on persistent postural-perceptual dizziness: A retrospective pilot study
Miwa T, Kanemaru SI

Retrospective Study

- 6825 Longitudinal changes in personalized platelet count metrics are good indicators of initial 3-year outcome in colorectal cancer
Herold Z, Herold M, Lohinszky J, Szasz AM, Dank M, Somogyi A
- 6845 Efficacy of Kegel exercises in preventing incontinence after partial division of internal anal sphincter during anal fistula surgery
Garg P, Yagnik VD, Kaur B, Menon GR, Dawka S

Observational Study

- 6855 Influence of the water jet system vs cavitron ultrasonic surgical aspirator for liver resection on the remnant liver
Hanaki T, Tsuda A, Sunaguchi T, Goto K, Morimoto M, Murakami Y, Kihara K, Matsunaga T, Yamamoto M, Tokuyasu N, Sakamoto T, Hasegawa T, Fujiwara Y

- 6865** Critical values of monitoring indexes for perioperative major adverse cardiac events in elderly patients with biliary diseases

Zhang ZM, Xie XY, Zhao Y, Zhang C, Liu Z, Liu LM, Zhu MW, Wan BJ, Deng H, Tian K, Guo ZT, Zhao XZ

- 6876** Comparative study of surface electromyography of masticatory muscles in patients with different types of bruxism

Lan KW, Jiang LL, Yan Y

Randomized Controlled Trial

- 6890** Dural puncture epidural technique provides better anesthesia quality in repeat cesarean delivery than epidural technique: Randomized controlled study

Wang SY, He Y, Zhu HJ, Han B

SYSTEMATIC REVIEWS

- 6900** Network pharmacology-based strategy for predicting therapy targets of Sanqi and Huangjing in diabetes mellitus

Cui XY, Wu X, Lu D, Wang D

META-ANALYSIS

- 6915** Endoscopic submucosal dissection for early signet ring cell gastric cancer: A systematic review and meta-analysis

Weng CY, Sun SP, Cai C, Xu JL, Lv B

- 6927** Prognostic value of computed tomography derived skeletal muscle mass index in lung cancer: A meta-analysis

Pan XL, Li HJ, Li Z, Li ZL

CASE REPORT

- 6936** Autosomal dominant osteopetrosis type II resulting from a *de novo* mutation in the *CLCN7* gene: A case report

Song XL, Peng LY, Wang DW, Wang H

- 6944** Clinical expression and mitochondrial deoxyribonucleic acid study in twins with 14484 Leber's hereditary optic neuropathy: A case report

Chuenkongkaew WL, Chinkulkitnivat B, Lertrit P, Chirapapaisan N, Kaewsutthi S, Suktitipat B, Mitrpant C

- 6954** Management of the enteroatmospheric fistula: A case report

Cho J, Sung K, Lee D

- 6960** Lower lip recurrent keratoacanthoma: A case report

Liu XG, Liu XG, Wang CJ, Wang HX, Wang XX

- 6966** Optic disc coloboma associated with macular retinoschisis: A case report

Zhang W, Peng XY

- 6974** A 7-year-old boy with recurrent cyanosis and tachypnea: A case report
Li S, Chen LN, Zhong L
- 6981** Schwannomatosis patient who was followed up for fifteen years: A case report
Li K, Liu SJ, Wang HB, Yin CY, Huang YS, Guo WT
- 6991** Intentional replantation combined root resection therapy for the treatment of type III radicular groove with two roots: A case report
Tan D, Li ST, Feng H, Wang ZC, Wen C, Nie MH
- 6999** Clinical features and genetic variations of severe neonatal hyperbilirubinemia: Five case reports
Lin F, Xu JX, Wu YH, Ma YB, Yang LY
- 7006** Percutaneous transhepatic access for catheter ablation of a patient with heterotaxy syndrome complicated with atrial fibrillation: A case report
Wang HX, Li N, An J, Han XB
- 7013** Secondary positioning of rotationally asymmetric refractive multifocal intraocular lens in a patient with glaucoma: A case report
Fan C, Zhou Y, Jiang J
- 7020** Laparoscopic repair of diaphragmatic hernia associating with radiofrequency ablation for hepatocellular carcinoma: A case report
Tsunoda J, Nishi T, Ito T, Inaguma G, Matsuzaki T, Seki H, Yasui N, Sakata M, Shimada A, Matsumoto H
- 7029** Hypopituitary syndrome with pituitary crisis in a patient with traumatic shock: A case report
Zhang XC, Sun Y
- 7037** Solitary plasmacytoma of the left rib misdiagnosed as angina pectoris: A case report
Yao J, He X, Wang CY, Hao L, Tan LL, Shen CJ, Hou MX
- 7045** Secondary coronary artery ostial lesions: Three case reports
Liu XP, Wang HJ, Gao JL, Ma GL, Xu XY, Ji LN, He RX, Qi BYE, Wang LC, Li CQ, Zhang YJ, Feng YB
- 7054** Bladder perforation injury after percutaneous peritoneal dialysis catheterization: A case report
Shi CX, Li ZX, Sun HT, Sun WQ, Ji Y, Jia SJ
- 7060** Myotonic dystrophy type 1 presenting with dyspnea: A case report
Jia YX, Dong CL, Xue JW, Duan XQ, Xu MY, Su XM, Li P
- 7068** Novel mutation in the *SALL1* gene in a four-generation Chinese family with uraemia: A case report
Fang JX, Zhang JS, Wang MM, Liu L
- 7076** Malignant transformation of primary mature teratoma of colon: A case report
Liu J

- 7082** Treatment of pyogenic liver abscess by surgical incision and drainage combined with platelet-rich plasma: A case report
Wang JH, Gao ZH, Qian HL, Li JS, Ji HM, Da MX
- 7090** Left bundle branch pacing in a ventricular pacing dependent patient with heart failure: A case report
Song BX, Wang XX, An Y, Zhang YY
- 7097** Solitary fibrous tumor of the liver: A case report and review of the literature
Xie GY, Zhu HB, Jin Y, Li BZ, Yu YQ, Li JT
- 7105** MutL homolog 1 germline mutation c.(453+1_454-1)_(545+1_546-1)del identified in lynch syndrome: A case report and review of literature
Zhang XW, Jia ZH, Zhao LP, Wu YS, Cui MH, Jia Y, Xu TM
- 7116** Malignant histiocytosis associated with mediastinal germ cell tumor: A case report
Yang PY, Ma XL, Zhao W, Fu LB, Zhang R, Zeng Q, Qin H, Yu T, Su Y
- 7124** Immunoglobulin G4 associated autoimmune cholangitis and pancreatitis following the administration of nivolumab: A case report
Agrawal R, Guzman G, Karimi S, Giulianotti PC, Lora AJM, Jain S, Khan M, Boulay BR, Chen Y
- 7130** Portal vein thrombosis in a noncirrhotic patient after hemihepatectomy: A case report and review of literature
Zhang SB, Hu ZX, Xing ZQ, Li A, Zhou XB, Liu JH
- 7138** Microvascular decompression for a patient with oculomotor palsy caused by posterior cerebral artery compression: A case report and literature review
Zhang J, Wei ZJ, Wang H, Yu YB, Sun HT
- 7147** Topical halometasone cream combined with fire needle pre-treatment for treatment of primary cutaneous amyloidosis: Two case reports
Su YQ, Liu ZY, Wei G, Zhang CM
- 7153** Simultaneous robot-assisted approach in a super-elderly patient with urothelial carcinoma and synchronous contralateral renal cell carcinoma: A case report
Yun JK, Kim SH, Kim WB, Kim HK, Lee SW
- 7163** Nursing a patient with latent autoimmune diabetes in adults with insulin-related lipodystrophy, allergy, and exogenous insulin autoimmune syndrome: A case report
He F, Xu LL, Li YX, Dong YX
- 7171** Incidental diagnosis of medullary thyroid carcinoma due to persistently elevated procalcitonin in a patient with COVID-19 pneumonia: A case report
Saha A, Mukhopadhyay M, Paul S, Bera A, Bandyopadhyay T
- 7178** Macular hole following phakic intraocular lens implantation: A case report
Li XJ, Duan JL, Ma JX, Shang QL

LETTER TO THE EDITOR

7184 Is every microorganism detected in the intensive care unit a nosocomial infection? Isn't prevention more important than detection?

Yildirim F, Karaman I, Yildirim M

ABOUT COVER

Editorial Board Member of *World Journal of Clinical Cases*, Jie-Feng Huang, PhD, Associate Chief Physician, Associate Professor, Department of Orthopaedics and Traumatology, The First Affiliated Hospital of Zhejiang Chinese Medical University, Hangzhou 310006, Zhejiang Province, China. 40983285@qq.com

AIMS AND SCOPE

The primary aim of *World Journal of Clinical Cases* (*WJCC*, *World J Clin Cases*) is to provide scholars and readers from various fields of clinical medicine with a platform to publish high-quality clinical research articles and communicate their research findings online.

WJCC mainly publishes articles reporting research results and findings obtained in the field of clinical medicine and covering a wide range of topics, including case control studies, retrospective cohort studies, retrospective studies, clinical trials studies, observational studies, prospective studies, randomized controlled trials, randomized clinical trials, systematic reviews, meta-analysis, and case reports.

INDEXING/ABSTRACTING

The *WJCC* is now indexed in Science Citation Index Expanded (also known as SciSearch®), Journal Citation Reports/Science Edition, Scopus, PubMed, and PubMed Central. The 2021 Edition of Journal Citation Reports® cites the 2020 impact factor (IF) for *WJCC* as 1.337; IF without journal self cites: 1.301; 5-year IF: 1.742; Journal Citation Indicator: 0.33; Ranking: 119 among 169 journals in medicine, general and internal; and Quartile category: Q3. The *WJCC*'s CiteScore for 2020 is 0.8 and Scopus CiteScore rank 2020: General Medicine is 493/793.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: *Hua-Ge Yan*; Production Department Director: *Xu Guo*; Editorial Office Director: *Jin-Lei Wang*.

NAME OF JOURNAL

World Journal of Clinical Cases

ISSN

ISSN 2307-8960 (online)

LAUNCH DATE

April 16, 2013

FREQUENCY

Thrice Monthly

EDITORS-IN-CHIEF

Bao-Gan Peng, Jerzy Tadeusz Chudek, George Kontogeorgos, Maurizio Serati, Ja Hyeon Ku

EDITORIAL BOARD MEMBERS

<https://www.wjgnet.com/2307-8960/editorialboard.htm>

PUBLICATION DATE

July 16, 2022

COPYRIGHT

© 2022 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjgnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>

A 7-year-old boy with recurrent cyanosis and tachypnea: A case report

Shu Li, Li-Na Chen, Lin Zhong

Specialty type: Pediatrics**Provenance and peer review:**

Unsolicited article; Externally peer reviewed.

Peer-review model: Single blind**Peer-review report's scientific quality classification**

Grade A (Excellent): A

Grade B (Very good): B

Grade C (Good): C

Grade D (Fair): 0

Grade E (Poor): 0

P-Reviewer: Oh SH, South Korea;

Taheri S, Iran; Taheri S, Iran

Received: October 31, 2021

Peer-review started: October 31, 2021

First decision: December 27, 2021

Revised: January 10, 2022

Accepted: May 13, 2022

Article in press: May 13, 2022

Published online: July 16, 2022



Shu Li, Li-Na Chen, Lin Zhong, Department of Pediatric Pulmonology and Immunology, West China Second University Hospital of Sichuan University, Chengdu 610066, Sichuan Province, China

Shu Li, Li-Na Chen, Lin Zhong, Key Laboratory of Birth Defects and Related Diseases of Women and Children, Ministry of Education, West China Second University Hospital of Sichuan University, Chengdu 610066, Sichuan Province, China

Shu Li, Li-Na Chen, Lin Zhong, NHC Key Laboratory of Chronobiology, Sichuan University, Chengdu 610066, Sichuan Province, China

Corresponding author: Lin Zhong, MD, Attending Doctor, Department of Pediatric Pulmonology and Immunology, West China Second University Hospital of Sichuan University, No. 1416 Chenglong Road, Chengdu 610066, Sichuan Province, China. linca_zhong@163.com

Abstract

BACKGROUND

Brain tumors are the most common solid tumors in children and comprise 25% of all malignancies in children. Common presentations include headache, nausea and vomiting, gait abnormality, papilledema, and epileptic seizure; however, some symptoms can be very insidious, with atypical and misleading manifestations.

CASE SUMMARY

Here, we report a 7-year-old boy who presented with recurrent cyanosis and tachypnea after exercise for 2 years. His body mass index was 26.43 kg/m². Hepatosplenomegaly, blood gas analysis, biochemical parameters, chest computed tomography scan, and echocardiograph suggested type II respiratory failure, pulmonary heart disease, and mild liver injury. Non-invasive breathing support, antibiotics, and anti-heart failure therapy were given. The patient's pulse oxygen saturation increased to over 95% when he was awake but dropped to 50%-60%, accompanied by cyanosis, during sleep while receiving high-flow nasal cannula oxygen. Sleep-related breathing disorder was suspected. In the intensive care unit, however, polysomnography was unavailable. Brain magnetic resonance imaging revealed a space-occupying (cerebellum and brainstem) lesion, which was later confirmed to be pleomorphic xanthoastrocytoma by surgery and histopathology by tissue biopsy.

CONCLUSION

When treating patients with cyanosis and tachypnea, a broad differential diagnosis should be considered, including brain tumor.

Key Words: Respiratory failure; Pulmonary heart disease; Sleep-related breathing disorders; Pleomorphic xanthoastrocytoma; Children; Case report

©The Author(s) 2022. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: Brain tumors are the most common solid tumors in children. Common presentations include headache, nausea and vomiting, gait abnormality, and epileptic seizure; however, some symptoms can be insidious and misleading. Here, we report a 7-year-old boy who presented with recurrent cyanosis and tachypnea after exercise for 2 years. Type II respiratory failure, pulmonary heart disease, liver injury, and severe obesity were diagnosed. Cyanosis during sleep (peripheral oxygen saturation dropping to 50%-60%) was observed during hospitalization. Sleep-related breathing disorder was suspected. Brain magnetic resonance imaging revealed a cerebellum and brainstem space-occupying lesion. Subsequent tissue biopsy pathology confirmed brainstem pleomorphic xanthoastrocytoma.

Citation: Li S, Chen LN, Zhong L. A 7-year-old boy with recurrent cyanosis and tachypnea: A case report. *World J Clin Cases* 2022; 10(20): 6974-6980

URL: <https://www.wjgnet.com/2307-8960/full/v10/i20/6974.htm>

DOI: <https://dx.doi.org/10.12998/wjcc.v10.i20.6974>

INTRODUCTION

Central nervous system tumors are the most common solid tumor encountered in pediatric practice, reportedly accounting for approximately 25% of all childhood cancers and classified as the second most common pediatric malignancy, after leukemia[1]. Common presentations include headache (33%), nausea and vomiting (32%), gait abnormality (27%), papilledema (13%), epileptic seizures (13%), and manifestations of increased intracranial pressure (10%). Other symptoms that may be present are strabismus, altered behavior, macrocephaly, and cranial nerve palsy.

Pleomorphic xanthoastrocytoma (PXA) is a rare type of brain astrocytoma that is typically seen in children and young adolescents. While PXAs occur almost exclusively supratentorially (98% of cases) and predominantly in the temporal lobes (50%)[2], they can form in the cerebellum, brain stem, and spinal cord. In most cases, regardless of location, the initial symptom is epilepsy.

Here, we report a case of PXA that presented in a rare location and with atypical clinical manifestations, which could be misleading in clinical practice.

CASE PRESENTATION

Chief complaints

A 7-year-old boy was admitted to our hospital to address recurrent cyanosis and tachypnea on exertion that had existed for 2 years.

History of present illness

The patient had been in his usual state of health until 2 years before admission, when intermittent facial cyanosis developed that worsened on exertion, accompanied by sweating, tachypnea, and swelling of the face and bilateral legs. Systemic pain developed, which mainly affected his chest, back, and lower limbs. The episode of pain was unclear and would resolve spontaneously. The patient had occasional dry cough without fever. He had previously been treated in a local hospital, but recurrent cyanosis and tachypnea after exercise persisted for the last 2 years.

One month before admission, the patient had one suspected episode of syncope, which manifested as falling to the ground without response to voice and had lasted for 2-3 min. He was subsequently sent to the Emergency Department of our hospital. His eyes did not appear in a staring state (*i.e.* no episodes of staring blankly), and he presented neither limb rigidity nor urinary incontinence. History of trauma or surgery was denied. The cranial computed tomography (CT) findings were normal. The echocardiography revealed enlargement of the right ventricle (17 mm), slight broadening of the pulmonary artery (25 mm), and normal left ventricle function (estimated ejection fraction of 70%). Tricuspid regurgitation (maximum velocity of 3.8 m/s) was also found, and pulmonary artery hypertension was suggested

(pressure gradient = 62 mmHg). Chest CT (Figure 1A) revealed bilateral pulmonary opacities, pulmonary congestion, slight enlargement of the cardiac silhouette, broadening of the pulmonary artery, mild pericardial effusion, and localized thickening of the bilateral pleural. Cefdinir, diuretics, and nifedipine were prescribed. Cyanosis and tachypnea were relieved. Unfortunately, although his symptoms had been relieved, they relapsed after withdrawal of the medications.

One week before admission, the tachypnea worsened, with fatigue, and the patient was again evaluated in our hospital. Upon admission, the white blood cell count was $10.5 \times 10^9/L$ (reference range: $3.6-9.7 \times 10^9/L$) and 71.2% of the cells were neutrophils (reference range: 23.6%-75.0%). In addition, the level of C-reactive protein was 10.9 mg/L (reference range: 0-8 mg/L) and the level of myocardial troponin I was 0.067 mg/L (reference range: 0-0.034 mg/L). A chest radiograph showed pulmonary congestion and enlarged cardiac silhouette. A diagnosis of cardiac insufficiency and pulmonary hypertension was made. Dopamine was administered intravenously (5 mg/kg/min). Therapy was started with vitamin C, hydrochlorothiazide, spironolactone, and captopril.

History of past illness

The patient had a history of recurrent respiratory infection and myoclonic jerks and snoring during sleep at night.

Personal and family history

The patient was born at full-term, with a birth weight of 2.9 kg. The family history was unremarkable.

Physical examination

On examination, the patient appeared critically ill. The temperature was 36.5 °C, pulse was 118 beats per min, blood pressure was 118/60 mmHg, respiration was 24, and oxygen saturation was 60%-70% while he was receiving oxygen through nasal cannula. His weight was 40 kg (> 97th percentile) and body mass index (BMI) was 26.43 kg/m². Orthopnea, nasal flaring, stretches of erythema on both cheeks, and soybean-sized cervical lymph nodes on both sides were present. Cyanosis was seen on his facial area and limbs. Coarse breath sounds with crackles were heard. The heart rhythm was strong and regular, with an augmented second heart sound; no cardiac murmur was heard. The abdomen was soft and nontender. The liver extended about 2 cm below the right costal margin with a moderate texture, and the spleen was 4 cm below the left costal margin. Muscle tone was normal. Notably, he had no clubbing, edema, or jugular vein distention. Other physical examination findings were unremarkable.

Laboratory examinations

Laboratory tests indicated impaired liver function, hypoxemia, and hypercapnia (Table 1). The 1,3-beta-D glucan assay and tuberculosis T-SPOT test were negative. The culture of sputum was unremarkable. An electrocardiogram obtained on admission day showed the following: Nodal tachycardia; axis leaning right (+ 130°); enlargement of the right atrium; ST-segment depression of 0.5 mv in leads II, III, arteriovenous fistula (AVF), and V1-V3; and T-wave depression in leads II, III, AVF, and V4-V5. Spirometry showed mild restrictive ventilation disturbance. The ratio of measured values to predicted values of forced vital capacity and forced expiratory volume in 1 s/forced vital capacity were 75.1% and 97.6%, respectively. Small airway obstruction was present (forced expiratory flow at 50% was 64.7%, FEF at 75% was 36.7%, and mid maximal expiratory flow 75/25 was 53.2%).

Imaging examinations

Chest CT (Figure 1B) revealed improvement of opacities, but the enlarged cardiac silhouette was persistent.

FINAL DIAGNOSIS

PXA, hypercapnic respiratory failure, pulmonary heart disease, liver injury, and severe obesity.

TREATMENT

After hospitalization, the patient was given oxygen therapy through a mask (3 L/min). Cefoperazone tazobactam, dopamine, and reduced glutathione were infused. Diuretics and captopril were administered orally. Mild facial cyanosis was still visible, and blood gas analysis indicated an abnormally elevated partial pressure of carbon dioxide (PCO₂) of 70 mmHg (9.3 kPa). The patient was transferred to the pediatric intensive care unit (PICU).

In the PICU, high-flow nasal cannula oxygen (flow 12 L/min, fraction of inspired oxygen 50%) was administered, and the pulse oxygen saturation (SaO₂) increased to over 95% when he was awake. Meanwhile, his liver function returned to normal level, and pulmonary artery pressure estimated by

Table 1 Laboratory data

Variable	Reference range	On admission	4 th hospital day
Complete blood count			
White cell count, 10 ⁹ /L	3.6-9.7	10.5	NA
Neutrophils, %	23.6-75	71.2	NA
Lymphocytes, %	19-64	21.7	NA
Hemoglobin, g/L	110-146	137	NA
Platelet count, 10 ⁹ /L	100-450	187	NA
CRP, mg/L	0-8	10.9	NA
Blood gas analysis			
pH	7.35-7.45	7.37	7.41
SaO ₂ , %	90.0-98.0	89.6	97.7
PO ₂ , kPa	10-14	7.6	11.3
PCO ₂ , kPa	4.8-5.9	7.5	9.3
HCO ₃ ⁻ , mmol/L	21-28	32.4	44.4
BE, mmol/L	-3-3	5.4	15.3
Na ⁺ , mmol/L	135.0-145.0	133	132
K ⁺ , mmol/L	3.5-5.5	5.1	4.3
Ca ²⁺ , mmol/L	1.09-1.30	0.99	1.12
AG, mmol/L	10-18	8.0	3.0
Organ biochemical parameters			
ALT, U/L	21-72	147	65
AST, U/L	17-59	134	44
LDH, U/L	313-618	1173	NA
Creatine, mmol/L	17.3-54.6	31	NA
Troponin I, mg/L	0-0.034	0.067	NA


AG: Anion gap; ALT: Alanine aminotransferase; AST: Aspartate transaminase; BE: Base excess; Ca²⁺: Calcium; CRP: C-reactive protein; HCO₃⁻: Bicarbonate; K⁺: Sodium; LDH: Lactate dehydrogenase; NA: Not available; Na⁺: Sodium; PCO₂: Partial pressure of carbon dioxide; PO₂: Partial pressure of oxygen; SaO₂: Pulse oxygen saturation.

echocardiography decreased. However, his SaO₂ decreased to 50%-60% during sleep, accompanied by cyanosis. Intracranial abnormality was considered, and magnetic resonance imaging was performed (Figure 2). A space-occupying lesion was seen at the right cerebellopontine angle, cerebellar tonsil, and medulla oblongata with partial involvement of the cervical spinal cord. The maximum cross-sectional area of the lesion was 1.7 cm × 2.3 cm with slightly hyperintense signal on T2-weighted imaging and slightly hypointense signal on T1-weighted imaging. The lesion showed significantly inhomogeneous enhancement. The pituitary body and the meninges were clean.

The patient was then transferred for neurosurgery. Histopathological examination of the resected brain tissue indicated PXA with the presence of *BRAF* V600E mutation, which is the most common form of gene mutation in PXA.


OUTCOME AND FOLLOW-UP

After neurosurgery, the patient's symptoms resolved transiently but gradually relapsed. Six months later, the boy died of respiratory failure.



DOI: 10.12998/wjcc.v10.i20.6974 Copyright ©The Author(s) 2022.

Figure 1 Computed tomography scan viewed in the lung window. A: Infiltrates and ground-glass opacities on both lungs were seen 1 mo before admission; B: On admission day, an improvement was seen in those changes after outpatient treatment.



DOI: 10.12998/wjcc.v10.i20.6974 Copyright ©The Author(s) 2022.

Figure 2 Magnetic resonance T2-weighted imaging. A: Sagittal scan showed a space-occupying a lesion with slightly hyperintense signal at the right cerebellopontine angle, cerebellar tonsil, and medulla oblongata without a clear boundary; B: Sagittal scan showing a space-occupying lesion with inhomogeneous enhancement; C: Transverse image showing a space-occupying lesion with slightly hyperintense signal at the right cerebellopontine angle and the cerebellar tonsil. White and black arrows indicate the space-occupying lesion of the cerebellum and brainstem.

DISCUSSION

The common causes of cyanosis and tachypnea in children are respiratory infections and heart diseases, with liver diseases as a less frequent cause. Here, we report a rare case of cyanosis and tachypnea that was mainly caused by a brain tumor, which was ultimately confirmed by histopathology to be PXA.

PXA generally has a favorable prognosis, with many cases being cured by surgery alone. The postoperative survival rates at 5 years and 10 years are 81% and 70%, respectively. Nevertheless, it is recognized that 15% to 20% of PXAs will undergo malignant progression. Because of the potential for malignant degeneration and the less predictable long-term prognosis, PXA has been classified as a World Health Organization grade II tumor malignancy[3], meaning that the tumor grows slowly and usually does not spread but is prone to recur postsurgery.

An underlying sleep-related breathing disorder (SRBD) was suggested by the observations of previous history of myoclonic jerks and snoring during sleep, decreased SaO₂ during sleep accompanied by cyanosis, and severe obesity. SRBDs have the potential to increase pulmonary arterial pressure during sleep but also during wakefulness in the long term[4]. Without appropriate intervention, SRBD progressively leads to pulmonary hypertension, pulmonary heart disease, and even death. In this case, several types of SRBD were considered that might have accounted for the disorder. Discriminating among these categories requires the performance of polysomnography (PSG).

Given the coexistence of elevated level of CO₂, alveolar hypoventilation syndrome was the most likely cause. It is defined as insufficient ventilation with elevated level of PCO₂ (in children PCO₂ > 50 mmHg, accounting for more than 25% of the total sleep time), either by direct determination of arterial blood gas or, more commonly, by proxy measures such as end-tidal or transcutaneous CO₂[5]. Alveolar hypoventilation syndrome is caused by some congenital or acquired conditions, including congenital

central hypoventilation syndrome (CCHS), acquired alveolar hypoventilation associated with central nervous system abnormality, neuromuscular, rib cage disorders, obesity, and certain medication intake [6].

Another type of SRBD that should be considered is acquired central alveolar hypoventilation, which is very rare and can be caused by brainstem damage resulting from tumor, ischemic damage, or infection[7]. The patient had PXA confirmed by surgery and pathological examination. The tumor was a focal glioma located at the cerebellum and brainstem, without a clear boundary. In this case, both the clinical manifestation and the location were rare. *BRAF* V600E mutation was found in the patient, which encodes an intracellular component of the mitogen-activated protein kinase pathway and has a high oncogenic potential. *BRAF* mutations can be found in 70% of typical PXA cases[8].

Except for the above two types of SRBD, obesity hypoventilation syndrome should also be taken into consideration. It is defined as a combination of obesity (BMI > 30 kg/m² in adults; > P₉₅ in children), daytime hypercapnia (PCO₂ > 45 mmHg), and sleep-disordered breathing, after ruling out other disorders that may cause alveolar hypoventilation, such as congenital central hypoventilation syndrome, sleep-related hypoventilation due to a medication or substance, and sleep-related hypoventilation due to a medical disorder. In this case, the patient's BMI was 26.43 kg/m², above the P₉₇ level (19.4 kg/m² in 7-year-old Chinese boys)[9], and blood gas analysis revealed PCO₂ of 56.5 mmHg and 70.1 mmHg during wakefulness, implicating obesity in the development of alveolar hypoventilation.

We believe that brain tumor and obesity may have both been implicated in the pathogenesis of alveolar hypoventilation in this case. The coexistence of other types of SRBD, such as central apnea syndrome or obstructive sleep apnea-hypopnea syndrome, might also occur, which cannot be ruled out without PSG examination.

Other diseases presenting with dyspnea and cyanosis should be differentiated. A potential differential diagnosis is hepatopulmonary syndrome (HPS), since the patient presents with elevated liver enzymes and hypoxemia. HPS is characterized by abnormalities in blood oxygenation caused by the presence of intrapulmonary vascular dilations (IPVD) in the context of liver disease, generally at the cirrhotic stage[10]. HPS comprises the presence of advanced chronic liver disease, gas exchange abnormalities, and the presence of IPVD, without intrinsic pulmonary diseases. The patient in this case did not have a previous history of liver disease. On admission, the liver extended about 2 cm below the right costal margin with a moderate texture and the spleen 4 cm below the left costal margin. The liver enzymes were slightly elevated (alanine aminotransferase at 147 U/L; aspartate transaminase at 134 U/L) with normal bilirubin. After 3 d of treatment, the liver extended about 1 cm below the right costal margin and the spleen 2 cm below the left costal margin. The liver enzyme returned to normal; thus, the liver disorder was reversible. Additionally, the chest CT showed interstitial changes and the spirometry revealed mild restrictive ventilation disturbance, which were evidence of pulmonary diseases, thereby ruling out HPS.

Arteriovenous shunts, especially pulmonary arteriovenous fistula (PAVF), is another differential diagnosis that should be considered. PAVF is an abnormal communication between the pulmonary veins and pulmonary arteries. Most individuals may have the condition since birth[11]. The common manifestations of PAVF are shortness of breath, hemoptysis, chest pain, dizziness, and syncope. Pulmonary CT angiography is the gold standard for diagnosis, presenting opacity and AVF. Although this patient had such symptoms as tachypnea, chest pain, and suspected syncope, which were similar to PAVF, he did not have hemoptysis, clubbing, or hereditary hemorrhagic telangiectasia, and the CT scans ruled out PAVF.

CCHS, previously known as Ondine's curse syndrome, is characterized by the impaired control of autonomic respiration during sleep and diminished sensitivity of CO₂ receptors. The primary cause is mutation of the *PHOXB* gene, which is found in 90% of the cases[12]. CCHS usually starts in the neonatal period or early infancy and is accompanied by neurological abnormalities, such as autonomic and neurocognitive dysfunctions. Our patient had no such manifestations, making this diagnosis untenable.

This study had some limitations. First, PSG was not conducted because the patient was critically ill, and magnetic resonance imaging had revealed a brain tumor, which necessitated immediate surgery; thus, the information on SRBD was incomplete. Second, cardiac catheterization data were lacking, so the evaluation of pulmonary artery pressure was not accurate.

CONCLUSION

This case shows that when treating patients with cyanosis and tachypnea, a broad differential diagnosis should be contemplated, including a brain tumor.

FOOTNOTES

Author contributions: Li S contributed to data collection and analyses and drafted the manuscript; Chen LN

contributed to data analyses and critical revision of the manuscript; Zhong L contributed to the conception and design and gave final approval of the manuscript to be published.

Supported by the Science and Technology Department of Sichuan Province, No. 2020YFS0105; and the West China Second University Hospital of Sichuan University, No. KL036.

Informed consent statement: Informed written consent was obtained from the patient for the publication of this report and any accompanying images.

Conflict-of-interest statement: The authors declare that they have no conflict of interest.

CARE Checklist (2016) statement: The authors have read the CARE Checklist (2016), and the manuscript was prepared and revised according to the CARE Checklist (2016).

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Country/Territory of origin: China

ORCID number: Shu Li 0000-0001-9126-1131; Li-Na Chen 0000-0001-9537-8847; Lin Zhong 0000-0003-0587-3460.

S-Editor: Guo XR

L-Editor: Filipodia

P-Editor: Chen YX

REFERENCES

- 1 **Cacciotti C**, Fleming A, Ramaswamy V. Advances in the molecular classification of pediatric brain tumors: a guide to the galaxy. *J Pathol* 2020; **251**: 249-261 [PMID: 32391583 DOI: 10.1002/path.5457]
- 2 **Andre D**, Furtado, Charles R. Fitz, Neuroimaging Part II. Handbook of Clinical Neurology (First Edition). Edinburgh: Elsevier, 2016; 136: 1139-1158
- 3 **Daniel J**, Brat, Arie Perry, 6-Astrocytic and Oligodendroglial Tumors. Practical Surgical Neuropathology: A Diagnostic Approach (Second Edition). Edinburgh: Elsevier, 2018: 91-123
- 4 **Thurnheer R**, Ulrich S, Bloch KE. Precapillary Pulmonary Hypertension and Sleep-Disordered Breathing: Is There a Link? *Respiration* 2017; **93**: 65-77 [PMID: 27884004 DOI: 10.1159/000452957]
- 5 **American Academy of sleep medicine**. International classification of sleep disorders 3rd ed, American Academy of Sleep Medicine; 2014 Sep 15; Darien, IL. 2014: 108-128
- 6 **Herkenrath SD**, Randerath WJ. More than Heart Failure: Central Sleep Apnea and Sleep-Related Hypoventilation. *Respiration* 2019; **98**: 95-110 [PMID: 31291632 DOI: 10.1159/000500728]
- 7 **Deak MC**, Kirsch DB. Sleep-disordered breathing in neurologic conditions. *Clin Chest Med* 2014; **35**: 547-556 [PMID: 25156770 DOI: 10.1016/]
- 8 **Shaikh N**, Brahmabhatt N, Kruser TJ, Kam KL, Appin CL, Wadhvani N, Chandler J, Kumthekar P, Lukas RV. Pleomorphic xanthoastrocytoma: a brief review. *CNS Oncol* 2019; **8**: CNS39 [PMID: 31535562 DOI: 10.2217/cns-2019-0009]
- 9 **Masa JF**, Pépin JL, Borel JC, Mokhlesi B, Murphy PB, Sánchez-Quiroga MÁ. Obesity hypoventilation syndrome. *Eur Respir Rev* 2019; **28** [PMID: 30872398 DOI: 10.1183/16000617.0097-2018]
- 10 **Grilo-Bensusan I**, Pascasio-Acevedo JM. Hepatopulmonary syndrome: What we know and what we would like to know. *World J Gastroenterol* 2016; **22**: 5728-5741 [PMID: 27433086 DOI: 10.3748/wjg.v22.i25.5728]
- 11 **Jha AK**, Sunder A. An unusual case of pulmonary arterio-venous fistula (PAVF). *J Family Med Prim Care* 2019; **8**: 3408-3411 [PMID: 31742178 DOI: 10.4103/jfmpe.jfmpe_580_19]
- 12 **Zaidi S**, Gandhi J, Vatsia S, Smith NL, Khan SA. Congenital central hypoventilation syndrome: An overview of etiopathogenesis, associated pathologies, clinical presentation, and management. *Auton Neurosci* 2018; **210**: 1-9 [PMID: 29249648 DOI: 10.1016/j.autneu.2017.11.003]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA
Telephone: +1-925-3991568
E-mail: bpgoffice@wjgnet.com
Help Desk: <https://www.f6publishing.com/helpdesk>
<https://www.wjgnet.com>

