

# World Journal of *Clinical Cases*

*World J Clin Cases* 2022 September 26; 10(27): 9550-9969



## Contents

Thrice Monthly Volume 10 Number 27 September 26, 2022

## OPINION REVIEW

- 9550 Psychiatric disorders and pain: The recurrence of a comorbidity  
*Vyshka G*

## REVIEW

- 9556 Cardiovascular disease and COVID-19, a deadly combination: A review about direct and indirect impact of a pandemic  
*Vidal-Perez R, Brandão M, Pazdernik M, Kresoja KP, Carpenito M, Maeda S, Casado-Arroyo R, Muscoli S, Pöss J, Fontes-Carvalho R, Vazquez-Rodriguez JM*
- 9573 Molecular factors, diagnosis and management of gastrointestinal tract neuroendocrine tumors: An update  
*Pavlidis ET, Pavlidis TE*

## MINIREVIEWS

- 9588 Human-induced pluripotent stem cell-atrial-specific cardiomyocytes and atrial fibrillation  
*Leowattana W, Leowattana T, Leowattana P*
- 9602 COVID-19 and the cardiovascular system-current knowledge and future perspectives  
*Chatzis DG, Magounaki K, Pantazopoulos I, Bhaskar SMM*

## ORIGINAL ARTICLE

## Case Control Study

- 9611 PDCA nursing in improving quality management efficacy in endoscopic submucosal dissection  
*He YH, Wang F*

## Retrospective Study

- 9619 Impact of COVID-19 pandemic on the ocular surface  
*Marta A, Marques JH, Almeida D, José D, Sousa P, Barbosa I*
- 9628 Anatomy and clinical application of suprascapular nerve to accessory nerve transfer  
*Wang JW, Zhang WB, Li F, Fang X, Yi ZQ, Xu XL, Peng X, Zhang WG*
- 9641 Therapeutic effect of two methods on avulsion fracture of tibial insertion of anterior cruciate ligament  
*Niu HM, Wang QC, Sun RZ*
- 9650 Efficacy of transcatheter arterial chemoembolization using pirarubicin-loaded microspheres combined with lobaplatin for primary liver cancer  
*Zhang C, Dai YH, Lian SF, Liu L, Zhao T, Wen JY*

- 9657** Prognostic significance of sex determining region Y-box 2, E-cadherin, and vimentin in esophageal squamous cell carcinoma

*Li C, Ma YQ*

- 9670** Clinical characteristics and prognosis of orbital solitary fibrous tumor in patients from a Chinese tertiary eye hospital

*Ren MY, Li J, Wu YX, Li RM, Zhang C, Liu LM, Wang JJ, Gao Y*

### Observational Study

- 9680** Altered heart rate variability and pulse-wave velocity after spinal cord injury

*Tsou HK, Shih KC, Lin YC, Li YM, Chen HY*

- 9693** Intra and extra pelvic multidisciplinary surgical approach of retroperitoneal sarcoma: Case series report

*Song H, Ahn JH, Jung Y, Woo JY, Cha J, Chung YG, Lee KH*

### META-ANALYSIS

- 9703** Meta-analysis of gemcitabine plus nab-paclitaxel combined with targeted agents in the treatment of metastatic pancreatic cancer

*Li ZH, Ma YJ, Jia ZH, Weng YY, Zhang P, Zhu SJ, Wang F*

- 9714** Clinical efficacy analysis of mesenchymal stem cell therapy in patients with COVID-19: A systematic review

*Cao JX, You J, Wu LH, Luo K, Wang ZX*

### CASE REPORT

- 9727** Treatment of gastric cancer with dermatomyositis as the initial symptom: Two case reports and review of literature

*Sun XF, Gao XD, Shen KT*

- 9734** Gallbladder hemorrhage—An uncommon surgical emergency: A case report

*Valenti MR, Cavallaro A, Di Vita M, Zanghi A, Longo Trischitta G, Cappellani A*

- 9743** Successful treatment of stage IIIB intrahepatic cholangiocarcinoma using neoadjuvant therapy with the PD-1 inhibitor camrelizumab: A case report

*Zhu SG, Li HB, Dai TX, Li H, Wang GY*

- 9750** Myocarditis as an extraintestinal manifestation of ulcerative colitis: A case report and review of the literature

*Wang YY, Shi W, Wang J, Li Y, Tian Z, Jiao Y*

- 9760** Endovascular treatment of traumatic renal artery pseudoaneurysm with a Stanford type A intramural haematoma: A case report

*Kim Y, Lee JY, Lee JS, Ye JB, Kim SH, Sul YH, Yoon SY, Choi JH, Choi H*

- 9768** Histiocytoid giant cellulitis-like Sweet syndrome at the site of sternal aspiration: A case report and review of literature

*Zhao DW, Ni J, Sun XL*

- 9776** Rare giant corneal keloid presenting 26 years after trauma: A case report  
*Li S, Lei J, Wang YH, Xu XL, Yang K, Jie Y*
- 9783** Efficacy evaluation of True Lift®, a nonsurgical facial ligament retightening injection technique: Two case reports  
*Huang P, Li CW, Yan YQ*
- 9790** Synchronous primary duodenal papillary adenocarcinoma and gallbladder carcinoma: A case report and review of literature  
*Chen J, Zhu MY, Huang YH, Zhou ZC, Shen YY, Zhou Q, Fei MJ, Kong FC*
- 9798** Solitary fibrous tumor of the renal pelvis: A case report  
*Liu M, Zheng C, Wang J, Wang JX, He L*
- 9805** Gastric metastasis presenting as submucosa tumors from renal cell carcinoma: A case report  
*Chen WG, Shan GD, Zhu HT, Chen LH, Xu GQ*
- 9814** Laparoscopic correction of hydronephrosis caused by left paraduodenal hernia in a child with cryptorchism: A case report  
*Wang X, Wu Y, Guan Y*
- 9821** Diagnosed corrected transposition of great arteries after cesarean section: A case report  
*Ichii N, Kakinuma T, Fujikawa A, Takeda M, Ohta T, Kagimoto M, Kaneko A, Izumi R, Kakinuma K, Saito K, Maeyama A, Yanagida K, Takeshima N, Ohwada M*
- 9828** Misdiagnosis of an elevated lesion in the esophagus: A case report  
*Ma XB, Ma HY, Jia XF, Wen FF, Liu CX*
- 9834** Diagnostic features and therapeutic strategies for malignant paraganglioma in a patient: A case report  
*Gan L, Shen XD, Ren Y, Cui HX, Zhuang ZX*
- 9845** Infant with reverse-transcription polymerase chain reaction confirmed COVID-19 and normal chest computed tomography: A case report  
*Ji GH, Li B, Wu ZC, Wang W, Xiong H*
- 9851** Pulmonary hypertension secondary to seronegative rheumatoid arthritis overlapping antisynthetase syndrome: A case report  
*Huang CY, Lu MJ, Tian JH, Liu DS, Wu CY*
- 9859** Monitored anesthesia care for craniotomy in a patient with Eisenmenger syndrome: A case report  
*Ri HS, Jeon Y*
- 9865** Emergency treatment and anesthesia management of internal carotid artery injury during neurosurgery: Four case reports  
*Wang J, Peng YM*

- 9873** Resolution of herpes zoster-induced small bowel pseudo-obstruction by epidural nerve block: A case report  
*Lin YC, Cui XG, Wu LZ, Zhou DQ, Zhou Q*
- 9879** Accidental venous port placement *via* the persistent left superior vena cava: Two case reports  
*Zhou RN, Ma XB, Wang L, Kang HF*
- 9886** Application of digital positioning guide plates for the surgical extraction of multiple impacted supernumerary teeth: A case report and review of literature  
*Wang Z, Zhao SY, He WS, Yu F, Shi SJ, Xia XL, Luo XX, Xiao YH*
- 9897** Iatrogenic aortic dissection during right transradial intervention in a patient with aberrant right subclavian artery: A case report  
*Ha K, Jang AY, Shin YH, Lee J, Seo J, Lee SI, Kang WC, Suh SY*
- 9904** Pneumomediastinum and subcutaneous emphysema secondary to dental extraction: Two case reports  
*Ye LY, Wang LF, Gao JX*
- 9911** Hemorrhagic shock due to submucosal esophageal hematoma along with mallory-weiss syndrome: A case report  
*Oba J, Usuda D, Tsuge S, Sakurai R, Kawai K, Matsubara S, Tanaka R, Suzuki M, Takano H, Shimoizawa S, Hotchi Y, Usami K, Tokunaga S, Osugi I, Katou R, Ito S, Mishima K, Kondo A, Mizuno K, Takami H, Komatsu T, Nomura T, Sugita M*
- 9921** Concurrent severe hepatotoxicity and agranulocytosis induced by *Polygonum multiflorum*: A case report  
*Shao YL, Ma CM, Wu JM, Guo FC, Zhang SC*
- 9929** Transient ischemic attack after mRNA-based COVID-19 vaccination during pregnancy: A case report  
*Chang CH, Kao SP, Ding DC*
- 9936** Drug-induced lung injury caused by acetaminophen in a Japanese woman: A case report  
*Fujii M, Kenzaka T*
- 9945** Familial mitochondrial encephalomyopathy, lactic acidosis, and stroke-like episode syndrome: Three case reports  
*Yang X, Fu LJ*
- 9954** Renal pseudoaneurysm after rigid ureteroscopic lithotripsy: A case report  
*Li YH, Lin YS, Hsu CY, Ou YC, Tung MC*

**LETTER TO THE EDITOR**

- 9961** Role of traditional Chinese medicine in the initiative practice for health  
*Li Y, Li SY, Zhong Y*
- 9964** Impact of the COVID-19 pandemic on healthcare workers' families  
*Helou M, El Osta N, Husni R*

- 9967**    Transition beyond the acute phase of the COVID-19 pandemic: Need to address the long-term health impacts of COVID-19

*Tsioutis C, Tofarides A, Spernovasilis N*

**ABOUT COVER**

Editorial Board Member of *World Journal of Clinical Cases*, Yusuf Tutar, PhD, Chairman, Director, Full Professor, Department of Basic Pharmaceutical Sciences, Division of Biochemistry, University of Health Sciences, Istanbul 34668, Turkey. ytutar@outlook.com

**AIMS AND SCOPE**

The primary aim of *World Journal of Clinical Cases* (WJCC, *World J Clin Cases*) is to provide scholars and readers from various fields of clinical medicine with a platform to publish high-quality clinical research articles and communicate their research findings online.

WJCC mainly publishes articles reporting research results and findings obtained in the field of clinical medicine and covering a wide range of topics, including case control studies, retrospective cohort studies, retrospective studies, clinical trials studies, observational studies, prospective studies, randomized controlled trials, randomized clinical trials, systematic reviews, meta-analysis, and case reports.

**INDEXING/ABSTRACTING**

The WJCC is now abstracted and indexed in Science Citation Index Expanded (SCIE, also known as SciSearch®), Journal Citation Reports/Science Edition, Current Contents®/Clinical Medicine, PubMed, PubMed Central, Scopus, Reference Citation Analysis, China National Knowledge Infrastructure, China Science and Technology Journal Database, and Superstar Journals Database. The 2022 Edition of Journal Citation Reports® cites the 2021 impact factor (IF) for WJCC as 1.534; IF without journal self cites: 1.491; 5-year IF: 1.599; Journal Citation Indicator: 0.28; Ranking: 135 among 172 journals in medicine, general and internal; and Quartile category: Q4. The WJCC's CiteScore for 2021 is 1.2 and Scopus CiteScore rank 2021: General Medicine is 443/826.

**RESPONSIBLE EDITORS FOR THIS ISSUE**

Production Editor: *Ying-Yi Yuan*; Production Department Director: *Xiang Li*; Editorial Office Director: *Jin-Lei Wang*.

**NAME OF JOURNAL**

*World Journal of Clinical Cases*

**ISSN**

ISSN 2307-8960 (online)

**LAUNCH DATE**

April 16, 2013

**FREQUENCY**

Thrice Monthly

**EDITORS-IN-CHIEF**

Bao-Gan Peng, Jerzy Tadeusz Chudek, George Kontogeorgos, Maurizio Serati, Ja Hyeon Ku

**EDITORIAL BOARD MEMBERS**

<https://www.wjgnet.com/2307-8960/editorialboard.htm>

**PUBLICATION DATE**

September 26, 2022

**COPYRIGHT**

© 2022 Baishideng Publishing Group Inc

**INSTRUCTIONS TO AUTHORS**

<https://www.wjgnet.com/bpg/gerinfo/204>

**GUIDELINES FOR ETHICS DOCUMENTS**

<https://www.wjgnet.com/bpg/GerInfo/287>

**GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH**

<https://www.wjgnet.com/bpg/gerinfo/240>

**PUBLICATION ETHICS**

<https://www.wjgnet.com/bpg/GerInfo/288>

**PUBLICATION MISCONDUCT**

<https://www.wjgnet.com/bpg/gerinfo/208>

**ARTICLE PROCESSING CHARGE**

<https://www.wjgnet.com/bpg/gerinfo/242>

**STEPS FOR SUBMITTING MANUSCRIPTS**

<https://www.wjgnet.com/bpg/GerInfo/239>

**ONLINE SUBMISSION**

<https://www.f6publishing.com>

## Endovascular treatment of traumatic renal artery pseudoaneurysm with a Stanford type A intramural haematoma: A case report

Yook Kim, Jin Young Lee, Jin Suk Lee, Jin-Bong Ye, Se Heon Kim, Young Hoon Sul, Su Young Yoon, Jung Hee Choi, Hanlim Choi

**Specialty type:** Surgery

**Provenance and peer review:**

Unsolicited article; Externally peer reviewed.

**Peer-review model:** Single blind

**Peer-review report's scientific quality classification**

Grade A (Excellent): 0  
Grade B (Very good): B  
Grade C (Good): C, C, C, C  
Grade D (Fair): 0  
Grade E (Poor): 0

**P-Reviewer:** Gálvez Salazar P, Ecuador; Lovi NK, Ghana; Tan B, China; Yang L, China

**Received:** May 4, 2022

**Peer-review started:** May 4, 2022

**First decision:** June 16, 2022

**Revised:** June 28, 2022

**Accepted:** August 21, 2022

**Article in press:** August 21, 2022

**Published online:** September 26, 2022



**Yook Kim,** Department of Radiology, Chungbuk National University Hospital, Cheongju-si 28644, Chungcheongbuk-do, South Korea

**Jin Young Lee, Jin Suk Lee, Jin-Bong Ye, Se Heon Kim,** Department of Trauma Surgery, Chungbuk National University Hospital, Cheongju-si 28644, Chungcheongbuk-do, South Korea

**Young Hoon Sul,** Department of Trauma Surgery, College of Medicine, Chungbuk National University, Cheongju-si 28644, Chungcheongbuk-do, South Korea

**Su Young Yoon,** Cardiovascular and Thoracic Surgery, Chungbuk National University Hospital, Cheongju-si 28644, Chungcheongbuk-do, South Korea

**Jung Hee Choi,** Anesthesiology and Pain Medicine, Chungbuk National University Hospital, Cheongju-si 28644, Chungcheongbuk-do, South Korea

**Hanlim Choi,** Department of Surgery, Chungbuk National University Hospital, Cheongju-si 28644, Chungcheongbuk-do, South Korea

**Corresponding author:** Jin Young Lee, Doctor, Surgeon, Department of Trauma Surgery, Chungbuk National University Hospital, 776 1Sunhwan-ro, Seowon-gu, Cheongju-si 28644, Chungcheongbuk-do, South Korea. [client80@hanmail.net](mailto:client80@hanmail.net)

### Abstract

#### BACKGROUND

Aortic intramural hematoma (IMH) associated with aortic branch tear and intramurally located pseudoaneurysm after blunt trauma has not been reported. Here, we report a case of progressive type A aortic IMH associated with a pseudoaneurysm arising from the injured proximal renal artery after blunt trauma.

#### CASE SUMMARY

During logging operations, a 66-year-old man experienced blunt force trauma after being injured by a fallen tree. He arrived at our trauma center with a left flank pain complaint. Computed tomography (CT) revealed a pseudoaneurysm arising from the proximal renal artery (localized within the aortic media) and Stanford type A IMH. A covered stent was deployed along the left main renal artery, bridging the pseudoaneurysm and covering the parent artery, successfully excluding the pseudoaneurysm as confirmed using aortography. However,



although the degree of the pseudoaneurysm decreased, follow-up CT revealed remnant pseudoaneurysm, likely caused by an endoleak. Subsequently, a covered stent was additionally installed through the previously deployed covered stent. Successful exclusion of the pseudoaneurysm was confirmed using final aortography. In the 7-mo follow-up CT scan, the IMH and pseudoaneurysm completely disappeared with no evidence of stent-related complications.

### CONCLUSION

Endovascular treatment such as stent-graft placement can be an effective and safe treatment for traumatic renal artery injury.

**Key Words:** Aortic intramural hematoma; Blunt trauma; Intimomedial tear; Renal arterial injury; Stent-graft; Case report

©The Author(s) 2022. Published by Baishideng Publishing Group Inc. All rights reserved.

**Core Tip:** We report the case of an adult male who presented with left flank pain after experiencing blunt trauma from being injured by a fallen tree. Investigations revealed a pseudoaneurysm arising from the proximal renal artery and Stanford type A IMH. The patient received endovascular treatment, and the successful exclusion of the pseudoaneurysm was confirmed using final aortography. A postoperative computed tomography scan showed the complete disappearance of the intramural hematoma and pseudoaneurysm with no stent-related complications. Consequently, we propose that endovascular treatment such as stent-graft placement can be an effective and safe treatment for traumatic renal artery injury.

**Citation:** Kim Y, Lee JY, Lee JS, Ye JB, Kim SH, Sul YH, Yoon SY, Choi JH, Choi H. Endovascular treatment of traumatic renal artery pseudoaneurysm with a Stanford type A intramural haematoma: A case report. *World J Clin Cases* 2022; 10(27): 9760-9767

**URL:** <https://www.wjgnet.com/2307-8960/full/v10/i27/9760.htm>

**DOI:** <https://dx.doi.org/10.12998/wjcc.v10.i27.9760>


## INTRODUCTION

Aortic intramural hematoma (IMH) is classically defined as a hematoma in the aortic wall, secondary to rupture of the vasa vasorum, and it is distinguished from aortic dissection (AD) by the absence on imaging of any evident intimal tear[1,2]. However, improved thin-section multidetector computed tomography (CT) images and surgical reports have confirmed small intimomedial tears in a variable percentage of patients diagnosed with IMH[1-4]. Recently, acute aortic syndrome (AAS) was defined as AD, IMH, and penetrating aortic ulcer (PAU)[5]. AAS is characterized by disruption of the media layer of the aorta with bleeding within IMH, along the aortic wall resulting in dissection, or transmurally through the wall in the case of ruptured PAU or trauma[5]. A total of 94% of the reported IMHs had a nontraumatic cause[2,4-6]. The natural history of IMH is highly variable; it may resolve or progress to aneurysm, dissection, or rupture. The morbidity and mortality caused by aortic IMH are similar to that of AD[5,7]. Uncomplicated IMH can be managed conservatively with antihypertensive medication[5]. Conversely, prompt surgical or endovascular intervention is required when IMH cases have significant predictors or symptoms of complications. The type of endovascular intervention depends on the Stanford classification, maximum aortic diameter, maximum IMH thickness, and intramural focal contrast enhancement[1,2,5,7]. To the best of our knowledge, aortic IMH caused by aortic branch artery tear with intramurally located pseudoaneurysm after blunt trauma has not been reported. Here, we report a case of progressive type A aortic IMH associated with a pseudoaneurysm arising from the injured proximal renal artery after blunt trauma, successfully managed using endovascular treatment.

## CASE PRESENTATION

### Chief complaints

A 66-year-old man arrived at our trauma center with a left flank pain complaint.



DOI: 10.12998/wjcc.v10.i27.9760 Copyright ©The Author(s) 2022.

**Figure 1** Initial computed tomography scans taken 1 h (A and B) after injury, and follow-up CT scans taken 15 d (C and D) after injury. A: The initial abdominal computed tomography (CT) scan also revealed focal out-pouching enhancing nodular lesion, suggesting a pseudoaneurysm (red arrow) arising from the proximal portion of the left renal artery with retroperitoneal hematoma (red asterisk); B: The initial chest CT scan taken one hour after injury showed Stanford type A intramural hematoma (IMH) (red arrowheads), which represent slightly hyperdensity on precontrast image (figure in red-border box); C: The CT scan reveals an increase in the size and extent of the pseudoaneurysm (red arrow) on the axial image; D: The pseudoaneurysm arising from the proximal renal artery (red arrowhead) was localized within aortic media (red asterisk), resulting in an intramural blood collection (red arrows) seen on the sagittal image.

### **History of present illness**

The patient's symptoms started 1 h after being injured by a fallen tree during logging operations.

### **History of past illness**

The patient's previous medical history was not significant.

### **Personal and family history**

The patient's family history was not significant.

### **Physical examination**


The patient was aerodynamically stable with a blood pressure of 19.3/11.3 kPa, heart rate of 72 beats per min, respiratory rate of 24 breaths per min, and body temperature of 36.0. There was no evidence of gross hematuria.

### **Laboratory examinations**

Laboratory findings were normal, including hemoglobin level, electrolytes, liver function, kidney function, and coagulation profile.

### **Imaging examinations**

Stanford type A IMH was detected on a chest CT scan (Figure 1A), and an abdominal CT scan revealed a focal out-pouching nodular lesion, suggesting the presence of a pseudoaneurysm arising from the proximal left renal artery with retroperitoneal hematoma (Figure 1B). We decided on conservative management by controlling systolic blood pressure to < 16.0 kPa due to the stable vital signs and small size of the pseudoaneurysm (< 0.5 cm) on the initial CT scan. However, a follow-up CT scan (axial image) taken 15 d after the initial CT scan revealed an increase in the size and extent of the pseudoaneurysm (Figure 1C), which was localized within the aortic media, resulting in an intramural blood collection (Figure 1D). In addition, the maximum diameter of the type A aortic IMH also increased on



DOI: 10.12998/wjcc.v10.i27.9760 Copyright ©The Author(s) 2022.

**Figure 2 Endovascular procedure performed 15 d and 17 d after injury.** A: The pseudoaneurysm (red arrow) arising from the left proximal renal artery was detected using abdominal aortography 15 d after injury; B: An 8 mm × 55 mm covered stent was deployed along the left renal main artery, bridging the pseudoaneurysm and covering the parent artery, successfully excluding the pseudoaneurysm; exclusion was confirmed using final aortography; C: Follow-up CT scan two days after the first procedure reveals a decrease in the size of the pseudoaneurysm (red arrow) arising from the renal artery; D: Second endovascular procedure performed three days after the first treatment. The additional covered stent (red arrows) was placed to fully cover the orifice of the left renal artery to prevent 1A endoleak, successfully excluding the pseudoaneurysm, as confirmed using final aortography.

the follow-up chest CT scan taken 15 d after the initial scan.

## FINAL DIAGNOSIS

The final diagnosis of the presented case is progressive type A IMH associated with renal artery injury after blunt trauma.

## TREATMENT

Considering the aggravated pseudoaneurysm and progressive IMH, an emergent endovascular procedure was performed immediately. The pseudoaneurysm arising from the left proximal renal artery was detected using abdominal aortography (Figure 2A), corresponding to the CT images. We opted for complete exclusion of the pseudoaneurysm with a covered stent to manage the pseudoaneurysm while preserving the blood flow to the kidney. Therefore, an 8 mm × 55 mm covered stent (Hercules; S&G Biotech Inc., Sungnam, Korea) was deployed along the left main renal artery, bridging the pseudoaneurysm and covering the parent artery, successfully excluding the pseudoaneurysm as confirmed using the final aortography (Figure 2B).

## OUTCOME AND FOLLOW-UP

A repeated CT scan performed 1 d after the first endovascular procedure revealed a decrease in the size of the pseudoaneurysm arising from the renal artery. However, a large intramural blood collection initially noted on the sagittal image remained (Figure 2C), likely caused by an endoleak. Therefore, we immediately performed a re-endovascular procedure. Although the degree of the pseudoaneurysm



DOI: 10.12998/wjcc.v10.i27.9760 Copyright ©The Author(s) 2022.

**Figure 3 Consecutive computed tomography scans performed 26 d, 45 d, and 7 mo after injury.** A: The computed tomography (CT) scan taken 26 d after injury revealed a decrease in the size and extent of the intramural blood collection (red arrow) compared with the CT scan taken seven days prior (figure in red-border box); B: The CT scan taken 45 d after injury showed that the intramural blood collection markedly decreased in size; C: In the 7-mo CT scan, the intramural hematoma (IMH) had completely disappeared; D: There was no evidence of stent-related complications such as perforation, obstruction, and stent migration.


decreased compared with the previous CT scan, evidence of the pseudoaneurysm was still observed using aortography. Subsequently, an additional 8 mm × 45 mm covered stent was installed through the previously deployed covered stent in the left main renal artery. The proximal portion of the newly covered stent was placed into the abdominal aortic lumen to prevent or resolve the endoleak. As a result of the complete coverage of the orifice of the left renal artery, the pseudoaneurysm was successfully excluded, as confirmed using the final aortography (Figure 2D). Eight days after the second endovascular procedure, a follow-up CT scan revealed that the size and extent of the intramural blood collection decreased compared with the previous scan taken 7 d prior (Figure 3A). The intramural blood collection disappeared 19 d after the second endovascular procedure, and the size of the aortic IMH also decreased, as seen on the follow-up CT scan (Figure 3B). Seven months after the injury, a follow-up CT scan showed that the IMH had completely disappeared (Figure 3C). There was no evidence of stent-related complications such as perforation, obstruction, or stent migration (Figure 3D).

## DISCUSSION

There are few reports of small intimomedial tears in IMH cases identified prospectively during imaging or surgery[3,4,8]. As mentioned in the introduction, IMH is classically considered secondary to the rupture of the vasa vasorum. However, with improved technology in the last two decades, more intimomedial tears are being identified in patients with IMH. Therefore, a better distinction may be that AD contains two intimomedial tears: an entry tear from the lumen into the media and a re-entry tear back into the aortic lumen. Conversely, IMH with an intimomedial tear often only has an entry tear[1,7]. In the present case, an intimal injury of the renal artery from the aorta occurred by blunt trauma, leading to the formation of a small orifice in the intimal-medial flap. The focal out-pouching saccular aneurysm arose from the proximal portion of the left renal artery, a pseudoaneurysm, associated with an aortic IMH.

The natural history of IMH is variable. It may resolve, enlarge, or progress to an aneurysm or dissection[9]. Multiple imaging prognostic factors indicating a higher risk of complications or progression have been identified[1,7]. When imaging an IMH, the following items should be considered: Maximum aortic diameter, maximum hematoma thickness, lack of focal contrast enhancement (*i.e.*,





DOI: 10.12998/wjcc.v10.i27.9760 Copyright ©The Author(s) 2022.

**Figure 4** Two-dimensional illustration of the treatment process for aortic intramural hematoma associated with a pseudoaneurysm arising from the renal artery after blunt trauma using covered stents. A: The intimomedial tear (yellow circle) of the proximal renal artery occurs first by trauma, leading to intramural blood collection, and finally within the intramural hematoma (yellow asterisk). This intramural blood collection is defined as a pseudoaneurysm (yellow arrow); B: Although the size of the pseudoaneurysm appeared to decrease after the deployment of the stent graft, a large blood pool was still present due to the endoleak caused by the residual blood flow (orange angled arrow) through the space between the uncovered area of the stent graft (yellow arrowhead) and intimomedial flap; C: The intramural blood collection (yellow circle) was markedly decreased in size and extent after installing an additional stent graft (yellow arrowhead); D: The intramural blood collection and hematoma completely disappeared seven months after the second session.

intramural blood collection or ulcer-like projection), and Stanford classification[1,2,5,9]. An intramural blood collection is a similar small localized area of contrast enhancement with the IMH and is more likely to occur in the descending aorta[9]. An intramural blood collection may appear as a string of contrast agents pooling on the coronal and sagittal images[1,2]. Continued improvements in the thin-section CT may allow for increased visibility of the connection to the branch artery and the aortic lumen [1,2,8]. Although studies of intramural blood collection are limited and the prognostic significance is uncertain, patients with a large intramural blood collection and those with a visible connection to a branch artery are at a higher risk of incomplete hematoma resorption. The blood collection may increase over time, necessitating endovascular treatment. In the present case, the multidetector CT differentiated the pseudoaneurysm as being confined within the IMH and not having a wall of its own. The pseudoaneurysm originated from the renal artery, extended along type A IMH, and gradually increased in size over time, as seen in the follow-up CT scans. This situation requires emergent surgical or endovascular treatment of the injured proximal renal artery.

Main renal artery injury after blunt trauma is rare, with an estimated incidence of only 0.05%[10-12]. A wide spectrum of treatment plans has been proposed for traumatic renal arterial injury, including observation, endovascular treatment, and surgical procedures. The preferred management is decided based on the hemodynamic stability or the need to control an acute and heavy hemorrhage. In the present case, the clinicians decided first to manage the pseudoaneurysm from the renal artery conservatively, considering the relatively small size of the pseudoaneurysm and the stable hemodynamics of the patient. However, a follow-up CT scan showed an increase in the size and extent of the pseudoaneurysm arising from the proximal renal artery, necessitating immediate endovascular intervention. The endovascular options may play an important role in diagnosing and treating renal artery injury with several advantages such as rapid recovery, a short length of hospital stays, and early resumption of physical activities[11]. Rundback *et al*[13] proposed an angiographic classification system by which renal arterial injuries could be divided into three main types for treatment. Type I lesions, as described in the present case, are saccular and arise from either the main renal artery trunk or proximally from a large segment artery. These lesions can be successfully treated with stent grafts, excluding the pseudoan-

eurysm arising from the main renal artery, while preserving the distal perfusion of the normal parenchyma to prevent kidney dysfunction. Several studies have reported endovascular stent-graft placement with promising results[10-12]. However, the endovascular stent-graft option also has some limitations depending on the available devices (such as stents), and we experienced them in the present case. After the stent-graft deployment along the main renal artery on the first attempt, the aortography seemed to show the proximal and distal parts of the pseudoaneurysm completely. However, a large intramural blood collection was still identified on the follow-up CT scan 1 d later.

Eventually, the pseudoaneurysm was managed after installing an additional stent graft with the proximal part placed in the middle of the aorta lumen to cover the injured area of the renal artery sufficiently (Figure 4). At the time of the urgent procedures, the only type of covered stent available in Korea was the expanded polytetrafluoroethylene-covered stent graft (Hercules: S&G Biotech Inc., Sungnam, Korea), which has an uncovered area of 2.5 mm at each end. Therefore, the injured area of the proximal renal artery may not have been fully covered by the stent graft in the first procedure, resulting in the type 1 endoleak. Considering that the pseudoaneurysm had small communication through an intimomedial tear of only 1-2 mm in diameter and that the tear origin of the renal artery was located too close to the aorta, the stent graft should have been safely placed as far as possible into the aorta beyond the intimomedial flap to prevent the endoleak, rather than adequately covering it.

## CONCLUSION

Understanding IMH pathogenesis, imaging findings, and increasing evidence suggesting that many IMHs are associated with intimomedial tears will help clinicians identify patients with IMHs and select the appropriate treatment, thus improving outcomes. Endovascular treatment, including stent-graft placement, can be an effective and safe treatment strategy for IMHs associated with intimomedial tears. A careful and repetitive approach, based on awareness of the various possible causes of endoleaks, could increase the success rate of endovascular treatment.

## FOOTNOTES

**Author contributions:** Lee JY and Kim Y contributed to manuscript drafting and reviewed the literature; Kim Y, Lee JS, Ye JB, Kim SH, Sul YH, Yoon SY, Choi JH, and Choi H collected clinical data and participated in treatment; all authors read and approved the final manuscript.

**Informed consent statement:** The institutional review board of Chungbuk National University Hospital approved a retrospective review of the medical records (2020-12-007) and waived the requirement to obtain informed consent from the patient.

**Conflict-of-interest statement:** The authors declare that they have no conflict of interest.

**CARE Checklist (2016) statement:** The authors have read the CARE Checklist (2016), and the manuscript was prepared and revised according to the CARE Checklist (2016).

**Open-Access:** This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

**Country/Territory of origin:** South Korea

**ORCID number:** Yook Kim 0000-0003-2162-419X; Jin Young Lee 0000-0002-4629-0934; Jin Suk Lee 0000-0002-6694-4683; Jin-Bong Ye 0000-0002-5597-4502; Se Heon Kim 0000-0002-1864-4096; Young Hoon Sul 0000-0003-3184-3396; Su Young Yoon 0000-0001-9264-1371; Jung Hee Choi 0000-0002-4142-6505; Hanlim Choi 0000-0003-0130-3292.

**S-Editor:** Chen YL

**L-Editor:** A

**P-Editor:** Chen YL

## REFERENCES

- 1 Chao CP, Walker TG, Kalva SP. Natural history and CT appearances of aortic intramural hematoma. *Radiographics* 2009; 29: 791-804 [PMID: 19448116 DOI: 10.1148/rq.293085122]

- 2 **Gutschow SE**, Walker CM, Martínez-Jiménez S, Rosado-de-Christenson ML, Stowell J, Kunin JR. Emerging Concepts in Intramural Hematoma Imaging. *Radiographics* 2016; **36**: 660-674 [PMID: [27163587](#) DOI: [10.1148/rg.2016150094](#)]
- 3 **Seitun S**, Rossi UG, Cademartiri F, Maffei E, Cronin P, Ferro C, Williams DM. MDCT findings of aortic branch artery pseudoaneurysms associated with type B intramural haematoma. *Radiol Med* 2012; **117**: 789-803 [PMID: [22228134](#) DOI: [10.1007/s11547-011-0779-1](#)]
- 4 **Williams DM**, Cronin P, Dasika N, Upchurch GR Jr, Patel HJ, Deeb MG, Abrams G. Aortic branch artery pseudoaneurysms accompanying aortic dissection. Part I. Pseudoaneurysm anatomy. *J Vasc Interv Radiol* 2006; **17**: 765-771 [PMID: [16687741](#) DOI: [10.1097/01.RVI.0000209217.94558.81](#)]
- 5 **Nienaber CA**, Powell JT. Management of acute aortic syndromes. *Eur Heart J* 2012; **33**: 26-35b [PMID: [21810861](#) DOI: [10.1093/eurheartj/ehs186](#)]
- 6 **Park KH**, Lim C, Choi JH, Sung K, Kim K, Lee YT, Park PW. Prevalence of aortic intimal defect in surgically treated acute type A intramural hematoma. *Ann Thorac Surg* 2008; **86**: 1494-1500 [PMID: [19049737](#) DOI: [10.1016/j.athoracsur.2008.06.061](#)]
- 7 **Sawaki S**, Hirate Y, Ashida S, Takanohashi A, Yagami K, Usui M. Clinical outcomes of medical treatment of acute type A intramural hematoma. *Asian Cardiovasc Thorac Ann* 2010; **18**: 354-359 [PMID: [20719786](#) DOI: [10.1177/0218492310375855](#)]
- 8 **Bosma MS**, Quint LE, Williams DM, Patel HJ, Jiang Q, Myles JD. Ulcerlike projections developing in noncommunicating aortic dissections: CT findings and natural history. *AJR Am J Roentgenol* 2009; **193**: 895-905 [PMID: [19696307](#) DOI: [10.2214/AJR.08.2073](#)]
- 9 **Kaji S**, Nishigami K, Akasaka T, Hozumi T, Takagi T, Kawamoto T, Okura H, Shono H, Horibata Y, Honda T, Yoshida K. Prediction of progression or regression of type A aortic intramural hematoma by computed tomography. *Circulation* 1999; **100**: II281-II286 [PMID: [10567317](#) DOI: [10.1161/01.cir.100.suppl\\_2.ii-281](#)]
- 10 **Lim KH**, Ryeom HK, Park J. Endovascular treatment of renal arterial perforation after blunt trauma: Case report. *Int J Surg Case Rep* 2018; **42**: 208-211 [PMID: [29275235](#) DOI: [10.1016/j.ijscr.2017.11.069](#)]
- 11 **Loffroy R**, Chevallier O, Gehin S, Midulla M, Berthod PE, Galland C, Briche P, Duperron C, Majbri N, Mousson C, Falvo N. Endovascular management of arterial injuries after blunt or iatrogenic renal trauma. *Quant Imaging Med Surg* 2017; **7**: 434-442 [PMID: [28932700](#) DOI: [10.21037/qims.2017.08.04](#)]
- 12 **Lopera JE**, Suri R, Kroma G, Gadani S, Dolmatch B. Traumatic occlusion and dissection of the main renal artery: endovascular treatment. *J Vasc Interv Radiol* 2011; **22**: 1570-1574 [PMID: [21937239](#) DOI: [10.1016/j.jvir.2011.08.002](#)]
- 13 **Rundback JH**, Rizvi A, Rozenblit GN, Poplauskys M, Maddineni S, Crea G, Agrawal U, Olson C, Matalon TA. Percutaneous stent-graft management of renal artery aneurysms. *J Vasc Interv Radiol* 2000; **11**: 1189-1193 [PMID: [11041477](#) DOI: [10.1016/s1051-0443\(07\)61362-1](#)]



Published by **Baishideng Publishing Group Inc**  
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

**Telephone:** +1-925-3991568

**E-mail:** [bpgoffice@wjgnet.com](mailto:bpgoffice@wjgnet.com)

**Help Desk:** <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

