

World Journal of *Clinical Cases*

World J Clin Cases 2022 December 6; 10(34): 12462-12803



FIELD OF VISION

- 12462 Problematics of neurosurgical service during the COVID-19 pandemic in Slovenia
Munda M, Bosnjak R, Velnar T

MINIREVIEWS

- 12470 Circulating angiotensin converting enzyme 2 and COVID-19
Leowattana W, Leowattana T, Leowattana P
- 12484 Evaluation of gut dysbiosis using serum and fecal bile acid profiles
Monma T, Iwamoto J, Ueda H, Tamamushi M, Kakizaki F, Konishi N, Yara S, Miyazaki T, Hirayama T, Ikegami T, Honda A
- 12494 Pediatric kidney transplantation during the COVID-19 pandemic
Tamura H

ORIGINAL ARTICLE**Clinical and Translational Research**

- 12500 *Coptis*, *Pinellia*, and *Scutellaria* as a promising new drug combination for treatment of *Helicobacter pylori* infection
Yu Z, Sheng WD, Yin X, Bin Y

Case Control Study

- 12515 Effects of illness perception on negative emotions and fatigue in chronic rheumatic diseases: Rumination as a possible mediator
Lu Y, Jin X, Feng LW, Tang C, Neo M, Ho RC

Retrospective Study

- 12532 Significance of incidental focal fluorine-18 fluorodeoxyglucose uptake in colon/rectum, thyroid, and prostate: With a brief literature review
Lee H, Hwang KH
- 12543 Follow-up study on ThinPrep cytology test-positive patients in tropical regions
Chen YC, Liang CN, Wang XF, Wang MF, Huang XN, Hu JD
- 12551 Effect of teach-back health education combined with structured psychological nursing on adverse emotion and patient cooperation during ^{99m}Tc -3PRGD2.SPECT/CT
Gong WN, Zhang YH, Niu J, Li XB
- 12559 Nosocomial infection and spread of SARS-CoV-2 infection among hospital staff, patients and caregivers
Cheng CC, Fann LY, Chou YC, Liu CC, Hu HY, Chu D

Observational Study

- 12566** Effectiveness and safety of generic and brand direct acting antivirals for treatment of chronic hepatitis C
Abdulla M, Al Ghareeb AM, Husain HAHY, Mohammed N, Al Qamish J
- 12578** Influence of group B streptococcus and vaginal cleanliness on the vaginal microbiome of pregnant women
Liao Q, Zhang XF, Mi X, Jin F, Sun HM, Wang QX

Randomized Controlled Trial

- 12587** Clinical study on tri-tongue acupuncture combined with low-frequency electrical stimulation for treating post-stroke dysarthria
Man B, Li WW, Xu JF, Wang Q

META-ANALYSIS

- 12594** Three-dimensional time-of-flight magnetic resonance angiography combined with high resolution T2-weighted imaging in preoperative evaluation of microvascular decompression
Liang C, Yang L, Zhang BB, Guo SW, Li RC

CASE REPORT

- 12605** Acute cytomegalovirus hepatitis in an immunocompetent patient: A case report
Wang JP, Lin BZ, Lin CL, Chen KY, Lin TJ
- 12610** Long-term results of extended Boari flap technique for management of complete ureteral avulsion: A case report
Zhong MZ, Huang WN, Huang GX, Zhang EP, Gan L
- 12617** Amyloid β -related angiitis of the central nervous system occurring after COVID-19 vaccination: A case report
Kizawa M, Iwasaki Y
- 12623** Pseudoileus caused by primary visceral myopathy in a Han Chinese patient with a rare MYH11 mutation: A case report
Li N, Song YM, Zhang XD, Zhao XS, He XY, Yu LF, Zou DW
- 12631** Emergent use of tube tip in pharynx technique in "cannot intubate cannot oxygenate" situation: A case report
Lin TC, Lai YW, Wu SH
- 12637** Inflammatory myofibroblastic tumor of the central nervous system: A case report
Su ZJ, Guo ZS, Wan HT, Hong XY
- 12648** Atypical aggressive vertebral hemangioma of the sacrum with postoperative recurrence: A case report
Wang GX, Chen YQ, Wang Y, Gao CP
- 12654** Closed reduction of hip dislocation associated with ipsilateral lower extremity fractures: A case report and review of the literature
Xu Y, Lv M, Yu SQ, Liu GP

- 12665** Repair of a large patellar cartilage defect using human umbilical cord blood-derived mesenchymal stem cells: A case report
Song JS, Hong KT, Song KJ, Kim SJ
- 12671** Abdominal bronchogenic cyst: A rare case report
Li C, Zhang XW, Zhao CA, Liu M
- 12678** Malignant fibrous histiocytoma of the axilla with breast cancer: A case report
Gao N, Yang AQ, Xu HR, Li L
- 12684** Rapid hemostasis of the residual inguinal access sites during endovascular procedures: A case report
Kim H, Lee K, Cho S, Joh JH
- 12690** Formation of granulation tissue on bilateral vocal cords after double-lumen endotracheal intubation: A case report
Xiong XJ, Wang L, Li T
- 12696** Giant cellular leiomyoma in the broad ligament of the uterus: A case report
Yan J, Li Y, Long XY, Li DC, Li SJ
- 12703** Pomolidomide for relapsed/refractory light chain amyloidosis after resistance to both bortezomib and daratumumab: A case report
Li X, Pan XH, Fang Q, Liang Y
- 12711** Ureteral- artificial iliac artery fistula: A case report
Feng T, Zhao X, Zhu L, Chen W, Gao YL, Wei JL
- 12717** How to manage isolated tension non-surgical pneumoperitonium during bronchoscopy? A case report
Baima YJ, Shi DD, Shi XY, Yang L, Zhang YT, Xiao BS, Wang HY, He HY
- 12726** Amiodarone-induced muscle tremor in an elderly patient: A case report
Zhu XY, Tang XH, Yu H
- 12734** Surgical treatment of Pitt-Hopkins syndrome associated with strabismus and early-onset myopia: Two case reports
Huang Y, Di Y, Zhang XX, Li XY, Fang WY, Qiao T
- 12742** Massive low-grade myxoid liposarcoma of the floor of the mouth: A case report and review of literature
Kugimoto T, Yamagata Y, Ohsako T, Hirai H, Nishii N, Kayamori K, Ikeda T, Harada H
- 12750** Gingival enlargement induced by cyclosporine in Medullary aplasia: A case report
Victory Rodríguez G, Ruiz Gutiérrez ADC, Gómez Sandoval JR, Lomeli Martínez SM
- 12761** Compound heterozygous mutations in PMFBP1 cause acephalic spermatozoa syndrome: A case report
Deng TQ, Xie YL, Pu JB, Xuan J, Li XM
- 12768** Colonic tubular duplication combined with congenital megacolon: A case report
Zhang ZM, Kong S, Gao XX, Jia XH, Zheng CN

- 12775** Perforated duodenal ulcer secondary to deferasirox use in a child successfully managed with laparoscopic drainage: A case report
Alshehri A, Alsinan TA
- 12781** Complication after nipple-areolar complex tattooing performed by a non-medical person: A case report
Byeon JY, Kim TH, Choi HJ
- 12787** Interventional urethral balloon dilatation before endoscopic visual internal urethrotomy for post-traumatic bulbous urethral stricture: A case report
Ha JY, Lee MS
- 12793** Regression of gastric endoscopic submucosal dissection induced polypoid nodular scar after *Helicobacter pylori* eradication: A case report
Jin BC, Ahn AR, Kim SH, Seo SY
- 12799** Congenital absence of the right coronary artery: A case report
Zhu XY, Tang XH

ABOUT COVER

Editorial Board Member of *World Journal of Clinical Cases*, Giuseppe Lanza, MD, MSc, PhD, Associate Professor, Department of Surgery and Medical-Surgical Specialties, University of Catania, Catania 95123, Italy. glanza@oasi.en.it

AIMS AND SCOPE

The primary aim of *World Journal of Clinical Cases* (*WJCC*, *World J Clin Cases*) is to provide scholars and readers from various fields of clinical medicine with a platform to publish high-quality clinical research articles and communicate their research findings online.

WJCC mainly publishes articles reporting research results and findings obtained in the field of clinical medicine and covering a wide range of topics, including case control studies, retrospective cohort studies, retrospective studies, clinical trials studies, observational studies, prospective studies, randomized controlled trials, randomized clinical trials, systematic reviews, meta-analysis, and case reports.

INDEXING/ABSTRACTING

The *WJCC* is now abstracted and indexed in Science Citation Index Expanded (SCIE, also known as SciSearch®), Journal Citation Reports/Science Edition, Current Contents®/Clinical Medicine, PubMed, PubMed Central, Scopus, Reference Citation Analysis, China National Knowledge Infrastructure, China Science and Technology Journal Database, and Superstar Journals Database. The 2022 Edition of Journal Citation Reports® cites the 2021 impact factor (IF) for *WJCC* as 1.534; IF without journal self cites: 1.491; 5-year IF: 1.599; Journal Citation Indicator: 0.28; Ranking: 135 among 172 journals in medicine, general and internal; and Quartile category: Q4. The *WJCC*'s CiteScore for 2021 is 1.2 and Scopus CiteScore rank 2021: General Medicine is 443/826.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: *Si Zhao*; Production Department Director: *Xu Guo*; Editorial Office Director: *Jin-Lei Wang*.

NAME OF JOURNAL

World Journal of Clinical Cases

ISSN

ISSN 2307-8960 (online)

LAUNCH DATE

April 16, 2013

FREQUENCY

Thrice Monthly

EDITORS-IN-CHIEF

Bao-Gan Peng, Jerzy Tadeusz Chudek, George Kontogeorgos, Maurizio Serati, Ja Hyeon Ku

EDITORIAL BOARD MEMBERS

<https://www.wjgnet.com/2307-8960/editorialboard.htm>

PUBLICATION DATE

December 6, 2022

COPYRIGHT

© 2022 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjgnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>

Closed reduction of hip dislocation associated with ipsilateral lower extremity fractures: A case report and review of the literature

Yong Xu, Ming Lv, Shu-Qiang Yu, Guang-Ping Liu

Specialty type: Orthopedics

Provenance and peer review:

Unsolicited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's scientific quality classification

Grade A (Excellent): 0

Grade B (Very good): B

Grade C (Good): C

Grade D (Fair): 0

Grade E (Poor): 0

P-Reviewer: Muthu S, India;

Zharikov YO, Russia

Received: September 15, 2022

Peer-review started: September 15, 2022

First decision: September 27, 2022

Revised: October 17, 2022

Accepted: November 11, 2022

Article in press: November 11, 2022

Published online: December 6, 2022



Yong Xu, Department of Orthopaedics, Chengdu Second People's Hospital, Chengdu 610000, Sichuan Province, China

Ming Lv, Guang-Ping Liu, Department of Orthopaedics, Zibo Central Hospital, Shandong University, Zibo 255000, Shandong Province, China

Shu-Qiang Yu, Department of Orthopaedics, The Fourth People's Hospital of Zibo, Zibo 255000, Shandong Province, China

Corresponding author: Guang-Ping Liu, MD, PhD, Chief Doctor, Surgeon, Department of Orthopaedics, Zibo Central Hospital, Shandong University, No. 54 Gongqingtuan Road, Zibo 255000, Shandong Province, China. guangpingpp@126.com

Abstract

BACKGROUND

Traumatic hip dislocation usually occurs following high-velocity trauma. It is imperative that the dislocation be reduced in a timely manner, especially in a closed manner, as an orthopedic emergency. However, closed reduction can hardly be achieved in patients who also have ipsilateral lower extremity fractures. Herein, we focus on hip dislocation associated with ipsilateral lower extremity fractures, excluding intracapsular fractures (femoral head and neck fractures), present an early closed hip joint reduction method for this injury pattern, and review the literature to discuss the appropriate closed reduction technique for this rare injury pattern.

CASE SUMMARY

We report a case of a 37-year-old male who sustained a left acetabular posterior wall fracture, an ipsilateral comminuted subtrochanteric fracture and dislocation of the hip. The hip dislocation was reduced urgently in a closed manner using the joy-stick technique with a T-shaped Schanz screw. The fractures were reduced and fixed as a 2nd-stage surgery procedure. At the 17-month postoperative follow-up, the patient had full range of motion of the affected hip.

CONCLUSION

Closed reduction of a hip dislocation associated with ipsilateral lower extremity fractures is rarely achieved by regular maneuvers. Attempts at closed reduction, by means of indirectly controlling the proximal fracture fragment or reconstructing the femoral leverage rapidly with the aid of various external reduction apparatuses, were shown to be effective in some scenarios. Mandatory open

reduction is indicated in cases of failed closed reduction, particularly in irreducible dislocations.

Key Words: Trauma; Hip dislocation; Close reduction; Open reduction; Fracture; Case report

©The Author(s) 2022. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: Hip dislocation associated with ipsilateral lower extremity fractures could be more efficiently managed with the aid of a T-shaped Schanz screw. Furthermore, this is the first review of similar techniques for early closed hip joint reduction for this injury pattern.

Citation: Xu Y, Lv M, Yu SQ, Liu GP. Closed reduction of hip dislocation associated with ipsilateral lower extremity fractures: A case report and review of the literature. *World J Clin Cases* 2022; 10(34): 12654-12664

URL: <https://www.wjgnet.com/2307-8960/full/v10/i34/12654.htm>

DOI: <https://dx.doi.org/10.12998/wjcc.v10.i34.12654>

INTRODUCTION

The hip joint is a ball and socket joint, in which its stability is maintained by the combination of bone and strong soft-tissue structures. Generally, traumatic hip dislocation occurs in young patients after high-energy trauma. The incidence of traumatic hip dislocation accounts for approximately 5% of all traumatic joint dislocations[1]. The common mechanisms of hip dislocation are motor vehicle accidents, falls from a height, motorcycle accidents, sports injuries and so on[2-4]. Due to the significant dislocation force, hip dislocation has a high rate of associated hip district fractures[5] or ipsilateral knee injuries[6,7]. Sahin *et al*[4] reported that 71% of patients diagnosed with hip dislocation had concomitant injuries, either systemic or musculoskeletal.

Traumatic hip dislocation is considered an orthopedic emergency. Urgent reduction should be performed to diminish the incidence and severity of major sciatic nerve injury[3], to reduce the duration of ischemia to the femoral head and minimize the incidence of osteonecrosis[2,8]. The longer the duration of hip dislocation, the higher the risk of osteonecrosis of the femoral head[9]. Generally, the priority is closed reduction as the initial treatment choice. In patients who have hip dislocation associated with ipsilateral lower extremity injuries (especially fractures), closed reduction can hardly be achieved because the concomitant injuries may be exacerbated by closed reduction maneuvers[8]. The early recognition and assessment of concomitant injuries is imperative to optimize the treatment procedures. The paucity of literature on closed reduction of this hip dislocation type poses a challenge for orthopedic surgeons for the reduction of this type of hip dislocation in a timely manner.

CASE PRESENTATION

Chief complaints

A 37-year-old male with hypotension was transferred to our trauma center from a local hospital after a vehicle crashed into the left side of his motorcycle on April 12, 2021. The patient complained of severe pain in the left hip and chest.

History of present illness

The patient had no present illness.

History of past illness

The patient had no past illness.

Personal and family history

The patient had no personal or family history.

Physical examination

The patient's blood pressure was 96/62 mmHg, and his heart rate was 85-110 beats/min with a normal sinus rhythm. Physical examination revealed minor pallor of the conjunctiva and shortening, external rotation and abduction deformities of the left leg. The femoral head was palpable in the gluteal region. Upon neurological examination, he was unable to dorsiflex his left foot and toes, and his sensory touch

was impaired in the calf and foot regions. There was no distal vascular deficit noted.

Laboratory examinations

The laboratory examinations appeared unremarkable, except for leukocytosis (white blood cell count, 15.98×10^9 cells/L) before resuscitation.

Imaging examinations

The computed tomography (CT) scan from a local hospital revealed hydropneumothorax, pulmonary contusions, multiple rib fractures, a left acetabular posterior wall fracture, an ipsilateral comminuted subtrochanteric fracture and dislocation of the hip (Figure 1).

FINAL DIAGNOSIS

The patient was diagnosed with a left hip dislocation fracture associated with an ipsilateral comminuted subtrochanteric fracture and a closed chest injury.

TREATMENT

After advanced trauma life support, the patient was urgently taken to the operating room and positioned in the right lateral decubitus position under deep conscious sedation through propofol. We inserted a T-shaped Schanz screw into the subtrochanteric cortex percutaneously under fluoroscopic guidance and manipulated the femoral head into the acetabulum using the joy-stick technique. The reduction was intraoperatively confirmed by fluoroscopy (Figure 2). Proximal tibial skeletal traction was performed to prevent femoral head redislocation, fracture immobilization and limb length restoration. Then, the patient was transferred to the intensive care unit, and fluid resuscitation and the optimization of physiological status continued. In the interim, to illustrate the acetabular fractures in detail for the development of the definitive surgical strategy, we performed post-reduction CT scans (Figure 3) which showed an acetabular posterior wall fracture, a marginal impaction fracture and an intra-articular loose body.

The 2nd-stage definitive fracture fixation was performed as soon as the patient's physiological status was stable enough. The patient was positioned in the right lateral decubitus position under general anesthesia. First, we explored the sciatic nerve and addressed the acetabular fractures through a Kocher-Langenbeck approach. After removal of the loose body, the impaction fracture was elevated and filled with an autograft. The acetabular posterior wall fracture was reduced and stabilized using 2 Lag screws and a buttress plate. Then, the femoral subtrochanteric fracture was reduced with a closed procedure and fixed with an antegrade intramedullary nail (Figure 4A). Intraoperative fluoroscopy was performed to confirm a concentric reduction, and hip stability was assessed by gently moving and applying force in the direction of the dislocation. Postoperatively, skin traction was applied to the affected leg for 2 wk until the stitches were removed. Then, non-weight-bearing activity with axillary crutches ambulating was allowed for another 4 wk.

OUTCOME AND FOLLOW-UP

Full weight-bearing activity was allowed when the radiographs demonstrated a solid union of the subtrochanteric fracture at 6 mo after surgery (Figure 4B). At the 17-mo postoperative follow-up, the patient had full range motion of the affected hip with residual foot droop. Radiographs revealed no evidence of avascular necrosis of the femoral head (Figure 4C).


DISCUSSION

Early closed reduction is the priority for traumatic hip dislocation since it can often be conveniently manipulated in the emergency room, it can shorten the duration of femoral head ischemia, and it can be carried out in conditions that are not suitable for the definitive management of associated fractures. Moreover, early closed reduction could also promptly relieve any distortion of the nerve from a dislocated femoral head or a displaced acetabular fracture in cases of nerve injury[10]. In contrast to open reduction, closed reduction tends to have better clinical outcomes, possibly because of less disruption of the remaining blood supply to the femoral head[11]. All kinds of closed reduction maneuvers, whether for anterior or posterior dislocation, rely on an intact ipsilateral lower extremity to transmit inline traction force and allow the femur to act as a lever to manipulate the femoral head into



DOI: 10.12998/wjcc.v10.i34.12654 Copyright ©The Author(s) 2022.

Figure 1 Three-dimensional computed tomography from a local hospital. The image shows a left acetabular posterior wall fracture, an ipsilateral comminuted subtrochanteric fracture and dislocation of the hip.



DOI: 10.12998/wjcc.v10.i34.12654 Copyright ©The Author(s) 2022.

Figure 2 Intraoperative fluoroscopy showing reduction of a dislocation. A: Prereduction and T-shaped Schanz screw; B: Postreduction.

the acetabulum. Regardless of the closed reduction maneuver and hip dislocation type, the surgeon needs to place his or her hand (or knee/arm/shoulder/forearm) underneath the ipsilateral knee of the affected hip and needs to apply a longitudinal traction force with internal and external rotation until the hip is reduced[8]. When a hip dislocation is associated with ipsilateral lower extremity fractures, closed reduction can hardly be achieved due to the ineffective traction fulcrum resulting from fractures around the knee or the absence of an intact femur necessary to transmit the traction force. We reviewed similar cases (Table 1) to discuss the appropriate closed reduction technique for this rare trauma pattern to improve the clinical outcomes.

The first reported method of closed reduction for posterior hip dislocation associated with an ipsilateral femoral shaft fracture seems to have been published by Wiltberger in 1948[13]. The attempt at early closed reduction with the aid of two threaded pins (reduction apparatus) placed approximately four inches apart into the lateral aspect of the left femoral trochanter had failed. After 3 d of tibial tubercle traction, the author threaded a two-foot length gas pipe over the crossbar of the reduction

Table 1 Review of reported cases of hip dislocation associated with ipsilateral lower extremity fractures

No.	Ref.	Age (yr)	Gender	Cause of injury	Dislocation type	Associated fractures of ipsilateral lower extremity	Method
1	Henry <i>et al</i> [12], 1934	64	F	-	P	Femoral shaft	Open
2	Wiltberger <i>et al</i> [13], 1948	35	M	Industrial accident	P	Femoral shaft	Closed
3	Ingram <i>et al</i> [14], 1954	17	M	Automobile vehicle accident	P	Femoral shaft	Closed
4	Murray <i>et al</i> [15], 1958	25/18	M/-	Motorcycle accident/-	P	Supracondylar of femur/tibia and supracondylar of femur	Open/closed
5	Helal <i>et al</i> [16], 1967	-	-	-	P	Femoral shaft	Closed
6	Lyddon <i>et al</i> [11], 1971	47	M	Automobile accident	Obturator	Femoral shaft	Closed
7	Ehtisham <i>et al</i> [17], 1976	19/19/; 19/17	-	Road traffic accident/ motorcycle accident (3)	A/P/P/A	Femoral shaft/femoral shaft/femoral shaft and roof of the acetabulum/femur, tibia and medial malleolus	Open
8	Schoenecker <i>et al</i> [18], 1978	18	M	Motor vehicle accident	A	Femoral shaft	Closed
9	Harper <i>et al</i> [19], 1982	17/19	M/F	Automobile accident	P	Femoral shaft	Closed
10	Barquet <i>et al</i> [20], 1983	25	M	Automobile collision	P	Femoral head, trochanter and shaft	Closed ¹
11	Verdonk <i>et al</i> [21], 1984	17/21	M	Motor cycle accident/-	P	Femoral shaft	Closed
12	Shannak[22], 1987	29	M	Automobile accident	Bi (A/P)	Femoral shaft/acetabulum, tibia and fibula	Closed
13	Carlsen <i>et al</i> [23], 1991	42	-	Motorcycle-car accident	P	Femoral shaft	Closed
14	Wu <i>et al</i> [24], 1993	16 cases	-	-	P (11) C (5)	Femoral shaft	Closed
15	Maqsood <i>et al</i> [25], 1996	21	M	Run over by a jeep	P	Femoral shaft	Closed
16	Maini <i>et al</i> [26], 2004	25	M	Railway accident	P	Femoral neck and greater trochanter	Open
17	Duygulu <i>et al</i> [27], 2006	52	M	Motor vehicle collision	P	Transverse and posterior wall acetabular fracture, femoral neck and shaft	Open
18	Sié EJB <i>et al</i> [28], 2006	24	M	Fall from a moving truck	P	Shaft, supra and intercondylar of femur, patella, tibial and medial malleolus	Closed ¹
19	Singh <i>et al</i> [29], 2006	35	M	Roadside accident	Inferior	IT fracture	Closed
20	Singh <i>et al</i> [30], 2008	55	M	Fall from a moving bus	Inferior	Open femoral subtrochanteric fracture	Closed
21	Alexa <i>et al</i> [31], 2009	41	M	Traffic accident	P	IT fracture	Open
22	Almosalamy <i>et al</i> [32], 2010	28	M	Car accident	P	Posterior wall acetabular and IT fracture	Open
23	Rodriguez-Martin <i>et al</i> [33], 2010	27	M	Car accident	P	Femoral head and intertrochanteric fracture	Open
24	Sen <i>et al</i> [34], 2011	32	M	Car accident	P	Femoral head, acetabular wall, knee dislocation and tibial plateau fracture	Open
25	Kuhn <i>et al</i> [35], 2013	44	M	Motor vehicle collision	P	Posterior wall acetabular and femoral per-trochanteric fracture	Closed
26	Radulescu <i>et al</i> [36], 2013	44	M	Precipitation	A	IT fracture	Open
27	Sinha <i>et al</i> [37], 2013	45	M	Fall from moving train	P	Transverse and posterior wall acetabular fracture, IT fracture	Open

28	Yousefi <i>et al</i> [38], 2013	43	M	Motor vehicle accident	P	Posterior wall acetabular and IT fracture	Open
29	Zhen[39], 2013	59	M	Car crash	Inferior	Posterior wall acetabular fracture and IT fracture	Total hip arthroplasty
30	Jamshidi <i>et al</i> [40], 2014	26	M	Motor vehicle accident	P	Posterior wall acetabular and IT fracture, tibia, fibula	Closed ¹
31	Chotai <i>et al</i> [41], 2015	25	M	Motor vehicle accident	P	IT fracture and proximal tibial	Open
32	Panigrahi <i>et al</i> [42], 2015	20	M	Road traffic accident	P	Femoral head, shaft and medial condylar fracture	Open
33	Alhammoud <i>et al</i> [43], 2016	30	M	Motor vehicle collision	P	Femoral head and shaft	Closed
34	Qi <i>et al</i> [44], 2016	43	M	Car accident	P	Transverse and posterior wall acetabular fracture, femoral shaft	Open
35	Ul Haq <i>et al</i> [45], 2016	26/36	M/F	Road traffic accident	P	IT fracture/femoral head and intertrochanteric fracture	Open
36	Uzun <i>et al</i> [46], 2017	20	M	Traffic accident	P	Transverse and posterior wall acetabular fracture, IT and lateral condyle	Open
37	Fageir <i>et al</i> [47], 2018	31	M	Fire truck accident	P	IT fracture	Open
38	Desai <i>et al</i> [48], 2019	19	M	Road traffic accident	P	IT fracture	Open
39	Rana <i>et al</i> [49], 2019	18	M	Road traffic accident	P	Femoral shaft	Closed
40	Benabbouha <i>et al</i> [50], 2020	56	F	Hit by a car	P	Posterior wall acetabular fracture, femoral shaft, tibia and fibula	Closed ¹
41	Gokulprasath <i>et al</i> [51], 2020	26	M	Road traffic accident	A	Subtrochanteric fracture	Closed
42	Iftekhar <i>et al</i> [52], 2020	24	M	Motor vehicle accident	P	Femoral shaft	Closed
43	Su <i>et al</i> [53], 2020	38/29	F/M	Train crash/traffic accident	P	Acetabulum, IT and neck fracture/IT and neck fracture	Open
44	Anand <i>et al</i> [54], 2021	50	M	Road traffic accident	P	Transverse-posterior acetabular fracture, IT and shaft, tibia, fibula	Open
45	Present case (2022)	37	M	Vehicle accident	P	Posterior wall acetabular fracture, and subtrochanteric fracture	Closed

¹The closed reduction maneuver was not mentioned in the literature.

IT: Intertrochanteric; A: Anterior; P: Posterior; C: Central; -: Not mentioned.

apparatus and successfully reduced the dislocation eventually. Although the dislocation was managed in a closed manner, delayed reduction increased the risk of femoral head necrosis. In clinical practice, early open reduction should be performed as an alternative once the attempt at closed reduction fails.

Ingram *et al*[14] introduced another closed reduction method in 1954. This seems to be the first description of an early closed reduction for hip dislocation with an ipsilateral femoral shaft fracture. He inserted a large Steinmann pin through the greater trochanter in an anteroposterior direction and clamped the Steinmann pin with large vice-grip pliers anteriorly and posteriorly at the skin edge. Then, closed reduction was accomplished by strong manual traction. In 1958, Murray DS[15] also successfully reduced hip dislocation by this method. This technique is effective, however, there is a high risk of puncturing the sciatic nerve behind the greater trochanter while the pin transfixes the greater trochanter percutaneously because of the nonanatomic position of the greater trochanter. In 1982, Harper *et al*[19] inserted the Steinmann pin in a posterior-anterior direction, taking care to stay lateral to the sciatic nerve, and successfully treated two patients with dislocations.

Some surgeons advocate closed reduction by manipulating the proximal fracture fragment with the aid of various apparatuses, such as Scuderi traction screw[11], Smith traction screw[16], large bone clamp[18], Lardennois hoop[21], tourniquet[23], Hoffmann half pin[24], and temporary external fixator [43]. However, Schoenecker *et al*[18] achieved 2 dislocations by gentle manual traction in a regular manner, ignoring the ipsilateral femoral fracture, which may involve distraction and angulation of the soft tissues at the fracture site and could inevitably jeopardize the neurovascular bundle.




Figure 3 Computed tomography scans. A, B: The images show an acetabular posterior wall fracture associated with a marginal impaction fracture (red arrow) and an intra-articular loose body (yellow arrow).




Figure 4 Left hip radiographs. A: Postoperative; B: 6 mo later; C: 17 mo later.

Among all the reduction apparatuses, the Schanz screw has been the most preferred[25,29,30,35,51]. It is inserted percutaneously under fluoroscopic guidance and is connected with a universal AO chuck or T handle universal chuck to manipulate the proximal fracture fragment to facilitate reduction. In our present case, we also performed closed reduction with the aid of a Schanz screw, although we were unaware of these previous reports at that time. The Schanz screw we used was T-shaped and was whole without any connector, which is often applied to pull the femoral head laterally in our reduction of acetabular fractures. Due to eliminating the potential loosening between the screw and the connector, the T-shaped Schanz screw played a role of pulling and levering the dislocated femoral head more easily as a whole and facilitated the joystick maneuver more efficiently to reduce the dislocation.

In 2019, Rana *et al*[49] recommended a novel approach to perform closed manipulation of a hip dislocation with a femoral shaft fracture. He restored the leverage of the fractured femur by temporarily fixing the femur with an external fixator instead of controlling the proximal fracture fragment with an external fixator. In 2020, Iftekhhar *et al*[52] also addressed hip dislocation with the same protocol. A temporary external fixator was applied to reduce and fix the femoral shaft fracture. The femoral lever arm was obtained, and hip reduction was achieved with a closed procedure. This practical technique reconstructs the continuity and leverage of the femur by means of external fixator osteosynthesis to transmit the traction force and manipulate the affected leg.

Although many closed reduction methods have been reported and do truly work, they are not always successful[54]. Open reduction could never be abandoned as an alternative, and its importance cannot be overemphasized. Many factors, such as the buttonholed femoral head through the capsule or abductors, a large interposed intra-articular fragment from the femoral head or acetabular wall, and soft tissue impingement, always contribute to irreducible dislocations[55-57]. Forced closed reduction may exacerbate these concomitant fractures or cause iatrogenic bone and peripheral neurovascular injuries[8, 58]. Multiple attempts at closed reduction may result in further traumatic injury to the chondral surface of the femoral head[3] and increase the risk for iatrogenic femoral neck fracture[59]. In this situation, mandatory open reduction should eventually be performed as an alternative after a failed closed

reduction attempt for the hemodynamically stable patients. Another situation is the presence of nonconcentric joint after reduction, which also needs surgery to eliminate intra-articular osteochondral fragments or suture the labral tear to obtain the reduction adequacy and excellent long-term clinical outcome[60].

Analyzing the reported similar cases (Table 1), the success rate of initial closed reduction has been reduced, although various closed reduction techniques have been reported. This is because surgeons have had to tackle more complex associated fractures rather than simple femoral shaft fractures in recent years. This may result from the increasing incidence and severity of such complex injury patterns due to increased high-velocity trauma, especially road traffic accidents. On the other hand, experienced trauma teams and modern resuscitation equipment have allowed an increasing number of patients with associated serious complex fractures to survive[61]. Another tendency was the increased prevalence of associated acetabular and per-trochanteric fractures. In such situations, immediate closed reduction has also been favored over open reduction, especially if the patient's general condition is too unstable for open procedures or if the patient's acetabular fractures need no surgical intervention. For patients who require a prolonged transfer to receive definitive surgery, management should also include immediate closed reduction as the primary procedure of the staged treatment strategy at the local hospital. Although hip dislocation associated with an ipsilateral femoral fracture or knee injury was considered a contraindication for closed reduction in some literature[3,8,59], attempts at closed reduction could also be performed with the aid of various external reduction apparatuses. The rational mechanism of effective indirect reduction techniques is controlling the proximal fracture fragment more easily or reconstructing the femoral leverage rapidly.

CONCLUSION

Closed reduction of a hip dislocation associated with ipsilateral lower extremity fractures is rarely achieved by regular maneuvers. Attempts at closed reduction, by means of indirectly controlling the proximal fracture fragment or reconstructing the femoral leverage rapidly with the aid of various external reduction apparatuses, were shown to be effective in some scenarios. Although closed reduction tends to have better clinical outcomes, mandatory open reduction is indicated in cases of failed closed reduction and particularly irreducible dislocations.

FOOTNOTES

Author contributions: Liu GP and Lv M conceived and formulated the surgery protocol; Xu Y and Yu SQ assembled previous reported records and drafted the manuscript; Liu GP performed the surgery and revised the manuscript for important intellectual content; and all authors read and approved the final manuscript.

Informed consent statement: Informed written consent was obtained from the patient.

Conflict-of-interest statement: The authors declare that they have no conflict of interest.

CARE Checklist (2016) statement: The authors have read the CARE Checklist (2016), and the manuscript was prepared and revised according to the CARE Checklist (2016).

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Country/Territory of origin: China

ORCID number: Yong Xu 0000-0003-2129-4696; Ming Lv 0000-0003-0895-4215; Shu-Qiang Yu 0000-0002-1383-1634; Guang-Ping Liu 0000-0002-4363-8368.

S-Editor: Wang JL

L-Editor: A

P-Editor: Wang JL

REFERENCES

- 1 Brav EA. Traumatic dislocation of the hip: Army experience and results over a twelveyear period. *J Bone Joint Surg* 1962;

- 44: 1115-1134 [DOI: [10.2106/00004623-196244060-00007](https://doi.org/10.2106/00004623-196244060-00007)]
- 2 **Fouk DM**, Mullis BH. Hip dislocation: evaluation and management. *J Am Acad Orthop Surg* 2010; **18**: 199-209 [PMID: [20357229](https://pubmed.ncbi.nlm.nih.gov/20357229/) DOI: [10.5435/00124635-201004000-00003](https://doi.org/10.5435/00124635-201004000-00003)]
 - 3 **Hillyard RF**, Fox J. Sciatic nerve injuries associated with traumatic posterior hip dislocations. *Am J Emerg Med* 2003; **21**: 545-548 [PMID: [14655233](https://pubmed.ncbi.nlm.nih.gov/14655233/) DOI: [10.1016/j.ajem.2003.08.014](https://doi.org/10.1016/j.ajem.2003.08.014)]
 - 4 **Sahin V**, Karakaş ES, Aksu S, Atlihan D, Turk CY, Halici M. Traumatic dislocation and fracture-dislocation of the hip: a long-term follow-up study. *J Trauma* 2003; **54**: 520-529 [PMID: [12634533](https://pubmed.ncbi.nlm.nih.gov/12634533/) DOI: [10.1097/01.TA.0000020394.32496.52](https://doi.org/10.1097/01.TA.0000020394.32496.52)]
 - 5 **Pape HC**, Rice J, Wolfram K, Gänsslen A, Pohlemann T, Krettek C. Hip dislocation in patients with multiple injuries. A followup investigation. *Clin Orthop Relat Res* 2000; 99-105 [PMID: [10943190](https://pubmed.ncbi.nlm.nih.gov/10943190/) DOI: [10.1097/00003086-200008000-00014](https://doi.org/10.1097/00003086-200008000-00014)]
 - 6 **Schmidt GL**, Sciulli R, Altman GT. Knee injury in patients experiencing a high-energy traumatic ipsilateral hip dislocation. *J Bone Joint Surg Am* 2005; **87**: 1200-1204 [PMID: [15930527](https://pubmed.ncbi.nlm.nih.gov/15930527/) DOI: [10.2106/JBJS.D.02306](https://doi.org/10.2106/JBJS.D.02306)]
 - 7 **Tabuenca J**, Truan JR. Knee injuries in traumatic hip dislocation. *Clin Orthop Relat Res* 2000; 78-83 [PMID: [10943187](https://pubmed.ncbi.nlm.nih.gov/10943187/) DOI: [10.1097/00003086-200008000-00011](https://doi.org/10.1097/00003086-200008000-00011)]
 - 8 **Dawson-Amoah K**, Raszewski J, Duplantier N, Waddell BS. Dislocation of the Hip: A Review of Types, Causes, and Treatment. *Ochsner J* 2018; **18**: 242-252 [PMID: [30275789](https://pubmed.ncbi.nlm.nih.gov/30275789/) DOI: [10.31486/toj.17.0079](https://doi.org/10.31486/toj.17.0079)]
 - 9 **Ahmed G**, Shiraz S, Riaz M, Ibrahim T. Late versus early reduction in traumatic hip dislocations: a meta-analysis. *Eur J Orthop Surg Traumatol* 2017; **27**: 1109-1116 [PMID: [28578459](https://pubmed.ncbi.nlm.nih.gov/28578459/) DOI: [10.1007/s00590-017-1988-7](https://doi.org/10.1007/s00590-017-1988-7)]
 - 10 **Cornwall R**, Radomisl TE. Nerve injury in traumatic dislocation of the hip. *Clin Orthop Relat Res* 2000; 84-91 [PMID: [10943188](https://pubmed.ncbi.nlm.nih.gov/10943188/) DOI: [10.1097/00003086-200008000-00012](https://doi.org/10.1097/00003086-200008000-00012)]
 - 11 **Lyddon DW Jr**, Hartman JT. Traumatic dislocation of the hip with ipsilateral femoral fractures. A case report. *J Bone Joint Surg Am* 1971; **53**: 1012-1016 [PMID: [5557603](https://pubmed.ncbi.nlm.nih.gov/5557603/) DOI: [10.2106/00004623-197153050-00018](https://doi.org/10.2106/00004623-197153050-00018)]
 - 12 **Henry AK**, Bayumi M. Fracture of the femur with luxation of the ipsilateral hip. *Br J Surg* 1934; **22**: 204-230 [DOI: [10.1002/bjs.1800228603](https://doi.org/10.1002/bjs.1800228603)]
 - 13 **Wiltberger BR**, Mitchell CL, Hedrick DW. Fracture of the femoral shaft complicated by hip dislocation; a method of treatment. *J Bone Joint Surg Am* 1948; **30A**: 225-228 [PMID: [18921642](https://pubmed.ncbi.nlm.nih.gov/18921642/) DOI: [10.2106/00004623-194830010-00024](https://doi.org/10.2106/00004623-194830010-00024)]
 - 14 **Ingram AJ**, Turner TC. Bilateral traumatic posterior dislocation of the hip complicated by bilateral fracture of the femoral shaft: report of a case. *J Bone Joint Surg Am* 1954; **36-A**: 1249-1255 [PMID: [13211718](https://pubmed.ncbi.nlm.nih.gov/13211718/) DOI: [10.2106/00004623-195436060-00014](https://doi.org/10.2106/00004623-195436060-00014)]
 - 15 **MURRAY DS**. Method of reducing fracture-dislocation of hip-joint with fractured femur on same side. *Br Med J* 1958; **2**: 427-428 [PMID: [13560889](https://pubmed.ncbi.nlm.nih.gov/13560889/) DOI: [10.1136/bmj.2.5093.427](https://doi.org/10.1136/bmj.2.5093.427)]
 - 16 **Helal B**, Skevis X. Unrecognised dislocation of the hip in fractures of the femoral shaft. *J Bone Joint Surg Br* 1967; **49**: 293-300 [PMID: [6026515](https://pubmed.ncbi.nlm.nih.gov/6026515/) DOI: [10.1302/0301-620X.49B2.293](https://doi.org/10.1302/0301-620X.49B2.293)]
 - 17 **Ehtisham SM**. Traumatic dislocation of hip joint with fracture of shaft of femur on the same side. *J Trauma* 1976; **16**: 196-205 [PMID: [1255836](https://pubmed.ncbi.nlm.nih.gov/1255836/) DOI: [10.1097/00005373-197603000-00004](https://doi.org/10.1097/00005373-197603000-00004)]
 - 18 **Schoenecker PL**, Manske PR, Sertl GO. Traumatic hip dislocation with ipsilateral femoral shaft fractures. *Clin Orthop Relat Res* 1978; 233-238 [PMID: [639395](https://pubmed.ncbi.nlm.nih.gov/639395/) DOI: [10.1097/00003086-197801000-00025](https://doi.org/10.1097/00003086-197801000-00025)]
 - 19 **Harper MC**. Traumatic dislocation of the hip with ipsilateral femoral shaft fracture: a method of treatment. *Injury* 1982; **13**: 391-394 [PMID: [7085052](https://pubmed.ncbi.nlm.nih.gov/7085052/) DOI: [10.1016/0020-1383\(82\)90091-2](https://doi.org/10.1016/0020-1383(82)90091-2)]
 - 20 **Barquet A**, Mussio A. Fracture-dislocation of the femoral head with associated ipsilateral trochanteric and shaft fracture of the femur. *Arch Orthop Trauma Surg* (1978) 1983; **102**: 61-63 [PMID: [6639314](https://pubmed.ncbi.nlm.nih.gov/6639314/) DOI: [10.1007/BF00443042](https://doi.org/10.1007/BF00443042)]
 - 21 **Verdonk R**, de Smet L. Hip dislocation combined with femoral shaft fracture. Two cases treated with the Lardennois hoop. *Acta Orthop Scand* 1984; **55**: 185-186 [PMID: [6711286](https://pubmed.ncbi.nlm.nih.gov/6711286/) DOI: [10.3109/17453678408992334](https://doi.org/10.3109/17453678408992334)]
 - 22 **Shannak AO**. Bilateral traumatic dislocation of the hips with ipsilateral femoral fracture. *Clin Orthop Relat Res* 1987; 126-129 [PMID: [3802627](https://pubmed.ncbi.nlm.nih.gov/3802627/) DOI: [10.1097/00003086-198702000-00017](https://doi.org/10.1097/00003086-198702000-00017)]
 - 23 **Carlsen AW**, Lind J. Traumatic dislocation of the hip with ipsilateral fracture of the femur: a method of reduction. *Injury* 1991; **22**: 68-69 [PMID: [2030040](https://pubmed.ncbi.nlm.nih.gov/2030040/) DOI: [10.1016/0020-1383\(91\)90170-j](https://doi.org/10.1016/0020-1383(91)90170-j)]
 - 24 **Wu CC**, Shih CH, Chen LH. Femoral shaft fractures complicated by fracture-dislocations of the ipsilateral hip. *J Trauma* 1993; **34**: 70-75 [PMID: [8437199](https://pubmed.ncbi.nlm.nih.gov/8437199/) DOI: [10.1097/00005373-199301000-00013](https://doi.org/10.1097/00005373-199301000-00013)]
 - 25 **Maqsood M**, Walker AP. Asymmetrical bilateral traumatic hip dislocation with ipsilateral fracture of the femoral shaft. *Injury* 1996; **27**: 521-522 [PMID: [8977845](https://pubmed.ncbi.nlm.nih.gov/8977845/) DOI: [10.1016/0020-1383\(96\)00053-8](https://doi.org/10.1016/0020-1383(96)00053-8)]
 - 26 **Maini L**, Mishra P, Jain P, Upadhyay A, Aggrawal A. Three part posterior fracture dislocation of the hip without fracture of the femoral head: review of literature and a case report. *Injury* 2004; **35**: 207-209 [PMID: [14736483](https://pubmed.ncbi.nlm.nih.gov/14736483/) DOI: [10.1016/s0020-1383\(03\)00166-9](https://doi.org/10.1016/s0020-1383(03)00166-9)]
 - 27 **Duygulu F**, Calis M, Argun M, Guney A. Unusual combination of femoral head dislocation associated acetabular fracture with ipsilateral neck and shaft fractures: A case report. *J Trauma* 2006; **61**: 1545-1548 [PMID: [17159705](https://pubmed.ncbi.nlm.nih.gov/17159705/) DOI: [10.1097/01.ta.0000224898.83708.e1](https://doi.org/10.1097/01.ta.0000224898.83708.e1)]
 - 28 **Sié EJB**, Kodo M, Traoré A, Mobiot C, Lambin Y. Floating knee with ipsilateral floating hip injury equivalent variant: A Case report. *Nigerian J Orthop Trauma* 2006; **5**: 50-53
 - 29 **Singh R**, Sharma SC, Goel T. Traumatic inferior hip dislocation in an adult with ipsilateral trochanteric fracture. *J Orthop Trauma* 2006; **20**: 220-222 [PMID: [16648705](https://pubmed.ncbi.nlm.nih.gov/16648705/) DOI: [10.1097/00005131-200603000-00010](https://doi.org/10.1097/00005131-200603000-00010)]
 - 30 **Singh AP**, Singh AP, Mittal V. Traumatic inferior hip dislocation with ipsilateral open subtrochanteric fracture: A rare case. *Injury Extra* 2008; **39**: 384-385 [DOI: [10.1016/j.injury.2008.08.002](https://doi.org/10.1016/j.injury.2008.08.002)]
 - 31 **Alexa O**, Puha B, Veliceasa B, Păduraru D. Posterior dislocation of the hip associated with ipsilateral trochanteric fracture--a very rare case. *Chirurgia (Bucur)* 2009; **104**: 761-764 [PMID: [20187480](https://pubmed.ncbi.nlm.nih.gov/20187480/)]
 - 32 **Almosalamy MH**, Mehrz MH, Al-Mukaimi A. Traumatic posterior hip fracture dislocation with ipsilateral intertrochanteric fracture and sciatic nerve palsy in an adult patient. *J Kuwait Med Assoc* 2010; **42**: 324-327
 - 33 **Rodriguez-Martin J**, Pretell-Mazzini J, Porrás-Moreno MA, Hernanz-Gonzalez Y, Resines-Erasun C. A polytrauma

- patient with an unusual posterior fracture-dislocation of the femoral head: a case report. *Strategies Trauma Limb Reconstr* 2010; **5**: 47-51 [PMID: 20360877 DOI: 10.1007/s11751-009-0078-8]
- 34 **Sen RK**, Tripathy SK, Krishnan V, Goyal T, Jagadeesh V. Ipsilateral fracture dislocations of the hip and knee joints with contralateral open fracture of the leg: a rare case and its management principles. *Chin J Traumatol* 2011; **14**: 183-187 [PMID: 21635808]
- 35 **Kuhn KM**, Boudreau JA, Watson JT. Rare combination of ipsilateral acetabular fracture-dislocation and pertrochanteric fracture. *Am J Orthop (Belle Mead NJ)* 2013; **42**: 372-375 [PMID: 24078956]
- 36 **Radulescu R**, Badila A, Japie I, Papuc A, Manolescu R. Anterior dislocation of the hip associated with intertrochanteric fracture of the femur - Case presentation. *J Med Life* 2013; **6**: 336-339 [PMID: 24146697]
- 37 **Sinha S**, Naik AK, Arya RK, Jain VK. Ipsilateral Traumatic Posterior Hip Dislocation, Posterior Wall and Transverse Acetabular Fracture with Trochanteric Fracture in an adult: Report of First Case. *J Orthop Case Rep* 2013; **3**: 31-34 [PMID: 27298928 DOI: 10.13107/jocr.2250-0685.128]
- 38 **Yousefi A**, Ashraf H, Mashhadinezhad A, Birjandinejad A. Posterior hip dislocation associated with posterior wall acetabular fracture and ipsilateral intertrochanteric fracture: a very rare case report. *Trauma Mon* 2013; **17**: 409-411 [PMID: 24350139 DOI: 10.5812/traumamon.8520]
- 39 **Zhen P**. Traumatic posterior hip dislocation with an ipsilateral comminuted femoral intertrochanteric fracture: a case report. *Orthop Surg* 2013; **5**: 219-221 [PMID: 24002841 DOI: 10.1111/os.12051]
- 40 **Jamshidi MH**, Mirkazemi M, Birjandinejad A. Posterior hip fracture -dislocation associated with ipsilateral intertrochanteric fracture; a rare case report. *Arch Bone Jt Surg* 2014; **2**: 75-78 [PMID: 25207320]
- 41 **Chotai PN**, Ebraheim NA, Hart R, Wassef A. Concomitant Posterior Hip Dislocation, Ipsilateral Intertrochanteric- and Proximal Tibial- Fractures with Popliteal Artery Injury: A Challenging Trauma Mélange. *Clin Pract* 2015; **5**: 776 [PMID: 26918095 DOI: 10.4081/cp.2015.776]
- 42 **Panigrahi R**, Mahapatra AK, Palo N, Priyadarshi A. Neglected Pipkin's fracture dislocation with bilateral femoral shaft fractures: an unusual combination. *J Surg Case Rep* 2015; **2015** [PMID: 25759170 DOI: 10.1093/jscr/rjv013]
- 43 **Alhammoud A**, Alnouri M, Arbash MA, Baco AM. Posterior Hip Dislocation with Ipsilateral Femoral Head and Shaft Fracture - Using a Temporary External Fixator as a Method for Closed Reduction. *J Orthop Case Rep* 2016; **6**: 44-46 [PMID: 28507965]
- 44 **Qi BC**, Zhao Y, Wang CX, Wang TJ, Zhang JT, Ju WN, Sun DH. Posterior dislocation of the hip with bilateral femoral fractures: An unusual combination. *Technol Health Care* 2016; **24**: 281-286 [PMID: 26578280 DOI: 10.3233/THC1111]
- 45 **Ul Haq R**, Kumar J, Dhammi IK, Jain AK. Posterior dislocation of hip with ipsilateral intertrochanteric fracture: A report of two cases. *Indian J Orthop* 2016; **50**: 571-576 [PMID: 27746503 DOI: 10.4103/0019-5413.189614]
- 46 **Uzun E**, Günay AE, Karaman İ, Mısıra A, Kızıkcapan TB, Özçamdallı M. Unusual combination of femoral head dislocation associated acetabular fracture with multipart femoral fractures: Case report. *Int J Surg Case Rep* 2017; **34**: 32-35 [PMID: 28415025 DOI: 10.1016/j.ijscr.2017.02.043]
- 47 **Fageir M**, Veettil MKP. Posterior dislocation of a native hip joint associated with ipsilateral per-trochanteric fracture: A rare case report. *Trauma Case Rep* 2018; **13**: 1-13 [PMID: 29644290 DOI: 10.1016/j.tcr.2017.09.003]
- 48 **Desai KB**, Dhamangaonkar AC. A case of posterior HIP dislocation with intertrochanteric fracture: Beware of the sharp calcar spike. *J Clin Orthop Trauma* 2019; **10**: 975-980 [PMID: 31528079 DOI: 10.1016/j.jcot.2019.04.014]
- 49 **Rana R**, Patra SK, Khuntia S, Jain M, Patro BP. Traumatic Posterior Dislocation of Hip with Ipsilateral Fracture of Shaft of Femur: Temporary Fixator-assisted Reduction and Final Fixation with Interlocking Nail. *Cureus* 2019; **11**: e5488 [PMID: 31656714 DOI: 10.7759/cureus.5488]
- 50 **Benabbouha A**, Boussouga M, Fjouji S, Lamkhanter A, Jaafar A. Simultaneous Ipsilateral Floating Hip and Knee: A Complex Combination and Difficult Surgical Challenge. *Case Rep Orthop* 2020; **2020**: 9197872 [PMID: 32095304 DOI: 10.1155/2020/9197872]
- 51 **Gokulprasath S**, Prashanth KRT. Anterior Hip Dislocation with Ipsilateral Subtrochanteric Femur Fracture in a Young Adult - A Rare Case Report. *J Orthop Case Rep* 2020; **10**: 63-66 [PMID: 33954139]
- 52 **Iftexhar N**, Rasool T, Zhitny VP, Rasool S. Ipsilateral Posterior Hip Dislocation and Femur Shaft Fracture - A Rare Presentation. *J Orthop Case Rep* 2020; **10**: 3-5 [PMID: 33585305]
- 53 **Su C**, Liu Y, Wu P, Yuan J, Lang J, Wu C, Zhang Y, Chen L. Ipsilateral femoral neck and intertrochanteric fractures with posterior dislocation of the hip: A report of two cases. *J Orthop Surg (Hong Kong)* 2020; **28**: 2309499019900449 [PMID: 32336196 DOI: 10.1177/2309499019900449]
- 54 **Anand V**, Deviprasad, Praba K, Babar S, Babu M. Ipsilateral floating hip and knee, or floating lower limb - A rare case report. *Trauma Case Rep* 2021; **31**: 100390 [PMID: 33365379 DOI: 10.1016/j.tcr.2020.100390]
- 55 **Pathinathan K**, Marage N, Fernando N. Bilateral irreducible asymmetrical fracture-dislocation of the hip: A case report and literature review. *Int J Surg Case Rep* 2021; **81**: 105803 [PMID: 33774447 DOI: 10.1016/j.ijscr.2021.105803]
- 56 **Fernandez M**, Williams T, Dubrana F, Di Francia R. A rare case of an irreducible Pipkin II fracture-dislocation of the femoral head in a young patient following low-energy trauma. *Int J Surg Case Rep* 2020; **70**: 234-237 [PMID: 32422586 DOI: 10.1016/j.ijscr.2020.04.005]
- 57 **Pascarella R**, Maresca A, Cappuccio M, Reggiani LM, Boriani S. Asymmetrical bilateral traumatic fracture dislocation of the hip: a report of two cases. *Chir Organi Mov* 2008; **92**: 109-111 [PMID: 18504531 DOI: 10.1007/s12306-008-0045-4]
- 58 **Chen W**, Gao Z, Ma L. Failed reduction of posterior hip dislocation accompanied by femoral head fracture: causes and resolving strategy. *Int Orthop* 2021; **45**: 1609-1614 [PMID: 33108471 DOI: 10.1007/s00264-020-04856-8]
- 59 **Mandell JC**, Marshall RA, Weaver MJ, Harris MB, Sodickson AD, Khurana B. Traumatic Hip Dislocation: What the Orthopedic Surgeon Wants to Know. *Radiographics* 2017; **37**: 2181-2201 [PMID: 29131775 DOI: 10.1148/rg.2017170012]
- 60 **Mitsionis GI**, Lykissas MG, Motsis E, Mitsiou D, Gkiatas I, Xenakis TA, Beris AE. Surgical management of posterior hip dislocations associated with posterior wall acetabular fracture: a study with a minimum follow-up of 15 years. *J Orthop Trauma* 2012; **26**: 460-465 [PMID: 22357088 DOI: 10.1097/BOT.0b013e31822c4d6e]

- 61 **Liebergall M**, Segal D, Peyser A, Mosheiff R. Combined injuries to the lower limbs. *Injury* 1999; **30** Suppl 2: B29-B33 [PMID: 10562858 DOI: 10.1016/s0020-1383(99)90005-0]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA
Telephone: +1-925-3991568
E-mail: bpgoffice@wjgnet.com
Help Desk: <https://www.f6publishing.com/helpdesk>
<https://www.wjgnet.com>

